

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 September 2001 (13.09.2001)

PCT

(10) International Publication Number  
WO 01/66750 A2

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
15/70, 15/81, 15/85, 5/10, 1/21, 1/19, C07K 14/705,  
16/28, C12Q 1/68, G01N 33/68

[US/US]; 3005 First Avenue, Seattle, WA 98121 (US).  
WOOD, Linda, S. [US/US]; 10193 Fox Hollow, Portage,  
MI 49024 (US).

(21) International Application Number: PCT/US01/07322

(74) Agents: DELUCA, Mark et al.; Woodcock Washburn  
Kurtz Mackiewicz & Norris LLP, 46th Floor, One Liberty  
Place, Philadelphia, PA 19103 (US).

(22) International Filing Date: 8 March 2001 (08.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/187,828	8 March 2000 (08.03.2000)	US
60/187,715	8 March 2000 (08.03.2000)	US
60/187,929	8 March 2000 (08.03.2000)	US
60/187,930	8 March 2000 (08.03.2000)	US
60/187,825	8 March 2000 (08.03.2000)	US
60/187,833	8 March 2000 (08.03.2000)	US
60/187,830	8 March 2000 (08.03.2000)	US
60/187,829	8 March 2000 (08.03.2000)	US
60/187,582	8 March 2000 (08.03.2000)	US
60/187,581	8 March 2000 (08.03.2000)	US
60/187,714	8 March 2000 (08.03.2000)	US
60/189,294	8 March 2000 (08.03.2000)	US
60/187,874	8 March 2000 (08.03.2000)	US
60/187,928	8 March 2000 (08.03.2000)	US
60/188,049	8 March 2000 (08.03.2000)	US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a  
patent (Rule 4.17(ii)) for all designations except US
- of inventorship (Rule 4.17(iv)) for US only
- of inventorship (Rule 4.17(iv)) for US only

Published:

- without international search report and to be republished  
upon receipt of that report

(71) Applicant (for all designated States except US): PHAR-  
MACIA & UPJOHN COMPANY [US/US]; 301 Henri-  
etta Street, Kalamazoo, MI 49001 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): VOGELI, Gabriel

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL G PROTEIN-COUPLED RECEPTORS

(57) Abstract: The present invention provides a gene encoding a G protein-coupled receptor termed nGPCR-x; constructs and re-  
combinant host cells incorporating the genes; the nGPCR-x polypeptides encoded by the gene; antibodies to the nGPCR-x polypep-  
tides; and methods of making and using all of the foregoing.

WO 01/66750 A2

## NOVEL G PROTEIN-COUPLED RECEPTORS

### CROSS-REFERENCE TO RELATED APPLICATIONS

5           The present application claims priority of Application Serial No. 60/187,828, filed March 8, 2000; Serial No. 60/187,715, filed March 8, 2000; Serial No. 60/187,929, filed March 8, 2000; Serial No. 60/187,930, filed March 8, 2000; Serial No. 60/187,825, filed March 8, 2000; Serial No. 60/187,833, filed March 8, 2000; Serial No. 60/187,830, filed March 8, 2000; Serial No. 60/187,829, filed March 8, 2000; Serial No. 60/187,582, filed  
10   March 8, 2000; Serial No. 60/187,581, filed March 8, 2000; Serial No. 60/187,714, filed March 8, 2000; Serial No. 60/189,294, filed March 8, 2000; Serial No. 60/187,874, filed March 8, 2000; Serial No. 60/187,928, filed March 8, 2000; Serial No. 60/188,049, filed March 8, 2000, each of which is hereby incorporated by reference in its entirety.

### 15   FIELD OF THE INVENTION

          The present invention relates generally to the fields of genetics and cellular and molecular biology. More particularly, the invention relates to novel G protein coupled receptors, to polynucleotides that encode such novel receptors, to reagents such as antibodies, probes, primers and kits comprising such antibodies, probes, primers related to  
20   the same, and to methods which use the novel G protein coupled receptors, polynucleotides or reagents.

### BACKGROUND OF THE INVENTION

          The G protein-coupled receptors (GPCRs) form a vast superfamily of cell surface  
25   receptors which are characterized by an amino-terminal extracellular domain, a carboxyl-terminal intracellular domain, and a serpentine structure that passes through the cell membrane seven times. Hence, such receptors are sometimes also referred to as seven transmembrane (7TM) receptors. These seven transmembrane domains define three extracellular loops and three intracellular loops, in addition to the amino- and carboxy-  
30   terminal domains. The extracellular portions of the receptor have a role in recognizing

and binding one or more extracellular binding partners (*e.g.*, ligands), whereas the intracellular portions have a role in recognizing and communicating with downstream molecules in the signal transduction cascade.

The G protein-coupled receptors bind a variety of ligands including calcium ions, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and even photons, and are important in the normal (and sometimes the aberrant) function of many cell types. [See generally Strosberg, *Eur. J. Biochem.* 196:1-10 (1991) and Bohm *et al.*, *Biochem J.* 322:1-18 (1997).] When a specific ligand binds to its corresponding receptor, the ligand typically stimulates the receptor to activate a specific heterotrimeric guanine-nucleotide-binding regulatory protein (G-protein) that is coupled to the intracellular portion of the receptor. The G protein in turn transmits a signal to an effector molecule within the cell, by either stimulating or inhibiting the activity of that effector molecule. These effector molecules include adenylate cyclase, phospholipases and ion channels. Adenylate cyclase and phospholipases are enzymes that are involved in the production of the second messenger molecules cAMP, inositol triphosphate and diacylglycerol. It is through this sequence of events that an extracellular ligand stimuli exerts intracellular changes through a G protein-coupled receptor. Each such receptor has its own characteristic primary structure, expression pattern, ligand-binding profile, and intracellular effector system.

Because of the vital role of G protein-coupled receptors in the communication between cells and their environment, such receptors are attractive targets for therapeutic intervention, for example by activating or antagonizing such receptors. For receptors having a known ligand, the identification of agonists or antagonists may be sought specifically to enhance or inhibit the action of the ligand. Some G protein-coupled receptors have roles in disease pathogenesis (*e.g.*, certain chemokine receptors that act as HIV co-receptors may have a role in AIDS pathogenesis), and are attractive targets for therapeutic intervention even in the absence of knowledge of the natural ligand of the receptor. Other receptors are attractive targets for therapeutic intervention by virtue of their expression pattern in tissues or cell types that are themselves attractive targets for therapeutic intervention. Examples of this latter category of receptors include receptors expressed in immune cells, which can be targeted to either inhibit autoimmune responses

or to enhance immune responses to fight pathogens or cancer, and receptors expressed in the brain or other neural organs and tissues, which are likely targets in the treatment of mental disorder, depression, bipolar disease, or other neurological disorders. This latter category of receptor is also useful as a marker for identifying and/or purifying (e.g., via fluorescence-activated cell sorting) cellular subtypes that express the receptor. Unfortunately, only a limited number of G protein receptors from the central nervous system (CNS) are known. Thus, a need exists for G protein-coupled receptors that have been identified and show promise as targets for therapeutic intervention in a variety of animals, including humans.

## SUMMARY OF THE INVENTION

The present invention relates to an isolated nucleic acid molecule that comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, or a fragment thereof. The nucleic acid molecule encodes at least a portion of nGPCR-x. In some embodiments, the nucleic acid molecule comprises a sequence that encodes a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, or a fragment thereof. In some embodiments, the nucleic acid molecule comprises a sequence homologous to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or a fragment thereof. In some embodiments, the nucleic acid molecule comprises a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, and fragments thereof.

According to some embodiments, the present invention provides vectors which comprise the nucleic acid molecule of the invention. In some embodiments, the vector is an expression vector.

According to some embodiments, the present invention provides host cells which comprise the vectors of the invention. In some embodiments, the host cells comprise expression vectors.

The present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence complementary to at least a portion of a sequence selected from the



group consisting of SEQ ID NO:1 to SEQ ID NO:134, said portion comprising at least 10 nucleotides.

The present invention provides a method of producing a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, or a homolog or fragment thereof. The method comprising the steps of introducing a recombinant expression vector that includes a nucleotide sequence that encodes the polypeptide into a compatible host cell, growing the host cell under conditions for expression of the polypeptide and recovering the polypeptide.

The present invention provides an isolated antibody which binds to an epitope on a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, or a homolog or fragment thereof.

The present invention provides a method of inducing an immune response in a mammal against a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, or a homolog or fragment thereof. The method comprises administering to a mammal an amount of the polypeptide sufficient to induce said immune response.

The present invention provides a method for identifying a compound which binds nGPCR-x. The method comprises the steps of contacting nGPCR-x with a compound and determining whether the compound binds nGPCR-x.

The present invention provides a method for identifying a compound which binds a nucleic acid molecule encoding nGPCR-x. The method comprises the steps of contacting said nucleic acid molecule encoding nGPCR-x with a compound and determining whether said compound binds said nucleic acid molecule.

The present invention provides a method for identifying a compound which modulates the activity of nGPCR-x. The method comprises the steps of contacting nGPCR-x with a compound and determining whether nGPCR-x activity has been modulated.

The present invention provides a method of identifying an animal homolog of nGPCR-x. The method comprises the steps screening a nucleic acid database of the animal with a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or a portion thereof and determining whether a portion of said library or database

is homologous to said sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or portion thereof.

The present invention provides a method of identifying an animal homolog of nGPCR-x. The methods comprises the steps screening a nucleic acid library of the animal with a nucleic acid molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or a portion thereof; and determining whether a portion of said library or database is homologous to said sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or a portion thereof.

Another aspect of the present invention relates to methods of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor. The methods comprise the steps of assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR-x that is expressed in the brain. The nGPCR-x comprise an amino acid sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, and allelic variants thereof. A diagnosis of the disorder or predisposition is made from the presence or absence of the mutation. The presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR-x in the nucleic acid correlates with an increased risk of developing the disorder.

The present invention further relates to methods of screening for a nGPCR-x hereditary mental disorder genotype in a human patient. The methods comprise the steps of providing a biological sample comprising nucleic acid from the patient, in which the nucleic acid includes sequences corresponding to alleles of nGPCR-x. The presence of one or more mutations in the nGPCR-x allele is indicative of a hereditary mental disorder genotype.

The present invention provides kits for screening a human subject to diagnose mental disorder or a genetic predisposition therefor. The kits include an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-x gene. The oligonucleotide comprises 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a wild type human nGPCR-x gene sequence or nGPCR-x coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution. The kit also

includes a media packaged with the oligonucleotide. The media contains information for identifying polymorphisms that correlate with mental disorder or a genetic predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

The present invention further relates to methods of identifying nGPCR-x allelic  
5 variants that correlates with mental disorders. The methods comprise the steps of providing biological samples that comprise nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny, and detecting in the nucleic acid the presence of one or more mutations in an nGPCR-x that is expressed in the brain. The nGPCR-x comprises an amino acid sequence selected from the group  
10 consisting of SEQ ID NO:135 to SEQ ID NO:268, and allelic variants thereof. The nucleic acid includes sequences corresponding to the gene or genes encoding nGPCR-x. The one or more mutations detected indicate an allelic variant that correlates with a mental disorder.

The present invention further relates to purified polynucleotides comprising  
15 nucleotide sequences encoding alleles of nGPCR-x from a human with mental disorder. The polynucleotide hybridizes to the complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 under the following hybridization conditions: (a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and (b) washing 2 times for 30  
20 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS. The polynucleotide that encodes nGPCR-x amino acid sequence of the human differs from a sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268 by at least one residue.

The present invention also provides methods for identifying a modulator of  
25 biological activity of nGPCR-x comprising the steps of contacting a cell that expresses nGPCR-x in the presence and in the absence of a putative modulator compound and measuring nGPCR-x biological activity in the cell. The decreased or increased nGPCR-x biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

30 The present invention further provides methods to identify compounds useful for the treatment of mental disorders. The methods comprise the steps of contacting a

composition comprising nGPCR-x with a compound suspected of binding nGPCR-x. The binding between nGPCR-x and the compound suspected of binding nGPCR-x is detected. Compounds identified as binding nGPCR-x are candidate compounds useful for the treatment of mental disorder. Compounds identified as binding nGPCR-x may be further  
5 tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity.

The present invention further provides methods for identifying a compound useful as a modulator of binding between nGPCR-x and a binding partner of nGPCR-x. The methods comprise the steps of contacting the binding partner and a composition  
10 comprising nGPCR-x in the presence and in the absence of a putative modulator compound and detecting binding between the binding partner and nGPCR-x. Decreased or increased binding between the binding partner and nGPCR-x in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a related disease or disorder.  
15 Compounds identified as modulating binding between nGPCR-x and a nGPCR-x binding partner may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity as modulators.

Another aspect of the present invention relates to methods of purifying a G protein from a sample containing a G protein. The methods comprise the steps of contacting the  
20 sample with an nGPCR-x for a time sufficient to allow the G protein to form a complex with the nGPCR-x; isolating the complex from remaining components of the sample; maintaining the complex under conditions which result in dissociation of the G protein from the nGPCR-x; and isolating said G protein from the nGPCR-x.

## 25 DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

### Definitions

Various definitions are made throughout this document. Most words have the meaning that would be attributed to those words by one skilled in the art. Words specifically defined either below or elsewhere in this document have the meaning  
30 provided in the context of the present invention as a whole and as are typically understood by those skilled in the art.

"Synthesized" as used herein and understood in the art, refers to polynucleotides produced by purely chemical, as opposed to enzymatic, methods. "Wholly" synthesized DNA sequences are therefore produced entirely by chemical means, and "partially" synthesized DNAs embrace those wherein only portions of the resulting DNA were  
5 produced by chemical means.

By the term "region" is meant a physically contiguous portion of the primary structure of a biomolecule. In the case of proteins, a region is defined by a contiguous portion of the amino acid sequence of that protein.

The term "domain" is herein defined as referring to a structural part of a  
10 biomolecule that contributes to a known or suspected function of the biomolecule. Domains may be co-extensive with regions or portions thereof; domains may also incorporate a portion of a biomolecule that is distinct from a particular region, in addition to all or part of that region. Examples of GPCR protein domains include, but are not limited to, the extracellular (*i.e.*, N-terminal), transmembrane and cytoplasmic (*i.e.*, C-  
15 terminal) domains, which are co-extensive with like-named regions of GPCRs; each of the seven transmembrane segments of a GPCR; and each of the loop segments (both extracellular and intracellular loops) connecting adjacent transmembrane segments.

As used herein, the term "activity" refers to a variety of measurable indicia suggesting or revealing binding, either direct or indirect; affecting a response, *i.e.* having a  
20 measurable affect in response to some exposure or stimulus, including, for example, the affinity of a compound for directly binding a polypeptide or polynucleotide of the invention, or, for example, measurement of amounts of upstream or downstream proteins or other similar functions after some stimulus or event.

Unless indicated otherwise, as used herein, the abbreviation in lower case (gpcr)  
25 refers to a gene, cDNA, RNA or nucleic acid sequence, while the upper case version (GPCR) refers to a protein, polypeptide, peptide, oligopeptide, or amino acid sequence. The term "nGPCR-x" refers to any of the nGPCRs taught herein, while specific reference to a nGPCR (for example nGPCR-2073) refers only to that specific nGPCR.

As used herein, the term "antibody" is meant to refer to complete, intact  
30 antibodies, and Fab, Fab', F(ab)2, and other fragments thereof. Complete, intact

antibodies include monoclonal antibodies such as murine monoclonal antibodies, chimeric antibodies and humanized antibodies.

As used herein, the term "binding" means the physical or chemical interaction between two proteins or compounds or associated proteins or compounds or combinations thereof. Binding includes ionic, non-ionic, Hydrogen bonds, Van der Waals, hydrophobic interactions, etc. The physical interaction, the binding, can be either direct or indirect, indirect being through or due to the effects of another protein or compound. Direct binding refers to interactions that do not take place through or due to the effect of another protein or compound but instead are without other substantial chemical intermediates. Binding may be detected in many different manners. As a non-limiting example, the physical binding interaction between a nGPCR-x of the invention and a compound can be detected using a labeled compound. Alternatively, functional evidence of binding can be detected using, for example, a cell transfected with and expressing a nGPCR-x of the invention. Binding of the transfected cell to a ligand of the nGPCR-x that was transfected into the cell provides functional evidence of binding. Other methods of detecting binding are well known to those of skill in the art.

As used herein, the term "compound" means any identifiable chemical or molecule, including, but not limited to, small molecule, peptide, protein, sugar, nucleotide, or nucleic acid, and such compound can be natural or synthetic.

As used herein, the term "complementary" refers to Watson-Crick basepairing between nucleotide units of a nucleic acid molecule.

As used herein, the term "contacting" means bringing together, either directly or indirectly, a compound into physical proximity to a polypeptide or polynucleotide of the invention. The polypeptide or polynucleotide can be in any number of buffers, salts, solutions *etc.* Contacting includes, for example, placing the compound into a beaker, microtiter plate, cell culture flask, or a microarray, such as a gene chip, or the like, which contains the nucleic acid molecule, or polypeptide encoding the nGPCR or fragment thereof.

As used herein, the phrase "homologous nucleotide sequence," or "homologous amino acid sequence," or variations thereof, refers to sequences characterized by a homology, at the nucleotide level or amino acid level, of at least the specified percentage.

Homologous nucleotide sequences include those sequences coding for isoforms of proteins. Such isoforms can be expressed in different tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. Homologous nucleotide sequences include nucleotide  
5 sequences encoding for a protein of a species other than humans, including, but not limited to, mammals. Homologous nucleotide sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the nucleotide sequence encoding other known GPCRs. Homologous amino acid sequences include  
10 those amino acid sequences which contain conservative amino acid substitutions and which polypeptides have the same binding and/or activity. A homologous amino acid sequence does not, however, include the amino acid sequence encoding other known GPCRs. Percent homology can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group,  
15 University Research Park, Madison WI), using the default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2, 482-489, which is incorporated herein by reference in its entirety).

As used herein, the term "isolated" nucleic acid molecule refers to a nucleic acid molecule (DNA or RNA) that has been removed from its native environment. Examples  
20 of isolated nucleic acid molecules include, but are not limited to, recombinant DNA molecules contained in a vector, recombinant DNA molecules maintained in a heterologous host cell, partially or substantially purified nucleic acid molecules, and synthetic DNA or RNA molecules.

As used herein, the terms "modulates" or "modifies" means an increase or decrease  
25 in the amount, quality, or effect of a particular activity or protein.

As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues which has a sufficient number of bases to be used in a polymerase chain reaction (PCR). This short sequence is based on (or designed from) a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an identical, similar or  
30 complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 50

nucleotides, preferably about 15 to 30 nucleotides. They are chemically synthesized and may be used as probes.

As used herein, the term "probe" refers to nucleic acid sequences of variable length, preferably between at least about 10 and as many as about 6,000 nucleotides, depending on use. They are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and much slower to hybridize than oligomers. They may be single- or double-stranded and carefully designed to have specificity in PCR, hybridization membrane-based, or ELISA-like technologies.

10 The term "preventing" refers to decreasing the probability that an organism contracts or develops an abnormal condition.

The term "treating" refers to having a therapeutic effect and at least partially alleviating or abrogating an abnormal condition in the organism.

The term "therapeutic effect" refers to the inhibition or activation factors causing or contributing to the abnormal condition. A therapeutic effect relieves to some extent one or more of the symptoms of the abnormal condition. In reference to the treatment of abnormal conditions, a therapeutic effect can refer to one or more of the following: (a) an increase in the proliferation, growth, and/or differentiation of cells; (b) inhibition (*i.e.*, slowing or stopping) of cell death; (c) inhibition of degeneration; (d) relieving to some extent one or more of the symptoms associated with the abnormal condition; and (e) enhancing the function of the affected population of cells. Compounds demonstrating efficacy against abnormal conditions can be identified as described herein.

25 The term "abnormal condition" refers to a function in the cells or tissues of an organism that deviates from their normal functions in that organism. An abnormal condition can relate to cell proliferation, cell differentiation, cell signaling, or cell survival. An abnormal condition may also include obesity, diabetic complications such as retinal degeneration, and irregularities in glucose uptake and metabolism, and fatty acid uptake and metabolism.

30 Abnormal cell proliferative conditions include cancers such as fibrotic and mesangial disorders, abnormal angiogenesis and vasculogenesis, wound healing, psoriasis, diabetes mellitus, and inflammation.



Abnormal differentiation conditions include, but are not limited to, neurodegenerative disorders, slow wound healing rates, and slow tissue grafting healing rates. Abnormal cell signaling conditions include, but are not limited to, psychiatric disorders involving excess neurotransmitter activity.

5 Abnormal cell survival conditions may also relate to conditions in which programmed cell death (apoptosis) pathways are activated or abrogated. A number of protein kinases are associated with the apoptosis pathways. Aberrations in the function of any one of the protein kinases could lead to cell immortality or premature cell death.

The term "administering" relates to a method of incorporating a compound into  
10 cells or tissues of an organism. The abnormal condition can be prevented or treated when the cells or tissues of the organism exist within the organism or outside of the organism. Cells existing outside the organism can be maintained or grown in cell culture dishes. For cells harbored within the organism, many techniques exist in the art to administer compounds, including (but not limited to) oral, parenteral, dermal, injection, and aerosol  
15 applications. For cells outside of the organism, multiple techniques exist in the art to administer the compounds, including (but not limited to) cell microinjection techniques, transformation techniques and carrier techniques.

The abnormal condition can also be prevented or treated by administering a compound to a group of cells having an aberration in a signal transduction pathway to an  
20 organism. The effect of administering a compound on organism function can then be monitored. The organism is preferably a mouse, rat, rabbit, guinea pig or goat, more preferably a monkey or ape, and most preferably a human.

By "amplification" it is meant increased numbers of DNA or RNA in a cell compared with normal cells. "Amplification" as it refers to RNA can be the detectable  
25 presence of RNA in cells, since in some normal cells there is no basal expression of RNA. In other normal cells, a basal level of expression exists, therefore in these cases amplification is the detection of at least 1 to 2-fold, and preferably more, compared to the basal level.

As used herein, the phrase "stringent hybridization conditions" or "stringent  
30 conditions" refers to conditions under which a probe, primer, or oligonucleotide will hybridize to its target sequence, but to no other sequences. Stringent conditions are

sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present in excess, at  $T_m$ , 50% of the probes are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes, primers or oligonucleotides (e.g. 10 to 50 nucleotides) and at least about 60°C for longer probes, primers or oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

The amino acid sequences are presented in the amino to carboxy direction, from left to right. The amino and carboxy groups are not presented in the sequence. The nucleotide sequences are presented by single strand only, in the 5' to 3' direction, from left to right. Nucleotides and amino acids are represented in the manner recommended by the IUPAC-IUB Biochemical Nomenclature Commission or (for amino acids) by three letters code.

## **Polynucleotides**

The present invention provides purified and isolated polynucleotides (e.g., DNA sequences and RNA transcripts, both sense and complementary antisense strands, both single- and double-stranded, including splice variants thereof) that encode unknown G protein-coupled receptors heretofore termed novel GPCRs, or nGPCRs. These genes are described herein and designated herein collectively as nGPCR-x (where x is 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426,

2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, and 74). Table 1 below identifies the novel gene sequence nGPCR-x designation, the SEQ ID NO: of the gene sequence, the SEQ ID NO: of the polypeptide encoded thereby, and the U.S. Provisional Application in which the gene sequence has been disclosed.

Table 1

nGPCR	Nucleotide Sequence (SEQ ID NO:)	Amino acid Sequence (SEQ ID NO:)	Originally filed in:	nGPCR	Nucleotide Sequence (SEQ ID NO:)	Amino acid Sequence (SEQ ID NO:)	Originally filed in:
2356	1	135	A	2403	68	202	H
2357	2	136	A	2404	69	203	H
2358	3	137	A	2405	70	204	H
2359	4	138	A	2406	71	205	H
2360	5	139	A	2407	72	206	H
2361	6	140	A	2408	73	207	H
2362	7	141	A	2409	74	208	H
2363	8	142	A	2410	75	209	H
2364	9	143	A	2411	76	210	H
2365	10	144	A	2412	77	211	I
2366	11	145	B	2413	78	212	I
2367	12	146	B	2414	79	213	I
2368	13	147	B	2415	80	214	I
2369	14	148	B	2416	81	215	I
2370	15	149	B	2417	82	216	I
2371	16	150	B	2418	83	217	I
2372	17	151	B	2419	84	218	I
2373	18	142	B	2420	85	219	I
2374	19	153	B	2421	86	220	I
2375	20	154	B	2422	87	221	J
2376	21	155	C	2423	88	222	J
2377	22	156	C	2424	89	223	J
2378	23	157	C	2425	90	224	J
2379	24	158	C	2426	91	225	J
2380	25	159	C	2427	92	226	J
2381	26	160	C	2428	93	227	J
2382	27	161	C	2429	94	228	J
2383	28	162	C	2430	95	229	J
2384	29	163	C	2431	96	230	J
2385	30	164	C	2432	97	231	K
2386	31	165	D	2433	98	232	K
2387	32	166	D	2434	99	233	K
2388	33	167	D	2435	100	234	K
2389	34	168	D	2436	101	235	K
2390	35	169	D	2437	102	236	K
2391	36	170	D	2438	103	237	K
2392	37	171	D	2439	104	238	K
2393	38	172	D	2440	105	239	K
2394	39	173	D	2441	106	240	K
2395	40	174	D	2442	107	241	L
2396	41	175	E	2443	108	242	L

2397	42	176	E	2444	109	243	L
2398	43	177	E	2445	110	244	L
2399	44	178	E	2446	111	245	L
2400	45	179	E	2447	112	246	L
2401	46	180	E	2448	113	247	L
75	47	181	F	2449	114	248	L
76	48	182	F	2450	115	249	L
77	49	183	F	2451	116	250	L
78	50	184	F	2451	117	251	M
79	51	185	F	2453	118	252	M
80	52	186	F	2454	119	253	M
81	53	187	F	2455	120	254	M
82	54	188	F	2456	121	255	M
83	55	189	F	2457	122	256	M
84	56	190	F	2458	123	257	M
85	57	191	G	2459	124	258	M
2337	58	192	G	2460	125	259	M
2338	59	193	G	2461	126	260	M
2339	60	194	G	2462	127	261	N
2340	61	195	G	2463	128	262	N
2341	62	196	G	2464	129	263	N
2342	63	197	G	2465	130	264	N
2343	64	198	G	2466	131	265	N
2344	65	199	G	2467	132	266	N
2345	66	200	G	2568	133	267	N
2402	67	201	H	74	134	268	O

## Legend

A= Ser. No. 60/187,828  
 C= Ser. No. 60/187,929  
 E= Ser. No. 60/187,825  
 G= Ser. No. 60/187,830  
 I= Ser. No. 60/187,582  
 K= Ser. No. 60/187,714  
 M= Ser. No. 60/187,874  
 O= Ser. No. 60/188,049

B= Ser. No. 60/187,715  
 D= Ser. No. 60/187,930  
 F= Ser. No. 60/187,833  
 H= Ser. No. 60/187,829  
 J= Ser. No. 60/187,581  
 L= Ser. No. 60/189,294  
 N= Ser. No. 60/187,928

When a specific nGPCR is identified (for example nGPCR-2344), it is understood that only that specific nGPCR is being referred to.

As described in Example 5 below, the gene encoding nGPCR-74 (nucleic acid sequence SEQ ID NO:134, amino acid sequence SEQ ID NO:268) has been detected in brain tissue indicating that this nGPCR protein is a neuroreceptor. It is well known that other nGPCR-x are expressed in many different tissues, including the brain. Accordingly, the nGPCR-x of the present invention may be useful, *inter alia*, for treating and/or diagnosing mental disorders. Following the techniques described in Example 5, below, those skilled in the art could readily ascertain if nGPCR-x is expressed in a particular tissue or region.

The invention provides purified and isolated polynucleotides (*e.g.*, cDNA, genomic DNA, synthetic DNA, RNA, or combinations thereof, whether single- or double-stranded) that comprise a nucleotide sequence encoding the amino acid sequence of the polypeptides of the invention. Such polynucleotides are useful for recombinantly expressing the receptor and also for detecting expression of the receptor in cells (*e.g.*, using Northern hybridization and *in situ* hybridization assays). Such polynucleotides also are useful in the design of antisense and other molecules for the suppression of the expression of nGPCR-x in a cultured cell, a tissue, or an animal; for therapeutic purposes; or to provide a model for diseases or conditions characterized by aberrant nGPCR-x expression. Specifically excluded from the definition of polynucleotides of the invention are entire isolated, non-recombinant native chromosomes of host cells. A preferred polynucleotide has a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, which correspond to naturally occurring nGPCR-x sequences. It will be appreciated that numerous other polynucleotide sequences exist that also encode nGPCR-x having the sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, due to the well-known degeneracy of the universal genetic code.

The invention also provides a purified and isolated polynucleotide comprising a nucleotide sequence that encodes a mammalian polypeptide, wherein the polynucleotide hybridizes to a polynucleotide having the sequence set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or the non-coding strand complementary thereto, under the following hybridization conditions:

(a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate; and

(b) washing 2 times for 30 minutes each at 60°C in a wash solution comprising 0.1% SSC, 1% SDS. Polynucleotides that encode a human allelic variant are highly preferred.

The present invention relates to molecules which comprise the gene sequences that encode the nGPCRs; constructs and recombinant host cells incorporating the gene sequences; the novel GPCR polypeptides encoded by the gene sequences; antibodies to the polypeptides and homologs; kits employing the polynucleotides and polypeptides, and methods of making and using all of the foregoing. In addition, the present invention

relates to homologs of the gene sequences and of the polypeptides and methods of making and using the same.

Genomic DNA of the invention comprises the protein-coding region for a polypeptide of the invention and is also intended to include allelic variants thereof. It is  
5 widely understood that, for many genes, genomic DNA is transcribed into RNA transcripts that undergo one or more splicing events wherein intron (*i.e.*, non-coding regions) of the transcripts are removed, or "spliced out." RNA transcripts that can be spliced by alternative mechanisms, and therefore be subject to removal of different RNA sequences but still encode a nGPCR-x polypeptide, are referred to in the art as splice variants which  
10 are embraced by the invention. Splice variants comprehended by the invention therefore are encoded by the same original genomic DNA sequences but arise from distinct mRNA transcripts. Allelic variants are modified forms of a wild-type gene sequence, the modification resulting from recombination during chromosomal segregation or exposure to conditions which give rise to genetic mutation. Allelic variants, like wild type genes,  
15 are naturally occurring sequences (as opposed to non-naturally occurring variants that arise from *in vitro* manipulation).

The invention also comprehends cDNA that is obtained through reverse transcription of an RNA polynucleotide encoding nGPCR-x (conventionally followed by second strand synthesis of a complementary strand to provide a double-stranded DNA).

20 Preferred DNA sequences encoding human nGPCR-x polypeptides are selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134. A preferred DNA of the invention comprises a double stranded molecule along with the complementary molecule (the "non-coding strand" or "complement") having a sequence unambiguously deducible from the coding strand according to Watson-Crick base-pairing rules for DNA. Also  
25 preferred are other polynucleotides encoding the nGPCR-x polypeptide selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, which differ in sequence from the polynucleotides selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, by virtue of the well-known degeneracy of the universal nuclear genetic code.

The invention further embraces other species, preferably mammalian, homologs of  
30 the human nGPCR-x DNA. Species homologs, sometimes referred to as "orthologs," in general, share at least 35%, at least 40%, at least 45%, at least 50%, at least 60%, at least

65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% homology with human DNA of the invention. Generally, percent sequence "homology" with respect to polynucleotides of the invention may be calculated as the percentage of nucleotide bases in the candidate sequence that are identical to nucleotides in the nGPCR-x sequence set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity.

Polynucleotides of the invention permit identification and isolation of polynucleotides encoding related nGPCR-x polypeptides, such as human allelic variants and species homologs, by well-known techniques including Southern and/or Northern hybridization, and polymerase chain reaction (PCR). Examples of related polynucleotides include human and non-human genomic sequences, including allelic variants, as well as polynucleotides encoding polypeptides homologous to nGPCR-x and structurally related polypeptides sharing one or more biological, immunological, and/or physical properties of nGPCR-x. Non-human species genes encoding proteins homologous to nGPCR-x can also be identified by Southern and/or PCR analysis and are useful in animal models for nGPCR-x disorders. Knowledge of the sequence of a human nGPCR-x DNA also makes possible through use of Southern hybridization or polymerase chain reaction (PCR) the identification of genomic DNA sequences encoding nGPCR-x expression control regulatory sequences such as promoters, operators, enhancers, repressors, and the like. Polynucleotides of the invention are also useful in hybridization assays to detect the capacity of cells to express nGPCR-x. Polynucleotides of the invention may also provide a basis for diagnostic methods useful for identifying a genetic alteration(s) in a nGPCR-x locus that underlies a disease state or states, which information is useful both for diagnosis and for selection of therapeutic strategies.

According to the present invention, the nGPCR-x nucleotide sequences disclosed herein may be used to identify homologs of the nGPCR-x, in other animals, including but not limited to humans and other mammals, and invertebrates. Any of the nucleotide sequences disclosed herein, or any portion thereof, can be used, for example, as probes to screen databases or nucleic acid libraries, such as, for example, genomic or cDNA libraries, to identify homologs, using screening procedures well known to those skilled in

the art. Accordingly, homologs having at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, more preferably at least 90%, more preferably at least 95%, and most preferably at least 100% homology with nGPCR-x sequences can be identified.

5       The disclosure herein of full-length polynucleotides encoding nGPCR-x polypeptides makes readily available to the worker of ordinary skill in the art every possible fragment of the full-length polynucleotide.

One preferred embodiment of the present invention provides an isolated nucleic acid molecule comprising a sequence homologous sequences selected from the group  
10       consisting of SEQ ID NO:1 to SEQ ID NO:134, and fragments thereof. Another preferred embodiment provides an isolated nucleic acid molecule comprising a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, and fragments thereof.

As used in the present invention, fragments of nGPCR-x-encoding polynucleotides comprise at least 10, and preferably at least 12, 14, 16, 18, 20, 25, 50, or 75 consecutive  
15       nucleotides of a polynucleotide encoding nGPCR-x. Preferably, fragment polynucleotides of the invention comprise sequences unique to the nGPCR-x-encoding polynucleotide sequence, and therefore hybridize under highly stringent or moderately stringent conditions only (*i.e.*, "specifically") to polynucleotides encoding nGPCR-x (or fragments thereof). Polynucleotide fragments of genomic sequences of the invention comprise not  
20       only sequences unique to the coding region, but also include fragments of the full-length sequence derived from introns, regulatory regions, and/or other non-translated sequences. Sequences unique to polynucleotides of the invention are recognizable through sequence comparison to other known polynucleotides, and can be identified through use of alignment programs routinely utilized in the art, *e.g.*, those made available in public  
25       sequence databases. Such sequences also are recognizable from Southern hybridization analyses to determine the number of fragments of genomic DNA to which a polynucleotide will hybridize. Polynucleotides of the invention can be labeled in a manner that permits their detection, including radioactive, fluorescent, and enzymatic labeling.

30       Fragment polynucleotides are particularly useful as probes for detection of full-length or fragments of nGPCR-x polynucleotides. One or more polynucleotides can be



included in kits that are used to detect the presence of a polynucleotide encoding nGPCR-x, or used to detect variations in a polynucleotide sequence encoding nGPCR-x.

The invention also embraces DNAs encoding nGPCR-x polypeptides that hybridize under moderately stringent or high stringency conditions to the non-coding strand, or complement, of the polynucleotides set forth in sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134.

Exemplary highly stringent hybridization conditions are as follows: hybridization at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% Dextran sulfate, and washing twice for 30 minutes at 60°C in a wash solution comprising 0.1X SSC and 1% SDS. It is understood in the art that conditions of equivalent stringency can be achieved through variation of temperature and buffer, or salt concentration as described Ausubel *et al.* (Eds.), Protocols in Molecular Biology, John Wiley & Sons (1994), pp. 6.0.3 to 6.4.10. Modifications in hybridization conditions can be empirically determined or precisely calculated based on the length and the percentage of guanosine/cytosine (GC) base pairing of the probe. The hybridization conditions can be calculated as described in Sambrook, *et al.*, (Eds.), Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press: Cold Spring Harbor, New York (1989), pp. 9.47 to 9.51.

With the knowledge of the nucleotide sequence information disclosed in the present invention, one skilled in the art can identify and obtain nucleotide sequences which encode nGPCR-x from different sources (*i.e.*, different tissues or different organisms) through a variety of means well known to the skilled artisan and as disclosed by, for example, Sambrook *et al.*, "Molecular cloning: a laboratory manual", Second Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), which is incorporated herein by reference in its entirety.

For example, DNA that encodes nGPCR-x may be obtained by screening of mRNA, cDNA, or genomic DNA with oligonucleotide probes generated from the nGPCR-x gene sequence information provided herein. Probes may be labeled with a detectable group, such as a fluorescent group, a radioactive atom or a chemiluminescent group in accordance with procedures known to the skilled artisan and used in conventional hybridization assays, as described by, for example, Sambrook *et al.*

A nucleic acid molecule comprising any of the nGPCR-x nucleotide sequences described above can alternatively be synthesized by use of the polymerase chain reaction (PCR) procedure, with the PCR oligonucleotide primers produced from the nucleotide sequences provided herein. See U.S. Patent Numbers 4,683,195 to Mullis *et al.* and 4,683,202 to Mullis. The PCR reaction provides a method for selectively increasing the concentration of a particular nucleic acid sequence even when that sequence has not been previously purified and is present only in a single copy in a particular sample. The method can be used to amplify either single- or double-stranded DNA. The essence of the method involves the use of two oligonucleotide probes to serve as primers for the template-dependent, polymerase mediated replication of a desired nucleic acid molecule.

A wide variety of alternative cloning and *in vitro* amplification methodologies are well known to those skilled in the art. Examples of these techniques are found in, for example, Berger *et al.*, *Guide to Molecular Cloning Techniques*, Methods in Enzymology 152, Academic Press, Inc., San Diego, CA (Berger), which is incorporated herein by reference in its entirety.

Automated sequencing methods can be used to obtain or verify the nucleotide sequence of nGPCR-x. The nGPCR-x nucleotide sequences of the present invention are believed to be 100% accurate. However, as is known in the art, nucleotide sequence obtained by automated methods may contain some errors. Nucleotide sequences determined by automation are typically at least about 90%, more typically at least about 95% to at least about 99.9% identical to the actual nucleotide sequence of a given nucleic acid molecule. The actual sequence may be more precisely determined using manual sequencing methods, which are well known in the art. An error in a sequence which results in an insertion or deletion of one or more nucleotides may result in a frame shift in translation such that the predicted amino acid sequence will differ from that which would be predicted from the actual nucleotide sequence of the nucleic acid molecule, starting at the point of the mutation.

The nucleic acid molecules of the present invention, and fragments derived therefrom, are useful for screening for restriction fragment length polymorphism (RFLP) associated with certain disorders, as well as for genetic mapping.

The polynucleotide sequence information provided by the invention makes possible large-scale expression of the encoded polypeptide by techniques well known and routinely practiced in the art.

#### Vectors

5 Another aspect of the present invention is directed to vectors, or recombinant expression vectors, comprising any of the nucleic acid molecules described above. Vectors are used herein either to amplify DNA or RNA encoding nGPCR-x and/or to express DNA which encodes nGPCR-x. Preferred vectors include, but are not limited to, plasmids, phages, cosmids, episomes, viral particles or viruses, and integratable DNA  
10 fragments (*i.e.*, fragments integratable into the host genome by homologous recombination). Preferred viral particles include, but are not limited to, adenoviruses, baculoviruses, parvoviruses, herpesviruses, poxviruses, adeno-associated viruses, Semliki Forest viruses, vaccinia viruses, and retroviruses. Preferred expression vectors include, but are not limited to, pcDNA3 (Invitrogen) and pSVL (Pharmacia Biotech). Other  
15 expression vectors include, but are not limited to, pSPORT™ vectors, pGEM™ vectors (Promega), pPROEXvectors™ (LTI, Bethesda, MD), Bluescript™ vectors (Stratagene), pQE™ vectors (Qiagen), pSE420™ (Invitrogen), and pYES2™ (Invitrogen).

Expression constructs preferably comprise GPCR-x-encoding polynucleotides operatively linked to an endogenous or exogenous expression control DNA sequence and  
20 a transcription terminator. Expression control DNA sequences include promoters, enhancers, operators, and regulatory element binding sites generally, and are typically selected based on the expression systems in which the expression construct is to be utilized. Preferred promoter and enhancer sequences are generally selected for the ability to increase gene expression, while operator sequences are generally selected for the ability  
25 to regulate gene expression. Expression constructs of the invention may also include sequences encoding one or more selectable markers that permit identification of host cells bearing the construct. Expression constructs may also include sequences that facilitate, and preferably promote, homologous recombination in a host cell. Preferred constructs of the invention also include sequences necessary for replication in a host cell.

30 Expression constructs are preferably utilized for production of an encoded protein, but may also be utilized simply to amplify a nGPCR-x-encoding polynucleotide sequence.

In preferred embodiments, the vector is an expression vector wherein the polynucleotide of the invention is operatively linked to a polynucleotide comprising an expression control sequence. Autonomously replicating recombinant expression constructs such as plasmid and viral DNA vectors incorporating polynucleotides of the invention are also provided.

5 Preferred expression vectors are replicable DNA constructs in which a DNA sequence encoding nGPCR-x is operably linked or connected to suitable control sequences capable of effecting the expression of the nGPCR-x in a suitable host. DNA regions are operably linked or connected when they are functionally related to each other. For example, a promoter is operably linked or connected to a coding sequence if it controls the  
10 transcription of the sequence. Amplification vectors do not require expression control domains, but rather need only the ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants. The need for control sequences in the expression vector will vary depending upon the host selected and the transformation method chosen. Generally, control sequences include a  
15 transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding and sequences which control the termination of transcription and translation.

Preferred vectors preferably contain a promoter that is recognized by the host organism. The promoter sequences of the present invention may be prokaryotic,  
20 eukaryotic or viral. Examples of suitable prokaryotic sequences include the  $P_R$  and  $P_L$  promoters of bacteriophage lambda (The bacteriophage Lambda, Hershey, A. D., Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY (1973), which is incorporated herein by reference in its entirety; Lambda II, Hendrix, R. W., Ed., Cold Spring Harbor Press, Cold Spring Harbor, NY (1980), which is incorporated herein by reference in its entirety);  
25 the *trp*, *recA*, heat shock, and *lacZ* promoters of *E. coli* and the SV40 early promoter (Benoist *et al. Nature*, 1981, 290, 304-310, which is incorporated herein by reference in its entirety). Additional promoters include, but are not limited to, mouse mammary tumor virus, long terminal repeat of human immunodeficiency virus, maloney virus, cytomegalovirus immediate early promoter, Epstein Barr virus, Rous sarcoma virus,  
30 human actin, human myosin, human hemoglobin, human muscle creatine, and human metallothionein.

Additional regulatory sequences can also be included in preferred vectors. Preferred examples of suitable regulatory sequences are represented by the Shine-Dalgarno of the replicase gene of the phage MS-2 and of the gene cII of bacteriophage lambda. The Shine-Dalgarno sequence may be directly followed by DNA encoding  
5 nGPCR-x and result in the expression of the mature nGPCR-x protein.

Moreover, suitable expression vectors can include an appropriate marker that allows the screening of the transformed host cells. The transformation of the selected host is carried out using any one of the various techniques well known to the expert in the art and described in Sambrook *et al.*, *supra*.

10 An origin of replication can also be provided either by construction of the vector to include an exogenous origin or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter may be sufficient. Alternatively, rather than using vectors which contain viral origins of replication, one skilled in the art can transform mammalian cells by the method of co-  
15 transformation with a selectable marker and nGPCR-x DNA. An example of a suitable marker is dihydrofolate reductase (DHFR) or thymidine kinase (*see*, U.S. Patent No. 4,399,216).

Nucleotide sequences encoding GPCR-x may be recombined with vector DNA in accordance with conventional techniques, including blunt-ended or staggered-ended  
20 termini for ligation, restriction enzyme digestion to provide appropriate termini, filling in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and ligation with appropriate ligases. Techniques for such manipulation are disclosed by Sambrook *et al.*, *supra* and are well known in the art. Methods for construction of mammalian expression vectors are disclosed in, for example, Okayama *et al.*, *Mol. Cell. Biol.*, **1983**, *3*, 280, Cosman *et al.*, *Mol. Immunol.*, **1986**, *23*, 935, Cosman *et al.*, *Nature*, **1984**, *312*, 768, EP-A-0367566, and WO 91/18982, each of which is  
25 incorporated herein by reference in its entirety.

#### Host cells

According to another aspect of the invention, host cells are provided, including  
30 prokaryotic and eukaryotic cells, comprising a polynucleotide of the invention (or vector of the invention) in a manner that permits expression of the encoded nGPCR-x

polypeptide. Polynucleotides of the invention may be introduced into the host cell as part of a circular plasmid, or as linear DNA comprising an isolated protein coding region or a viral vector. Methods for introducing DNA into the host cell that are well known and routinely practiced in the art include transformation, transfection, electroporation, nuclear  
5 injection, or fusion with carriers such as liposomes, micelles, ghost cells, and protoplasts. Expression systems of the invention include bacterial, yeast, fungal, plant, insect, invertebrate, vertebrate, and mammalian cells systems.

The invention provides host cells that are transformed or transfected (stably or transiently) with polynucleotides of the invention or vectors of the invention. As stated  
10 above, such host cells are useful for amplifying the polynucleotides and also for expressing the nGPCR-x polypeptide or fragment thereof encoded by the polynucleotide.

In still another related embodiment, the invention provides a method for producing a nGPCR-x polypeptide (or fragment thereof) comprising the steps of growing a host cell of the invention in a nutrient medium and isolating the polypeptide or variant thereof from  
15 the cell or the medium. Because nGPCR-x is a seven transmembrane receptor, it will be appreciated that, for some applications, such as certain activity assays, the preferable isolation may involve isolation of cell membranes containing the polypeptide embedded therein, whereas for other applications a more complete isolation may be preferable.

According to some aspects of the present invention, transformed host cells having  
20 an expression vector comprising any of the nucleic acid molecules described above are provided. Expression of the nucleotide sequence occurs when the expression vector is introduced into an appropriate host cell. Suitable host cells for expression of the polypeptides of the invention include, but are not limited to, prokaryotes, yeast, and eukaryotes. If a prokaryotic expression vector is employed, then the appropriate host cell  
25 would be any prokaryotic cell capable of expressing the cloned sequences. Suitable prokaryotic cells include, but are not limited to, bacteria of the genera *Escherichia*, *Bacillus*, *Salmonella*, *Pseudomonas*, *Streptomyces*, and *Staphylococcus*.

If an eukaryotic expression vector is employed, then the appropriate host cell would be any eukaryotic cell capable of expressing the cloned sequence. Preferably,  
30 eukaryotic cells are cells of higher eukaryotes. Suitable eukaryotic cells include, but are not limited to, non-human mammalian tissue culture cells and human tissue culture cells.

Preferred host cells include, but are not limited to, insect cells, HeLa cells, Chinese hamster ovary cells (CHO cells), African green monkey kidney cells (COS cells), human HEK-293 cells, and murine 3T3 fibroblasts. Propagation of such cells in cell culture has become a routine procedure (*see*, Tissue Culture, Academic Press, Kruse and Patterson, eds. (1973), which is incorporated herein by reference in its entirety).

In addition, a yeast host may be employed as a host cell. Preferred yeast cells include, but are not limited to, the genera *Saccharomyces*, *Pichia*, and *Kluveromyces*. Preferred yeast hosts are *S. cerevisiae* and *P. pastoris*. Preferred yeast vectors can contain an origin of replication sequence from a 2T yeast plasmid, an autonomously replication sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Shuttle vectors for replication in both yeast and *E. coli* are also included herein.

Alternatively, insect cells may be used as host cells. In a preferred embodiment, the polypeptides of the invention are expressed using a baculovirus expression system (*see*, Luckow *et al.*, *Bio/Technology*, 1988, 6, 47, Baculovirus Expression Vectors: A Laboratory Manual, O'Rielly *et al.* (Eds.), W.H. Freeman and Company, New York, 1992, and U.S. Patent No. 4,879,236, each of which is incorporated herein by reference in its entirety). In addition, the MAXBAC™ complete baculovirus expression system (Invitrogen) can, for example, be used for production in insect cells.

Host cells of the invention are a valuable source of immunogen for development of antibodies specifically immunoreactive with nGPCR-x. Host cells of the invention are also useful in methods for the large-scale production of nGPCR-x polypeptides wherein the cells are grown in a suitable culture medium and the desired polypeptide products are isolated from the cells, or from the medium in which the cells are grown, by purification methods known in the art, *e.g.*, conventional chromatographic methods including immunoaffinity chromatography, receptor affinity chromatography, hydrophobic interaction chromatography, lectin affinity chromatography, size exclusion filtration, cation or anion exchange chromatography, high pressure liquid chromatography (HPLC), reverse phase HPLC, and the like. Still other methods of purification include those methods wherein the desired protein is expressed and purified as a fusion protein having a specific tag, label, or chelating moiety that is recognized by a specific binding partner or

agent. The purified protein can be cleaved to yield the desired protein, or can be left as an intact fusion protein. Cleavage of the fusion component may produce a form of the desired protein having additional amino acid residues as a result of the cleavage process.

Knowledge of nGPCR-x DNA sequences allows for modification of cells to permit, or increase, expression of endogenous nGPCR-x. Cells can be modified (e.g. by homologous recombination) to provide increased expression by replacing, in whole or in part, the naturally occurring nGPCR-x promoter with all or part of a heterologous promoter so that the cells express nGPCR-x at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to endogenous nGPCR-x encoding sequences. (See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955.) It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamoyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the nGPCR-x coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the nGPCR-x coding sequences in the cells.

#### **Knock-outs**

The DNA sequence information provided by the present invention also makes possible the development (e.g., by homologous recombination or "knock-out" strategies; see Capecchi, *Science* 244:1288-1292 (1989), which is incorporated herein by reference) of animals that fail to express functional nGPCR-x or that express a variant of nGPCR-x. Such animals (especially small laboratory animals such as rats, rabbits, and mice) are useful as models for studying the *in vivo* activities of nGPCR-x and modulators of nGPCR-x.

#### **Antisense**

Also made available by the invention are anti-sense polynucleotides that recognize and hybridize to polynucleotides encoding nGPCR-x. Full-length and fragment anti-sense polynucleotides are provided. Fragment antisense molecules of the invention include (i) those that specifically recognize and hybridize to nGPCR-x RNA (as determined by sequence comparison of DNA encoding nGPCR-x to DNA encoding other known



molecules). Identification of sequences unique to nGPCR-x encoding polynucleotides can be deduced through use of any publicly available sequence database, and/or through use of commercially available sequence comparison programs. After identification of the desired sequences, isolation through restriction digestion or amplification using any of the various polymerase chain reaction techniques well known in the art can be performed. Anti-sense polynucleotides are particularly relevant to regulating expression of nGPCR-x by those cells expressing nGPCR-x mRNA.

Antisense nucleic acids (preferably 10 to 30 base-pair oligonucleotides) capable of specifically binding to nGPCR-x expression control sequences or nGPCR-x RNA are introduced into cells (e.g., by a viral vector or colloidal dispersion system such as a liposome). The antisense nucleic acid binds to the nGPCR-x target nucleotide sequence in the cell and prevents transcription and/or translation of the target sequence. Phosphorothioate and methylphosphonate antisense oligonucleotides are specifically contemplated for therapeutic use by the invention. The antisense oligonucleotides may be further modified by adding poly-L-lysine, transferrin polylysine, or cholesterol moieties at their 5' end. Suppression of nGPCR-x expression at either the transcriptional or translational level is useful to generate cellular or animal models for diseases/conditions characterized by aberrant nGPCR-x expression.

Antisense oligonucleotides, or fragments of sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or sequences complementary or homologous thereto, derived from the nucleotide sequences of the present invention encoding nGPCR-x are useful as diagnostic tools for probing gene expression in various tissues. For example, tissue can be probed *in situ* with oligonucleotide probes carrying detectable groups by conventional autoradiography techniques to investigate native expression of this enzyme or pathological conditions relating thereto. Antisense oligonucleotides are preferably directed to regulatory regions of sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or mRNA corresponding thereto, including, but not limited to, the initiation codon, TATA box, enhancer sequences, and the like.

**Transcription factors**

The nGPCR-x sequences taught in the present invention facilitate the design of novel transcription factors for modulating nGPCR-x expression in native cells and animals, and cells transformed or transfected with nGPCR-x polynucleotides. For example, the Cys<sub>2</sub>-His<sub>2</sub> zinc finger proteins, which bind DNA via their zinc finger domains, have been shown to be amenable to structural changes that lead to the recognition of different target sequences. These artificial zinc finger proteins recognize specific target sites with high affinity and low dissociation constants, and are able to act as gene switches to modulate gene expression. Knowledge of the particular nGPCR-x target sequence of the present invention facilitates the engineering of zinc finger proteins specific for the target sequence using known methods such as a combination of structure-based modeling and screening of phage display libraries (Segal *et al.*, Proc. Natl. Acad. Sci. (USA) 96:2758-2763 (1999); Liu *et al.*, Proc. Natl. Acad. Sci. (USA) 94:5525-5530 (1997); Greisman *et al.*, Science 275:657-661 (1997); Choo *et al.*, J. Mol. Biol. 273:525-532 (1997)). Each zinc finger domain usually recognizes three or more base pairs. Since a recognition sequence of 18 base pairs is generally sufficient in length to render it unique in any known genome, a zinc finger protein consisting of 6 tandem repeats of zinc fingers would be expected to ensure specificity for a particular sequence (Segal *et al.*) The artificial zinc finger repeats, designed based on nGPCR-x sequences, are fused to activation or repression domains to promote or suppress nGPCR-x expression (Liu *et al.*) Alternatively, the zinc finger domains can be fused to the TATA box-binding factor (TBP) with varying lengths of linker region between the zinc finger peptide and the TBP to create either transcriptional activators or repressors (Kim *et al.*, Proc. Natl. Acad. Sci. (USA) 94:3616-3620 (1997). Such proteins and polynucleotides that encode them, have utility for modulating nGPCR-x expression *in vivo* in both native cells, animals and humans; and/or cells transfected with nGPCR-x-encoding sequences. The novel transcription factor can be delivered to the target cells by transfecting constructs that express the transcription factor (gene therapy), or by introducing the protein. Engineered zinc finger proteins can also be designed to bind RNA sequences for use in therapeutics as alternatives to antisense or catalytic RNA methods (McColl *et al.*, Proc. Natl. Acad. Sci. (USA) 96:9521-9526 (1997); Wu *et al.*, Proc. Natl. Acad. Sci. (USA) 92:344-348 (1995)). The present invention contemplates methods of designing such transcription factors based

on the gene sequence of the invention, as well as customized zinc finger proteins, that are useful to modulate nGPCR-x expression in cells (native or transformed) whose genetic complement includes these sequences.

### Polypeptides

5       The invention also provides purified and isolated mammalian nGPCR-x polypeptides encoded by a polynucleotide of the invention. Presently preferred is a human nGPCR-x polypeptide comprising the amino acid sequence set out in sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, or fragments thereof comprising an epitope specific to the polypeptide. By "epitope specific to" is meant a  
10       portion of the nGPCR receptor that is recognizable by an antibody that is specific for the nGPCR, as defined in detail below.

Although the sequences provided are particular human sequences, the invention is intended to include within its scope other human allelic variants; non-human mammalian forms of nGPCR-x, and other vertebrate forms of nGPCR-x.

15       It will be appreciated that extracellular epitopes are particularly useful for generating and screening for antibodies and other binding compounds that bind to receptors such as nGPCR-x. Thus, in another preferred embodiment, the invention provides a purified and isolated polypeptide comprising at least one extracellular domain (*e.g.*, the N-terminal extracellular domain or one of the three extracellular loops) of  
20       nGPCR-x. Purified and isolated polypeptides comprising the N-terminal extracellular domain of nGPCR-x are highly preferred. Also preferred is a purified and isolated polypeptide comprising a nGPCR-x fragment selected from the group consisting of the N-terminal extracellular domain of nGPCR-x, transmembrane domains of nGPCR-x, an extracellular loop connecting transmembrane domains of nGPCR-x, an intracellular loop  
25       connecting transmembrane domains of nGPCR-x, the C-terminal cytoplasmic region of nGPCR-x, and fusions thereof. Such fragments may be continuous portions of the native receptor. However, it will also be appreciated that knowledge of the nGPCR-x gene and protein sequences as provided herein permits recombining of various domains that are not contiguous in the native protein. Using a FORTRAN computer program called  
30       "tmtest.all" [Parodi *et al.*, Comput. Appl. Biosci. 5:527-535 (1994)], nGPCR-x was shown to contain transmembrane-spanning domains.

The invention also embraces polypeptides that have at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55% or at least 50% identity and/or homology to the preferred polypeptide of the invention. Percent amino acid sequence "identity" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in the nGPCR-x sequence after aligning both sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Percent sequence "homology" with respect to the preferred polypeptide of the invention is defined herein as the percentage of amino acid residues in the candidate sequence that are identical with the residues in the nGPCR-x sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and also considering any conservative substitutions as part of the sequence identity.

In one aspect, percent homology is calculated as the percentage of amino acid residues in the smaller of two sequences which align with identical amino acid residue in the sequence being compared, when four gaps in a length of 100 amino acids may be introduced to maximize alignment (Dayhoff, in Atlas of Protein Sequence and Structure, Vol. 5, p. 124, National Biochemical Research Foundation, Washington, D.C. (1972), incorporated herein by reference).

Polypeptides of the invention may be isolated from natural cell sources or may be chemically synthesized, but are preferably produced by recombinant procedures involving host cells of the invention. Use of mammalian host cells is expected to provide for such post-translational modifications (e.g., glycosylation, truncation, lipidation, and phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention. Glycosylated and non-glycosylated forms of nGPCR-x polypeptides are embraced by the invention.

The invention also embraces variant (or analog) nGPCR-x polypeptides. In one example, insertion variants are provided wherein one or more amino acid residues supplement a nGPCR-x amino acid sequence. Insertions may be located at either or both termini of the protein, or may be positioned within internal regions of the nGPCR-x amino

acid sequence. Insertional variants with additional residues at either or both termini can include, for example, fusion proteins and proteins including amino acid tags or labels.

Insertion variants include nGPCR-x polypeptides wherein one or more amino acid residues are added to a nGPCR-x acid sequence or to a biologically active fragment thereof.

Variant products of the invention also include mature nGPCR-x products, *i.e.*, nGPCR-x products wherein leader or signal sequences are removed, with additional amino terminal residues. The additional amino terminal residues may be derived from another protein, or may include one or more residues that are not identifiable as being derived from specific proteins. nGPCR-x products with an additional methionine residue at position -1 (Met<sup>-1</sup>-nGPCR-x) are contemplated, as are variants with additional methionine and lysine residues at positions -2 and -1 (Met<sup>-2</sup>-Lys<sup>-1</sup>-nGPCR-x). Variants of nGPCR-x with additional Met, Met-Lys, Lys residues (or one or more basic residues in general) are particularly useful for enhanced recombinant protein production in bacterial host cells.

The invention also embraces nGPCR-x variants having additional amino acid residues that result from use of specific expression systems. For example, use of commercially available vectors that express a desired polypeptide as part of a glutathione-S-transferase (GST) fusion product provides the desired polypeptide having an additional glycine residue at position -1 after cleavage of the GST component from the desired polypeptide. Variants that result from expression in other vector systems are also contemplated.

Insertional variants also include fusion proteins wherein the amino terminus and/or the carboxy terminus of nGPCR-x is/are fused to another polypeptide.

In another aspect, the invention provides deletion variants wherein one or more amino acid residues in a nGPCR-x polypeptide are removed. Deletions can be effected at one or both termini of the nGPCR-x polypeptide, or with removal of one or more non-terminal amino acid residues of nGPCR-x. Deletion variants, therefore, include all fragments of a nGPCR-x polypeptide.

The invention also embraces polypeptide fragments of sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, wherein the fragments maintain

biological (*e.g.*, ligand binding and/or intracellular signaling) immunological properties of a nGPCR-x polypeptide.

In one preferred embodiment of the invention, an isolated nucleic acid molecule comprises a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, and fragments thereof, wherein the nucleic acid molecule encoding at least a portion of nGPCR-x. In a more preferred embodiment, the isolated nucleic acid molecule comprises a sequence that encodes a polypeptide comprising sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, and fragments thereof.

As used in the present invention, polypeptide fragments comprise at least 5, 10, 15, 20, 25, 30, 35, or 40 consecutive amino acids of sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268. Preferred polypeptide fragments display antigenic properties unique to, or specific for, human nGPCR-x and its allelic and species homologs. Fragments of the invention having the desired biological and immunological properties can be prepared by any of the methods well known and routinely practiced in the art.

In still another aspect, the invention provides substitution variants of nGPCR-x polypeptides. Substitution variants include those polypeptides wherein one or more amino acid residues of a nGPCR-x polypeptide are removed and replaced with alternative residues. In one aspect, the substitutions are conservative in nature; however, the invention embraces substitutions that are also non-conservative. Conservative substitutions for this purpose may be defined as set out in Tables 2, 3, or 4 below.

Variant polypeptides include those wherein conservative substitutions have been introduced by modification of polynucleotides encoding polypeptides of the invention. Amino acids can be classified according to physical properties and contribution to secondary and tertiary protein structure. A conservative substitution is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are set out in Table 2 (from WO 97/09433, page 10, published March 13, 1997 (PCT/GB96/02197, filed 9/6/96), immediately below.

Table 2  
Conservative Substitutions I

<u>SIDE CHAIN CHARACTERISTIC</u>	<u>AMINO ACID</u>
Aliphatic	
Non-polar	G A P I L V
Polar - uncharged	C S T M N Q
Polar - charged	D E K R
Aromatic	H F W Y
Other	N Q D E

Alternatively, conservative amino acids can be grouped as described in Lehninger, [Biochemistry, Second Edition; Worth Publishers, Inc. NY, NY (1975), pp.71-77] as set out in Table 3, below.

Table 3  
Conservative Substitutions II

<u>SIDE CHAIN CHARACTERISTIC</u>	<u>AMINO ACID</u>
Non-polar (hydrophobic)	
A. Aliphatic:	A L I V P
B. Aromatic:	F W
C. Sulfur-containing:	M
D. Borderline:	G
Uncharged-polar	
A. Hydroxyl:	S T Y
B. Amides:	N Q
C. Sulfhydryl:	C
D. Borderline:	G
Positively Charged (Basic):	K R H
Negatively Charged (Acidic):	D E

As still another alternative, exemplary conservative substitutions are set out in Table 4, below.

Table 4  
Conservative Substitutions III

<u>Original Residue</u>	<u>Exemplary Substitution</u>
Ala (A)	Val, Leu, Ile
Arg (R)	Lys, Gln, Asn
Asn (N)	Gln, His, Lys, Arg
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (E)	Asp
His (H)	Asn, Gln, Lys, Arg

Ile (I)	Leu, Val, Met, Ala, Phe,
Leu (L)	Ile, Val, Met, Ala, Phe
Lys (K)	Arg, Gln, Asn
Met (M)	Leu, Phe, Ile
Phe (F)	Leu, Val, Ile, Ala
Pro (P)	Gly
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Trp, Phe, Thr, Ser
Val (V)	Ile, Leu, Met, Phe, Ala

It should be understood that the definition of polypeptides of the invention is intended to include polypeptides bearing modifications other than insertion, deletion, or substitution of amino acid residues. By way of example, the modifications may be covalent in nature, and include for example, chemical bonding with polymers, lipids, other organic, and inorganic moieties. Such derivatives may be prepared to increase circulating half-life of a polypeptide, or may be designed to improve the targeting capacity of the polypeptide for desired cells, tissues, or organs. Similarly, the invention further embraces nGPCR-x polypeptides that have been covalently modified to include one or more water-soluble polymer attachments such as polyethylene glycol, polyoxyethylene glycol, or polypropylene glycol. Variants that display ligand binding properties of native nGPCR-x and are expressed at higher levels, as well as variants that provide for constitutively active receptors, are particularly useful in assays of the invention; the variants are also useful in providing cellular, tissue and animal models of diseases/conditions characterized by aberrant nGPCR-x activity.

In a related embodiment, the present invention provides compositions comprising purified polypeptides of the invention. Preferred compositions comprise, in addition to the polypeptide of the invention, a pharmaceutically acceptable (*i.e.*, sterile and non-toxic) liquid, semisolid, or solid diluent that serves as a pharmaceutical vehicle, excipient, or medium. Any diluent known in the art may be used. Exemplary diluents include, but are not limited to, water, saline solutions, polyoxyethylene sorbitan monolaurate, magnesium stearate, methyl- and propylhydroxybenzoate, talc, alginates, starches, lactose, sucrose, dextrose, sorbitol, mannitol, glycerol, calcium phosphate, mineral oil, and cocoa butter.



5 Variants that display ligand binding properties of native nGPCR-x and are expressed at higher levels, as well as variants that provide for constitutively active receptors, are particularly useful in assays of the invention; the variants are also useful in assays of the invention and in providing cellular, tissue and animal models of diseases/conditions characterized by aberrant nGPCR-x activity.

10 The G protein-coupled receptor functions through a specific heterotrimeric guanine-nucleotide-binding regulatory protein (G-protein) coupled to the intracellular portion of the G protein-coupled receptor molecule. Accordingly, the G protein-coupled receptor has a specific affinity to G protein. G proteins specifically bind to guanine nucleotides. Isolation of G proteins provides a means to isolate guanine nucleotides. G proteins may be isolated using commercially available anti-G protein antibodies or isolated G protein-coupled receptors. Similarly, G proteins may be detected in a sample isolated using commercially available detectable anti-G protein antibodies or isolated G protein-coupled receptors.

15 According to the present invention, the isolated nGPCR-x proteins of the present invention are useful to isolate and purify G proteins from samples such as cell lysates. Example 15 below sets forth an example of isolation of G proteins using isolated nGPCR-x proteins. Such methodology may be used in place of the use of commercially available anti-G protein antibodies which are used to isolate G proteins. Moreover, G proteins may be detected using n-GPCR-x proteins in place of commercially available detectable anti-G protein antibodies. Since nGPCR-x proteins specifically bind to G proteins, they can be employed in any specific use where G protein specific affinity is required such as those uses where commercially available anti-G protein antibodies are employed.

#### Antibodies

25 Also comprehended by the present invention are antibodies (e.g., monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies, bifunctional/bispecific antibodies, humanized antibodies, human antibodies, and complementary determining region (CDR)-grafted antibodies, including compounds which include CDR sequences which specifically recognize a polypeptide of the invention) specific for nGPCR-x or fragments thereof. Preferred antibodies of the invention are human antibodies that are produced and identified according to methods described in WO93/11236, published June

30

20, 1993, which is incorporated herein by reference in its entirety. Antibody fragments, including Fab, Fab', F(ab')<sub>2</sub>, and F<sub>v</sub>, are also provided by the invention. The term "specific for," when used to describe antibodies of the invention, indicates that the variable regions of the antibodies of the invention recognize and bind nGPCR-x polypeptides exclusively (*i.e.*, are able to distinguish nGPCR-x polypeptides from other known GPCR polypeptides by virtue of measurable differences in binding affinity, despite the possible existence of localized sequence identity, homology, or similarity between nGPCR-x and such polypeptides). It will be understood that specific antibodies may also interact with other proteins (for example, *S. aureus* protein A or other antibodies in ELISA techniques) through interactions with sequences outside the variable region of the antibodies, and, in particular, in the constant region of the molecule. Screening assays to determine binding specificity of an antibody of the invention are well known and routinely practiced in the art. For a comprehensive discussion of such assays, see Harlow *et al.* (Eds.), Antibodies A Laboratory Manual; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988), Chapter 6. Antibodies that recognize and bind fragments of the nGPCR-x polypeptides of the invention are also contemplated, provided that the antibodies are specific for nGPCR-x polypeptides. Antibodies of the invention can be produced using any method well known and routinely practiced in the art.

The invention provides an antibody that is specific for the nGPCR-x of the invention. Antibody specificity is described in greater detail below. However, it should be emphasized that antibodies that can be generated from polypeptides that have previously been described in the literature and that are capable of fortuitously cross-reacting with nGPCR-x (*e.g.*, due to the fortuitous existence of a similar epitope in both polypeptides) are considered "cross-reactive" antibodies. Such cross-reactive antibodies are not antibodies that are "specific" for nGPCR-x. The determination of whether an antibody is specific for nGPCR-x or is cross-reactive with another known receptor is made using any of several assays, such as Western blotting assays, that are well known in the art. For identifying cells that express nGPCR-x and also for modulating nGPCR-x-ligand binding activity, antibodies that specifically bind to an extracellular epitope of the nGPCR-x are preferred.

In one preferred variation, the invention provides monoclonal antibodies. Hybridomas that produce such antibodies also are intended as aspects of the invention. In yet another variation, the invention provides a humanized antibody. Humanized antibodies are useful for *in vivo* therapeutic indications.

5 In another variation, the invention provides a cell-free composition comprising polyclonal antibodies, wherein at least one of the antibodies is an antibody of the invention specific for nGPCR-x. Antisera isolated from an animal is an exemplary composition, as is a composition comprising an antibody fraction of an antisera that has been resuspended in water or in another diluent, excipient, or carrier.

10 In still another related embodiment, the invention provides an anti-idiotypic antibody specific for an antibody that is specific for nGPCR-x.

It is well known that antibodies contain relatively small antigen binding domains that can be isolated chemically or by recombinant techniques. Such domains are useful nGPCR-x binding molecules themselves, and also may be reintroduced into human antibodies, or fused to toxins or other polypeptides. Thus, in still another embodiment, the invention provides a polypeptide comprising a fragment of a nGPCR-x-specific antibody, wherein the fragment and the polypeptide bind to the nGPCR-x. By way of non-limiting example, the invention provides polypeptides that are single chain antibodies and CDR-grafted antibodies.

20 Non-human antibodies may be humanized by any of the methods known in the art. In one method, the non-human CDRs are inserted into a human antibody or consensus antibody framework sequence. Further changes can then be introduced into the antibody framework to modulate affinity or immunogenicity.

25 Antibodies of the invention are useful for, *e.g.*, therapeutic purposes (by modulating activity of nGPCR-x), diagnostic purposes to detect or quantitate nGPCR-x, and purification of nGPCR-x. Kits comprising an antibody of the invention for any of the purposes described herein are also comprehended. In general, a kit of the invention also includes a control antigen for which the antibody is immunospecific.

#### Compositions

30 Mutations in the nGPCR-x gene that result in loss of normal function of the nGPCR-x gene product underlie nGPCR-x-related human disease states. The invention

comprehends gene therapy to restore nGPCR-x activity to treat those disease states. Delivery of a functional nGPCR-x gene to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and more particularly viral vectors (*e.g.*, adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (*e.g.*, liposomes or chemical treatments). See, for example, Anderson, *Nature*, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, *Science*, 244: 1275-1281 (1989); Verma, *Scientific American*: 68-84 (1990); and Miller, *Nature*, 357: 455-460 (1992). Alternatively, it is contemplated that in other human disease states, preventing the expression of, or inhibiting the activity of, nGPCR-x will be useful in treating disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of nGPCR-x.

Another aspect of the present invention is directed to compositions, including pharmaceutical compositions, comprising any of the nucleic acid molecules or recombinant expression vectors described above and an acceptable carrier or diluent. Preferably, the carrier or diluent is pharmaceutically acceptable. Suitable carriers are described in the most recent edition of *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in this field, which is incorporated herein by reference in its entirety. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils may also be used. The formulations are sterilized by commonly used techniques.

Also within the scope of the invention are compositions comprising polypeptides, polynucleotides, or antibodies of the invention that have been formulated with, *e.g.*, a pharmaceutically acceptable carrier.

The invention also provides methods of using antibodies of the invention. For example, the invention provides a method for modulating ligand binding of a nGPCR-x comprising the step of contacting the nGPCR-x with an antibody specific for the nGPCR-x, under conditions wherein the antibody binds the receptor.

As discussed above, it is well known that GPCRs are expressed in many different tissues and regions, including in the brain. GPCRs that may be expressed in the brain, such as nGPCR-x, provide an indication that aberrant nGPCR-x signaling activity may

correlate with one or more neurological or psychological disorders. The invention also provides a method for treating a neurological or psychiatric disorder comprising the step of administering to a mammal in need of such treatment an amount of an antibody-like polypeptide of the invention that is sufficient to modulate ligand binding to a nGPCR-x in  
5 neurons of the mammal. nGPCR-x may also be expressed in other tissues, including but not limited to, peripheral blood lymphocytes, pancreas, ovary, uterus, testis, salivary gland, thyroid gland, kidney, adrenal gland, liver, bone marrow, prostate, fetal liver, colon, muscle, and fetal brain, and may be found in many other tissues. Within the brain, nGPCR-x mRNA transcripts may be found in many tissues, including, but not limited to,  
10 frontal lobe, hypothalamus, pons, cerebellum, caudate nucleus, and medulla.

#### Kits

The present invention is also directed to kits, including pharmaceutical kits. The kits can comprise any of the nucleic acid molecules described above, any of the polypeptides described above, or any antibody which binds to a polypeptide of the  
15 invention as described above, as well as a negative control. The kit preferably comprises additional components, such as, for example, instructions, solid support, reagents helpful for quantification, and the like.

In another aspect, the invention features methods for detection of a polypeptide in a sample as a diagnostic tool for diseases or disorders, wherein the method comprises the  
20 steps of: (a) contacting the sample with a nucleic acid probe which hybridizes under hybridization assay conditions to a nucleic acid target region of a polypeptide having sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, said probe comprising the nucleic acid sequence encoding the polypeptide, fragments thereof, and the complements of the sequences and fragments; and (b) detecting the presence or  
25 amount of the probe:target region hybrid as an indication of the disease.

In preferred embodiments of the invention, the disease is selected from the group consisting of thyroid disorders (*e.g.* thyrotoxicosis, myxoedema); renal failure; inflammatory conditions (*e.g.*, Crohn's disease); diseases related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS  
30 disorders (*e.g.*, pain including migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety

disorder, post-traumatic-stress disorder, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); and sexual dysfunction, among others.

As described above and in Example 5 below, the gene encoding nGPCR-74 (nucleic acid sequence SEQ ID NO:134, amino acid sequence SEQ ID NO:268) has been detected in brain tissue indicating that this nGPCR protein is a neuroreceptor. It is well known that other nGPCR-x are expressed in many different tissues, including the brain. Accordingly, the nGPCR-x of the present invention may be useful, *inter alia*, for treating and/or diagnosing mental disorders. Following the techniques described in Example 5, below, those skilled in the art could readily ascertain if nGPCR-x is expressed in a particular tissue or region.

Kits may be designed to detect either expression of polynucleotides encoding nGPCR-x expressed in the brain or the nGPCR-x proteins themselves in order to identify tissue as being neurological. For example, oligonucleotide hybridization kits can be provided which include a container having an oligonucleotide probe specific for the nGPCR-x-specific DNA and optionally, containers with positive and negative controls and/or instructions. Similarly, PCR kits can be provided which include a container having primers specific for the nGPCR-x-specific sequences, DNA and optionally, containers with size markers, positive and negative controls and/or instructions.

Hybridization conditions should be such that hybridization occurs only with the genes in the presence of other nucleic acid molecules. Under stringent hybridization conditions only highly complementary nucleic acid sequences hybridize. Preferably, such

conditions prevent hybridization of nucleic acids having 1 or 2 mismatches out of 20 contiguous nucleotides. Such conditions are defined supra.

The diseases for which detection of genes in a sample could be diagnostic include diseases in which nucleic acid (DNA and/or RNA) is amplified in comparison to normal  
5 cells. By "amplification" is meant increased numbers of DNA or RNA in a cell compared with normal cells.

The diseases that could be diagnosed by detection of nucleic acid in a sample preferably include central nervous system and metabolic diseases. The test samples suitable for nucleic acid probing methods of the present invention include, for example,  
10 cells or nucleic acid extracts of cells, or biological fluids. The samples used in the above-described methods will vary based on the assay format, the detection method and the nature of the tissues, cells or extracts to be assayed. Methods for preparing nucleic acid extracts of cells are well known in the art and can be readily adapted in order to obtain a sample that is compatible with the method utilized.

15 Alternatively, immunoassay kits can be provided which have containers container having antibodies specific for the nGPCR-x-protein and optionally, containers with positive and negative controls and/or instructions.

Kits may also be provided useful in the identification of GPCR binding partners such as natural ligands or modulators (agonists or antagonists). Substances useful for  
20 treatment of disorders or diseases preferably show positive results in one or more *in vitro* assays for an activity corresponding to treatment of the disease or disorder in question. Substances that modulate the activity of the polypeptides preferably include, but are not limited to, antisense oligonucleotides, agonists and antagonists, and inhibitors of protein kinases.

#### 25 **Methods of inducing immune response**

Another aspect of the present invention is directed to methods of inducing an immune response in a mammal against a polypeptide of the invention by administering to the mammal an amount of the polypeptide sufficient to induce an immune response. The amount will be dependent on the animal species, size of the animal, and the like but can be  
30 determined by those skilled in the art.

#### **Methods of identifying ligands**

The invention also provides assays to identify compounds that bind nGPCR-x. One such assay comprises the steps of: (a) contacting a composition comprising a nGPCR-x with a compound suspected of binding nGPCR-x; and (b) measuring binding between the compound and nGPCR-x. In one variation, the composition comprises a cell  
5 expressing nGPCR-x on its surface. In another variation, isolated nGPCR-x or cell membranes comprising nGPCR-x are employed. The binding may be measured directly, *e.g.*, by using a labeled compound, or may be measured indirectly by several techniques, including measuring intracellular signaling of nGPCR-x induced by the compound (or measuring changes in the level of nGPCR-x signaling). Following steps (a) and (b),  
10 compounds identified as binding nGPCR-x may be tested in other assays including, but not limited to, *in vivo* models, to confirm or quantitate binding to nGPCR-x.

Specific binding molecules, including natural ligands and synthetic compounds, can be identified or developed using isolated or recombinant nGPCR-x products, nGPCR-x variants, or preferably, cells expressing such products. Binding partners are useful for  
15 purifying nGPCR-x products and detection or quantification of nGPCR-x products in fluid and tissue samples using known immunological procedures. Binding molecules are also manifestly useful in modulating (*i.e.*, blocking, inhibiting or stimulating) biological activities of nGPCR-x, especially those activities involved in signal transduction.

The DNA and amino acid sequence information provided by the present invention  
20 also makes possible identification of binding partner compounds with which a nGPCR-x polypeptide or polynucleotide will interact. Methods to identify binding partner compounds include solution assays, *in vitro* assays wherein nGPCR-x polypeptides are immobilized, and cell-based assays. Identification of binding partner compounds of nGPCR-x polypeptides provides candidates for therapeutic or prophylactic intervention in  
25 pathologies associated with nGPCR-x normal and aberrant biological activity.

The invention includes several assay systems for identifying nGPCR-x binding partners. In solution assays, methods of the invention comprise the steps of (a) contacting a nGPCR-x polypeptide with one or more candidate binding partner compounds and (b) identifying the compounds that bind to the nGPCR-x polypeptide. Identification of the  
30 compounds that bind the nGPCR-x polypeptide can be achieved by isolating the nGPCR-x polypeptide/binding partner complex, and separating the binding partner compound from



the nGPCR-x polypeptide. An additional step of characterizing the physical, biological, and/or biochemical properties of the binding partner compound is also comprehended in another embodiment of the invention, wherein compounds identified as binding nGPCR-x may be tested in other assays including, but not limited to, *in vivo* models, to confirm or  
5 quantitate binding to nGPCR-x. In one aspect, the nGPCR-x polypeptide/binding partner complex is isolated using an antibody immunospecific for either the nGPCR-x polypeptide or the candidate binding partner compound.

In still other embodiments, either the nGPCR-x polypeptide or the candidate binding partner compound comprises a label or tag that facilitates its isolation, and  
10 methods of the invention to identify binding partner compounds include a step of isolating the nGPCR-x polypeptide/binding partner complex through interaction with the label or tag. An exemplary tag of this type is a poly-histidine sequence, generally around six histidine residues, that permits isolation of a compound so labeled using nickel chelation. Other labels and tags, such as the FLAG<sup>®</sup> tag (Eastman Kodak, Rochester, NY), well  
15 known and routinely used in the art, are embraced by the invention.

In one variation of an *in vitro* assay, the invention provides a method comprising the steps of (a) contacting an immobilized nGPCR-x polypeptide with a candidate binding partner compound and (b) detecting binding of the candidate compound to the nGPCR-x polypeptide. In an alternative embodiment, the candidate binding partner compound is  
20 immobilized and binding of nGPCR-x is detected. Immobilization is accomplished using any of the methods well known in the art, including covalent bonding to a support, a bead, or a chromatographic resin, as well as non-covalent, high affinity interactions such as antibody binding, or use of streptavidin/biotin binding wherein the immobilized compound includes a biotin moiety. Detection of binding can be accomplished (i) using a radioactive  
25 label on the compound that is not immobilized, (ii) using of a fluorescent label on the non-immobilized compound, (iii) using an antibody immunospecific for the non-immobilized compound, (iv) using a label on the non-immobilized compound that excites a fluorescent support to which the immobilized compound is attached, as well as other techniques well known and routinely practiced in the art.

30 The invention also provides cell-based assays to identify binding partner compounds of a nGPCR-x polypeptide. In one embodiment, the invention provides a

method comprising the steps of contacting a nGPCR-x polypeptide expressed on the surface of a cell with a candidate binding partner compound and detecting binding of the candidate binding partner compound to the nGPCR-x polypeptide. In a preferred embodiment, the detection comprises detecting a calcium flux or other physiological event  
5 in the cell caused by the binding of the molecule.

Another aspect of the present invention is directed to methods of identifying compounds that bind to either nGPCR-x or nucleic acid molecules encoding nGPCR-x, comprising contacting nGPCR-x, or a nucleic acid molecule encoding the same, with a compound, and determining whether the compound binds nGPCR-x or a nucleic acid  
10 molecule encoding the same. Binding can be determined by binding assays which are well known to the skilled artisan, including, but not limited to, gel-shift assays, Western blots, radiolabeled competition assay, phage-based expression cloning, co-fractionation by chromatography, co-precipitation, cross linking, interaction trap/two-hybrid analysis, southwestern analysis, ELISA, and the like, which are described in, for example, *Current*  
15 *Protocols in Molecular Biology*, 1999, John Wiley & Sons, NY, which is incorporated herein by reference in its entirety. The compounds to be screened include (which may include compounds which are suspected to bind nGPCR-x, or a nucleic acid molecule encoding the same), but are not limited to, extracellular, intracellular, biologic or chemical origin. The methods of the invention also embrace ligands, especially neuropeptides, that  
20 are attached to a label, such as a radiolabel (e.g.,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^3\text{H}$ ), a fluorescence label, a chemiluminescent label, an enzymic label and an immunogenic label. Modulators falling within the scope of the invention include, but are not limited to, non-peptide molecules such as non-peptide mimetics, non-peptide allosteric effectors, and peptides. The nGPCR-x polypeptide or polynucleotide employed in such a test may either be free in  
25 solution, attached to a solid support, borne on a cell surface or located intracellularly or associated with a portion of a cell. One skilled in the art can, for example, measure the formation of complexes between nGPCR-x and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between nGPCR-x and its substrate caused by the compound being tested.

30 In another embodiment of the invention, high throughput screening for compounds having suitable binding affinity to nGPCR-x is employed. Briefly, large numbers of

different test compounds are synthesized on a solid substrate. The peptide test compounds are contacted with nGPCR-x and washed. Bound nGPCR-x is then detected by methods well known in the art. Purified polypeptides of the invention can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-  
5 neutralizing antibodies can be used to capture the protein and immobilize it on the solid support.

Generally, an expressed nGPCR-x can be used for HTS binding assays in conjunction with its defined ligand, in this case the corresponding neuropeptide that activates it. The identified peptide is labeled with a suitable radioisotope, including, but  
10 not limited to,  $^{125}\text{I}$ ,  $^3\text{H}$ ,  $^{35}\text{S}$  or  $^{32}\text{P}$ , by methods that are well known to those skilled in the art. Alternatively, the peptides may be labeled by well-known methods with a suitable fluorescent derivative (Baindur *et al.*, *Drug Dev. Res.*, 1994, 33, 373-398; Rogers, *Drug Discovery Today*, 1997, 2, 156-160). Radioactive ligand specifically bound to the receptor in membrane preparations made from the cell line expressing the recombinant protein can  
15 be detected in HTS assays in one of several standard ways, including filtration of the receptor-ligand complex to separate bound ligand from unbound ligand (Williams, *Med. Res. Rev.*, 1991, 11, 147-184; Sweetnam *et al.*, *J. Natural Products*, 1993, 56, 441-455). Alternative methods include a scintillation proximity assay (SPA) or a FlashPlate format in which such separation is unnecessary (Nakayama, *Cur. Opinion Drug Disc. Dev.*, 1998,  
20 1, 85-91 Bossé *et al.*, *J. Biomolecular Screening*, 1998, 3, 285-292.). Binding of fluorescent ligands can be detected in various ways, including fluorescence energy transfer (FRET), direct spectrophotofluorometric analysis of bound ligand, or fluorescence polarization (Rogers, *Drug Discovery Today*, 1997, 2, 156-160; Hill, *Cur. Opinion Drug Disc. Dev.*, 1998, 1, 92-97).

25 Other assays may be used to identify specific ligands of a nGPCR-x receptor, including assays that identify ligands of the target protein through measuring direct binding of test ligands to the target protein, as well as assays that identify ligands of target proteins through affinity ultrafiltration with ion spray mass spectroscopy/HPLC methods or other physical and analytical methods. Alternatively, such binding interactions are  
30 evaluated indirectly using the yeast two-hybrid system described in Fields *et al.*, *Nature*, 340:245-246 (1989), and Fields *et al.*, *Trends in Genetics*, 10:286-292 (1994), both of

which are incorporated herein by reference. The two-hybrid system is a genetic assay for detecting interactions between two proteins or polypeptides. It can be used to identify proteins that bind to a known protein of interest, or to delineate domains or residues critical for an interaction. Variations on this methodology have been developed to clone  
5 genes that encode DNA binding proteins, to identify peptides that bind to a protein, and to screen for drugs. The two-hybrid system exploits the ability of a pair of interacting proteins to bring a transcription activation domain into close proximity with a DNA binding domain that binds to an upstream activation sequence (UAS) of a reporter gene, and is generally performed in yeast. The assay requires the construction of two hybrid  
10 genes encoding (1) a DNA-binding domain that is fused to a first protein and (2) an activation domain fused to a second protein. The DNA-binding domain targets the first hybrid protein to the UAS of the reporter gene; however, because most proteins lack an activation domain, this DNA-binding hybrid protein does not activate transcription of the reporter gene. The second hybrid protein, which contains the activation domain, cannot  
15 by itself activate expression of the reporter gene because it does not bind the UAS. However, when both hybrid proteins are present, the noncovalent interaction of the first and second proteins tethers the activation domain to the UAS, activating transcription of the reporter gene. For example, when the first protein is a GPCR gene product, or fragment thereof, that is known to interact with another protein or nucleic acid, this assay  
20 can be used to detect agents that interfere with the binding interaction. Expression of the reporter gene is monitored as different test agents are added to the system. The presence of an inhibitory agent results in lack of a reporter signal.

The yeast two-hybrid assay can also be used to identify proteins that bind to the gene product. In an assay to identify proteins that bind to a nGPCR-x receptor, or  
25 fragment thereof, a fusion polynucleotide encoding both a nGPCR-x receptor (or fragment) and a UAS binding domain (*i.e.*, a first protein) may be used. In addition, a large number of hybrid genes each encoding a different second protein fused to an activation domain are produced and screened in the assay. Typically, the second protein is encoded by one or more members of a total cDNA or genomic DNA fusion library, with  
30 each second protein-coding region being fused to the activation domain. This system is applicable to a wide variety of proteins, and it is not even necessary to know the identity

or function of the second binding protein. The system is highly sensitive and can detect interactions not revealed by other methods; even transient interactions may trigger transcription to produce a stable mRNA that can be repeatedly translated to yield the reporter protein.

5        Other assays may be used to search for agents that bind to the target protein. One such screening method to identify direct binding of test ligands to a target protein is described in U.S. Patent No. 5,585,277, incorporated herein by reference. This method relies on the principle that proteins generally exist as a mixture of folded and unfolded states, and continually alternate between the two states. When a test ligand binds to the folded form of a target protein (*i.e.*, when the test ligand is a ligand of the target protein),  
10        the target protein molecule bound by the ligand remains in its folded state. Thus, the folded target protein is present to a greater extent in the presence of a test ligand which binds the target protein, than in the absence of a ligand. Binding of the ligand to the target protein can be determined by any method that distinguishes between the folded and unfolded states of the target protein. The function of the target protein need not be known  
15        in order for this assay to be performed. Virtually any agent can be assessed by this method as a test ligand, including, but not limited to, metals, polypeptides, proteins, lipids, polysaccharides, polynucleotides and small organic molecules.

Another method for identifying ligands of a target protein is described in Wieboldt  
20        *et al.*, Anal. Chem., 69:1683-1691 (1997), incorporated herein by reference. This technique screens combinatorial libraries of 20-30 agents at a time in solution phase for binding to the target protein. Agents that bind to the target protein are separated from other library components by simple membrane washing. The specifically selected molecules that are retained on the filter are subsequently liberated from the target protein  
25        and analyzed by HPLC and pneumatically assisted electrospray (ion spray) ionization mass spectroscopy. This procedure selects library components with the greatest affinity for the target protein, and is particularly useful for small molecule libraries.

Other embodiments of the invention comprise using competitive screening assays in which neutralizing antibodies capable of binding a polypeptide of the invention  
30        specifically compete with a test compound for binding to the polypeptide. In this manner, the antibodies can be used to detect the presence of any peptide that shares one or more

antigenic determinants with nGPCR-x. Radiolabeled competitive binding studies are described in A.H. Lin *et al. Antimicrobial Agents and Chemotherapy*, 1997, vol. 41, no. 10, pp. 2127-2131, the disclosure of which is incorporated herein by reference in its entirety.

5 As described above and in Example 5 below, the gene encoding nGPCR-74 (nucleic acid sequence SEQ ID NO:134, amino acid sequence SEQ ID NO:268) has been detected in brain tissue indicating that this nGPCR protein is a neuroreceptor. It is well known that other nGPCR-x are expressed in many different tissues, including the brain. Accordingly, natural binding partners of these molecules include neurotransmitters.

#### 10 Identification of modulating agents

The invention also provides methods for identifying a modulator of binding between a nGPCR-x and a nGPCR-x binding partner, comprising the steps of: (a) contacting a nGPCR-x binding partner and a composition comprising a nGPCR-x in the presence and in the absence of a putative modulator compound; (b) detecting binding  
15 between the binding partner and the nGPCR-x; and (c) identifying a putative modulator compound or a modulator compound in view of decreased or increased binding between the binding partner and the nGPCR-x in the presence of the putative modulator, as compared to binding in the absence of the putative modulator. Following steps (a) and (b), compounds identified as modulating binding between nGPCR-x and a nGPCR-x binding  
20 partner may be tested in other assays including, but not limited to, *in vivo* models, to confirm or quantitate modulation of binding to nGPCR-x.

nGPCR-x binding partners that stimulate nGPCR-x activity are useful as agonists in disease states or conditions characterized by insufficient nGPCR-x signaling (*e.g.*, as a result of insufficient activity of a nGPCR-x ligand). nGPCR-x binding partners that block  
25 ligand-mediated nGPCR-x signaling are useful as nGPCR-x antagonists to treat disease states or conditions characterized by excessive nGPCR-x signaling. In addition nGPCR-x modulators in general, as well as nGPCR-x polynucleotides and polypeptides, are useful in diagnostic assays for such diseases or conditions.

In another aspect, the invention provides methods for treating a disease or  
30 abnormal condition by administering to a patient in need of such treatment a substance

that modulates the activity or expression of a polypeptide having sequences selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268.

Agents that modulate (*i.e.*, increase, decrease, or block) nGPCR-x activity or expression may be identified by incubating a putative modulator with a cell containing a nGPCR-x polypeptide or polynucleotide and determining the effect of the putative modulator on nGPCR-x activity or expression. The selectivity of a compound that modulates the activity of nGPCR-x can be evaluated by comparing its effects on nGPCR-x to its effect on other GPCR compounds. Following identification of compounds that modulate nGPCR-x activity or expression, such compounds may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity. Selective modulators may include, for example, antibodies and other proteins, peptides, or organic molecules that specifically bind to a nGPCR-x polypeptide or a nGPCR-x-encoding nucleic acid. Modulators of nGPCR-x activity will be therapeutically useful in treatment of diseases and physiological conditions in which normal or aberrant nGPCR-x activity is involved. nGPCR-x polynucleotides, polypeptides, and modulators may be used in the treatment of such diseases and conditions as infections, such as viral infections caused by HIV-1 or HIV-2; pain; cancers; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); Parkinson's disease; and psychotic and neurological disorders, including schizophrenia, migraine, ADHH, major depression, anxiety, mental disorder, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Tourette's Syndrome, among others. nGPCR-x polynucleotides and polypeptides, as well as nGPCR-x modulators, may also be used in diagnostic assays for such diseases or conditions.

Methods of the invention to identify modulators include variations on any of the methods described above to identify binding partner compounds, the variations including techniques wherein a binding partner compound has been identified and the binding assay is carried out in the presence and absence of a candidate modulator. A modulator is identified in those instances where binding between the nGPCR-x polypeptide and the binding partner compound changes in the presence of the candidate modulator compared

to binding in the absence of the candidate modulator compound. A modulator that increases binding between the nGPCR-x polypeptide and the binding partner compound is described as an enhancer or activator, and a modulator that decreases binding between the nGPCR-x polypeptide and the binding partner compound is described as an inhibitor.

5 Following identification of modulators, such compounds may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity as modulators.

The invention also comprehends high-throughput screening (HTS) assays to identify compounds that interact with or inhibit biological activity (*i.e.*, affect enzymatic activity, binding activity, *etc.*) of a nGPCR-x polypeptide. HTS assays permit screening

10 of large numbers of compounds in an efficient manner. Cell-based HTS systems are contemplated to investigate nGPCR-x receptor-ligand interaction. HTS assays are designed to identify "hits" or "lead compounds" having the desired property, from which modifications can be designed to improve the desired property. Chemical modification of

15 the "hit" or "lead compound" is often based on an identifiable structure/activity relationship between the "hit" and the nGPCR-x polypeptide.

Another aspect of the present invention is directed to methods of identifying compounds which modulate (*i.e.*, increase or decrease) an activity of nGPCR-x comprising contacting nGPCR-x with a compound, and determining whether the

20 compound modifies activity of nGPCR-x. The activity in the presence of the test compared is measured to the activity in the absence of the test compound. Where the activity of the sample containing the test compound is higher than the activity in the sample lacking the test compound, the compound will have increased activity. Similarly, where the activity of the sample containing the test compound is lower than the activity in

25 the sample lacking the test compound, the compound will have inhibited activity. Following the identification of compounds that modulate an activity of nGPCR-x, such compounds can be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity.

The present invention is particularly useful for screening compounds by using

30 nGPCR-x in any of a variety of drug screening techniques. The compounds to be screened include (which may include compounds which are suspected to modulate nGPCR-x



activity), but are not limited to, extracellular, intracellular, biologic or chemical origin. The nGPCR-x polypeptide employed in such a test may be in any form, preferably, free in solution, attached to a solid support, borne on a cell surface or located intracellularly. One skilled in the art can, for example, measure the formation of complexes between nGPCR-x  
5 and the compound being tested. Alternatively, one skilled in the art can examine the diminution in complex formation between nGPCR-x and its substrate caused by the compound being tested.

The activity of nGPCR-x polypeptides of the invention can be determined by, for example, examining the ability to bind or be activated by chemically synthesized peptide  
10 ligands. Alternatively, the activity of nGPCR-x polypeptides can be assayed by examining their ability to bind calcium ions, hormones, chemokines, neuropeptides, neurotransmitters, nucleotides, lipids, odorants, and photons. Alternatively, the activity of the nGPCR-x polypeptides can be determined by examining the activity of effector molecules including, but not limited to, adenylate cyclase, phospholipases and ion  
15 channels. Thus, modulators of nGPCR-x polypeptide activity may alter a GPCR receptor function, such as a binding property of a receptor or an activity such as G protein-mediated signal transduction or membrane localization. In various embodiments of the method, the assay may take the form of an ion flux assay, a yeast growth assay, a non-hydrolyzable GTP assay such as a [<sup>35</sup>S]-GTP  $\gamma$ S assay, a cAMP assay, an inositol  
20 triphosphate assay, a diacylglycerol assay, an Aequorin assay, a Luciferase assay, a FLIPR assay for intracellular Ca<sup>2+</sup> concentration, a mitogenesis assay, a MAP Kinase activity assay, an arachidonic acid release assay (e.g., using [<sup>3</sup>H]-arachidonic acid), and an assay for extracellular acidification rates, as well as other binding or function-based assays of nGPCR-x activity that are generally known in the art. In several of these embodiments,  
25 the invention comprehends the inclusion of any of the G proteins known in the art, such as G<sub>16</sub>, G<sub>15</sub>, or chimeric G<sub>q45</sub>, G<sub>qs5</sub>, G<sub>qo5</sub>, G<sub>q25</sub>, and the like. nGPCR-x activity can be determined by methodologies that are used to assay for FaRP activity, which is well known to those skilled in the art. Biological activities of nGPCR-x receptors according to the invention include, but are not limited to, the binding of a natural or an unnatural  
30 ligand, as well as any one of the functional activities of GPCRs known in the art. Non-limiting examples of GPCR activities include transmembrane signaling of various forms,

which may involve G protein association and/or the exertion of an influence over G protein binding of various guanidylate nucleotides; another exemplary activity of GPCRs is the binding of accessory proteins or polypeptides that differ from known G proteins.

The modulators of the invention exhibit a variety of chemical structures, which can be generally grouped into non-peptide mimetics of natural GPCR receptor ligands, peptide and non-peptide allosteric effectors of GPCR receptors, and peptides that may function as activators or inhibitors (competitive, uncompetitive and non-competitive) (e.g., antibody products) of GPCR receptors. The invention does not restrict the sources for suitable modulators, which may be obtained from natural sources such as plant, animal or mineral extracts, or non-natural sources such as small molecule libraries, including the products of combinatorial chemical approaches to library construction, and peptide libraries. Examples of peptide modulators of GPCR receptors exhibit the following primary structures: GLGPRPLRFamide, GNSFLRFamide, GGPQGPLRFamide, GPSGPLRFamide, PDVDHVFLRFamide, and pyro-EDVDHVFLRFamide.

Other assays can be used to examine enzymatic activity including, but not limited to, photometric, radiometric, HPLC, electrochemical, and the like, which are described in, for example, *Enzyme Assays: A Practical Approach*, eds. R. Eisinger and M. J. Danson, 1992, Oxford University Press, which is incorporated herein by reference in its entirety.

The use of cDNAs encoding GPCRs in drug discovery programs is well-known; assays capable of testing thousands of unknown compounds per day in high-throughput screens (HTSs) are thoroughly documented. The literature is replete with examples of the use of radiolabeled ligands in HTS binding assays for drug discovery (see Williams, *Medicinal Research Reviews*, 1991, 11, 147-184.; Sweetnam, et al., *J. Natural Products*, 1993, 56, 441-455 for review). Recombinant receptors are preferred for binding assay HTS because they allow for better specificity (higher relative purity), provide the ability to generate large amounts of receptor material, and can be used in a broad variety of formats (see Hodgson, *Bio/Technology*, 1992, 10, 973-980; each of which is incorporated herein by reference in its entirety).

A variety of heterologous systems is available for functional expression of recombinant receptors that are well known to those skilled in the art. Such systems include bacteria (Strosberg, et al., *Trends in Pharmacological Sciences*, 1992, 13, 95-98),

yeast (Pausch, *Trends in Biotechnology*, 1997, 15, 487-494), several kinds of insect cells (Vanden Broeck, *Int. Rev. Cytology*, 1996, 164, 189-268), amphibian cells (Jayawickreme *et al.*, *Current Opinion in Biotechnology*, 1997, 8, 629-634) and several mammalian cell lines (CHO, HEK-293, COS, etc.; see Gerhardt, *et al.*, *Eur. J. Pharmacology*, 1997, 334, 1-23). These examples do not preclude the use of other possible cell expression systems, including cell lines obtained from nematodes (PCT application WO 98/37177).

In preferred embodiments of the invention, methods of screening for compounds that modulate nGPCR-x activity comprise contacting test compounds with nGPCR-x and assaying for the presence of a complex between the compound and nGPCR-x. In such assays, the ligand is typically labeled. After suitable incubation, free ligand is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular compound to bind to nGPCR-x.

It is well known that activation of heterologous receptors expressed in recombinant systems results in a variety of biological responses, which are mediated by G proteins expressed in the host cells. Occupation of a GPCR by an agonist results in exchange of bound GDP for GTP at a binding site on the  $G_\alpha$  subunit; one can use a radioactive, non-hydrolyzable derivative of GTP,  $GTP\gamma[^{35}S]$ , to measure binding of an agonist to the receptor (Sim *et al.*, *Neuroreport*, 1996, 7, 729-733). One can also use this binding to measure the ability of antagonists to bind to the receptor by decreasing binding of  $GTP\gamma[^{35}S]$  in the presence of a known agonist. One could therefore construct a HTS based on  $GTP\gamma[^{35}S]$  binding, though this is not the preferred method.

The G proteins required for functional expression of heterologous GPCRs can be native constituents of the host cell or can be introduced through well-known recombinant technology. The G proteins can be intact or chimeric. Often, a nearly universally competent G protein (e.g.,  $G_{\alpha 16}$ ) is used to couple any given receptor to a detectable response pathway. G protein activation results in the stimulation or inhibition of other native proteins, events that can be linked to a measurable response.

Examples of such biological responses include, but are not limited to, the following: the ability to survive in the absence of a limiting nutrient in specifically engineered yeast cells (Pausch, *Trends in Biotechnology*, 1997, 15, 487-494); changes in intracellular  $Ca^{2+}$  concentration as measured by fluorescent dyes (Murphy, *et al.*, *Cur.*

*Opinion Drug Disc. Dev.*, 1998, 1, 192-199). Fluorescence changes can also be used to monitor ligand-induced changes in membrane potential or intracellular pH; an automated system suitable for HTS has been described for these purposes (Schroeder, *et al.*, *J. Biomolecular Screening*, 1996, 1, 75-80). Melanophores prepared from *Xenopus laevis* show a ligand-dependent change in pigment organization in response to heterologous GPCR activation; this response is adaptable to HTS formats (Jayawickreme *et al.*, *Cur. Opinion Biotechnology*, 1997, 8, 629-634). Assays are also available for the measurement of common second messengers, including cAMP, phosphoinositides and arachidonic acid, but these are not generally preferred for HTS.

Preferred methods of HTS employing these receptors include permanently transfected CHO cells, in which agonists and antagonists can be identified by the ability to specifically alter the binding of GTP $\gamma$ [ $^{35}$ S] in membranes prepared from these cells. In another embodiment of the invention, permanently transfected CHO cells could be used for the preparation of membranes which contain significant amounts of the recombinant receptor proteins; these membrane preparations would then be used in receptor binding assays, employing the radiolabeled ligand specific for the particular receptor. Alternatively, a functional assay, such as fluorescent monitoring of ligand-induced changes in internal Ca $^{2+}$  concentration or membrane potential in permanently transfected CHO cells containing each of these receptors individually or in combination would be preferred for HTS. Equally preferred would be an alternative type of mammalian cell, such as HEK-293 or COS cells, in similar formats. More preferred would be permanently transfected insect cell lines, such as *Drosophila* S2 cells. Even more preferred would be recombinant yeast cells expressing the *Drosophila melanogaster* receptors in HTS formats well known to those skilled in the art (*e.g.*, Pausch, *Trends in Biotechnology*, 1997, 15, 487-494).

The invention contemplates a multitude of assays to screen and identify inhibitors of ligand binding to nGPCR-x receptors. In one example, the nGPCR-x receptor is immobilized and interaction with a binding partner is assessed in the presence and absence of a candidate modulator such as an inhibitor compound. In another example, interaction between the nGPCR-x receptor and its binding partner is assessed in a solution assay, both in the presence and absence of a candidate inhibitor compound. In either assay, an

inhibitor is identified as a compound that decreases binding between the nGPCR-x receptor and its binding partner. Following the identification of compounds which inhibit ligand binding to nGPCR-x receptors, such compounds may be further tested in other assays including, but not limited to, *in vivo* models, in order to confirm or quantitate their activity. Another contemplated assay involves a variation of the dihybrid assay wherein an inhibitor of protein/protein interactions is identified by detection of a positive signal in a transformed or transfected host cell, as described in PCT publication number WO 95/20652, published August 3, 1995.

Candidate modulators contemplated by the invention include compounds selected from libraries of either potential activators or potential inhibitors. There are a number of different libraries used for the identification of small molecule modulators, including: (1) chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules. Chemical libraries consist of random chemical structures, some of which are analogs of known compounds or analogs of compounds that have been identified as "hits" or "leads" in other drug discovery screens, some of which are derived from natural products, and some of which arise from non-directed synthetic organic chemistry. Natural product libraries are collections of microorganisms, animals, plants, or marine organisms which are used to create mixtures for screening by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of plants or marine organisms. Natural product libraries include polyketides, non-ribosomal peptides, and variants (non-naturally occurring) thereof. For a review, see Science 282:63-68 (1998). Combinatorial libraries are composed of large numbers of peptides, oligonucleotides, or organic compounds as a mixture. These libraries are relatively easy to prepare by traditional automated synthesis methods, PCR, cloning, or proprietary synthetic methods. Of particular interest are non-peptide combinatorial libraries. Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of combinatorial chemistry and libraries created therefrom, see Myers, Curr. Opin. Biotechnol. 8:701-707 (1997). Identification of modulators through use of the various libraries described herein permits modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to modulate activity.

Still other candidate inhibitors contemplated by the invention can be designed and include soluble forms of binding partners, as well as such binding partners as chimeric, or fusion, proteins. A "binding partner" as used herein broadly encompasses non-peptide modulators, as well as such peptide modulators as neuropeptides other than natural  
5 ligands, antibodies, antibody fragments, and modified compounds comprising antibody domains that are immunospecific for the expression product of the identified nGPCR-x gene.

The polypeptides of the invention are employed as a research tool for identification, characterization and purification of interacting, regulatory proteins.  
10 Appropriate labels are incorporated into the polypeptides of the invention by various methods known in the art and the polypeptides are used to capture interacting molecules. For example, molecules are incubated with the labeled polypeptides, washed to remove unbound polypeptides, and the polypeptide complex is quantified. Data obtained using different concentrations of polypeptide are used to calculate values for the number,  
15 affinity, and association of polypeptide with the protein complex.

Labeled polypeptides are also useful as reagents for the purification of molecules with which the polypeptide interacts including, but not limited to, inhibitors. In one embodiment of affinity purification, a polypeptide is covalently coupled to a chromatography column. Cells and their membranes are extracted, and various cellular  
20 subcomponents are passed over the column. Molecules bind to the column by virtue of their affinity to the polypeptide. The polypeptide-complex is recovered from the column, dissociated and the recovered molecule is subjected to protein sequencing. This amino acid sequence is then used to identify the captured molecule or to design degenerate oligonucleotides for cloning the corresponding gene from an appropriate cDNA library.

25 Alternatively, compounds may be identified which exhibit similar properties to the ligand for the nGPCR-x of the invention, but which are smaller and exhibit a longer half time than the endogenous ligand in a human or animal body. When an organic compound is designed, a molecule according to the invention is used as a "lead" compound. The design of mimetics to known pharmaceutically active compounds is a well-known  
30 approach in the development of pharmaceuticals based on such "lead" compounds. Mimetic design, synthesis and testing are generally used to avoid randomly screening a

large number of molecules for a target property. Furthermore, structural data deriving from the analysis of the deduced amino acid sequences encoded by the DNAs of the present invention are useful to design new drugs, more specific and therefore with a higher pharmacological potency.

5        Comparison of the protein sequence of the present invention with the sequences present in all the available databases showed a significant homology with the transmembrane portion of G protein coupled receptors. Accordingly, computer modeling can be used to develop a putative tertiary structure of the proteins of the invention based on the available information of the transmembrane domain of other proteins. Thus, novel  
10    ligands based on the predicted structure of nGPCR-x can be designed.

      In a particular embodiment, the novel molecules identified by the screening methods according to the invention are low molecular weight organic molecules, in which case a composition or pharmaceutical composition can be prepared thereof for oral intake, such as in tablets. The compositions, or pharmaceutical compositions, comprising the  
15    nucleic acid molecules, vectors, polypeptides, antibodies and compounds identified by the screening methods described herein, can be prepared for any route of administration including, but not limited to, oral, intravenous, cutaneous, subcutaneous, nasal, intramuscular or intraperitoneal. The nature of the carrier or other ingredients will depend on the specific route of administration and particular embodiment of the invention to be  
20    administered. Examples of techniques and protocols that are useful in this context are, *inter alia*, found in Remington's Pharmaceutical Sciences, 16<sup>th</sup> edition, Osol, A (ed.), 1980, which is incorporated herein by reference in its entirety.

      The dosage of these low molecular weight compounds will depend on the disease state or condition to be treated and other clinical factors such as weight and condition of  
25    the human or animal and the route of administration of the compound. For treating human or animals, between approximately 0.5 mg/kg of body weight to 500 mg/kg of body weight of the compound can be administered. Therapy is typically administered at lower dosages and is continued until the desired therapeutic outcome is observed.

      The present compounds and methods, including nucleic acid molecules,  
30    polypeptides, antibodies, compounds identified by the screening methods described herein, have a variety of pharmaceutical applications and may be used, for example, to

treat or prevent unregulated cellular growth, such as cancer cell and tumor growth. In a particular embodiment, the present molecules are used in gene therapy. For a review of gene therapy procedures, see *e.g.* Anderson, *Science*, 1992, 256, 808-813, which is incorporated herein by reference in its entirety.

5       The present invention also encompasses a method of agonizing (stimulating) or antagonizing a nGPCR-x natural binding partner associated activity in a mammal comprising administering to said mammal an agonist or antagonist to one of the above disclosed polypeptides in an amount sufficient to effect said agonism or antagonism. One embodiment of the present invention, then, is a method of treating diseases in a mammal  
10 with an agonist or antagonist of the protein of the present invention comprises administering the agonist or antagonist to a mammal in an amount sufficient to agonize or antagonize nGPCR-x-associated functions.

In an effort to discover novel treatments for diseases, biomedical researchers and chemists have designed, synthesized, and tested molecules that modulate the function of G  
15 protein coupled receptors. Some small organic molecules form a class of compounds that modulate the function of G protein coupled receptors.

Exemplary diseases and conditions amenable to treatment based on the present invention include, but are not limited to, thyroid disorders (*e.g.*, thyrotoxicosis, myxoedema); renal failure; inflammatory conditions (*e.g.*, Crohn's disease); diseases  
20 related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (*e.g.*, pain including migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety disorder, post-traumatic-stress disorder, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease  
25 or Tourette's Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis,  
30 myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative



disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); sexual dysfunction, among others.

Methods of determining the dosages of compounds to be administered to a patient and modes of administering compounds to an organism are disclosed in U.S. Application  
5 Serial No. 08/702,282, filed August 23, 1996 and International patent publication number WO 96/22976, published August 1 1996, both of which are incorporated herein by reference in their entirety, including any drawings, figures or tables. Those skilled in the art will appreciate that such descriptions are applicable to the present invention and can be  
10 easily adapted to it.

The proper dosage depends on various factors such as the type of disease being treated, the particular composition being used and the size and physiological condition of the patient. Therapeutically effective doses for the compounds described herein can be estimated initially from cell culture and animal models. For example, a dose can be  
15 formulated in animal models to achieve a circulating concentration range that initially takes into account the  $IC_{50}$  as determined in cell culture assays. The animal model data can be used to more accurately determine useful doses in humans.

Plasma half-life and biodistribution of the drug and metabolites in the plasma, tumors and major organs can also be determined to facilitate the selection of drugs most  
20 appropriate to inhibit a disorder. Such measurements can be carried out. For example, HPLC analysis can be performed on the plasma of animals treated with the drug and the location of radiolabeled compounds can be determined using detection methods such as X-ray, CAT scan and MRI. Compounds that show potent inhibitory activity in the screening assays, but have poor pharmacokinetic characteristics, can be optimized by altering the  
25 chemical structure and retesting. In this regard, compounds displaying good pharmacokinetic characteristics can be used as a model.

Toxicity studies can also be carried out by measuring the blood cell composition. For example, toxicity studies can be carried out in a suitable animal model as follows: 1) the compound is administered to mice (an untreated control mouse should also be used);  
30 2) blood samples are periodically obtained via the tail vein from one mouse in each treatment group; and 3) the samples are analyzed for red and white blood cell counts,

blood cell composition and the percent of lymphocytes versus polymorphonuclear cells. A comparison of results for each dosing regime with the controls indicates if toxicity is present.

At the termination of each toxicity study, further studies can be carried out by sacrificing the animals (preferably, in accordance with the American Veterinary Medical Association guidelines Report of the American Veterinary Medical Assoc. Panel on Euthanasia, Journal of American Veterinary Medical Assoc., 202:229-249, 1993). Representative animals from each treatment group can then be examined by gross necropsy for immediate evidence of metastasis, unusual illness or toxicity. Gross abnormalities in tissue are noted and tissues are examined histologically. Compounds causing a reduction in body weight or blood components are less preferred, as are compounds having an adverse effect on major organs. In general, the greater the adverse effect the less preferred the compound.

For the treatment of many diseases, the expected daily dose of a hydrophobic pharmaceutical agent is between 1 to 500 mg/day, preferably 1 to 250 mg/day, and most preferably 1 to 50 mg/day. Drugs can be delivered less frequently provided plasma levels of the active moiety are sufficient to maintain therapeutic effectiveness. Plasma levels should reflect the potency of the drug. Generally, the more potent the compound the lower the plasma levels necessary to achieve efficacy.

As discussed above, it is well known that GPCRs are expressed in many different tissues and regions, including in the brain. nGPCR-x mRNA transcripts may be found in many other tissues, including, but not limited to peripheral blood lymphocytes, pancreas, ovary, uterus, testis, salivary gland, kidney, adrenal gland, liver, bone marrow, prostate, fetal liver, colon, muscle, and fetal brain, and may be found in many other tissues. Within the brain, nGPCR-x mRNA transcripts may be found in many tissues, including, but not limited to, frontal lobe, hypothalamus, pons, cerebellum, caudate nucleus, and medulla.

Sequences selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 will, as detailed above, enable screening the endogenous neurotransmitters/hormones/ligands which activate, agonize, or antagonize nGPCR-x and for compounds with potential utility in treating disorders including, but not limited to, thyroid disorders (e.g., thyrotoxicosis, myxoedema); renal failure; inflammatory

conditions (*e.g.*, Crohn's disease); diseases related to cell differentiation and homeostasis; rheumatoid arthritis; autoimmune disorders; movement disorders; CNS disorders (*e.g.*, pain including schizophrenia, migraine; stroke; psychotic and neurological disorders, including anxiety, mental disorder, manic depression, anxiety, generalized anxiety disorder, post-traumatic-stress disorder, depression, bipolar disorder, delirium, dementia, severe mental retardation; dyskinesias, such as Huntington's disease or Tourette's Syndrome; attention disorders including ADD and ADHD, and degenerative disorders such as Parkinson's, Alzheimer's; movement disorders, including ataxias, supranuclear palsy, *etc.*); infections, such as viral infections caused by HIV-1 or HIV-2; metabolic and cardiovascular diseases and disorders (*e.g.*, type 2 diabetes, impaired glucose tolerance, dyslipidemia, obesity, anorexia, hypotension, hypertension, thrombosis, myocardial infarction, cardiomyopathies, atherosclerosis, *etc.*); proliferative diseases and cancers (*e.g.*, different cancers such as breast, colon, lung, *etc.*, and hyperproliferative disorders such as psoriasis, prostate hyperplasia, *etc.*); hormonal disorders (*e.g.*, male/female hormonal replacement, polycystic ovarian syndrome, alopecia, *etc.*); sexual dysfunction, among others.

For example, nGPCR-x may be useful in the treatment of respiratory ailments such as asthma, where T cells are implicated by the disease. Contraction of airway smooth muscle is stimulated by thrombin. Cicala *et al* (1999) Br J Pharmacol 126:478-484. Additionally, in bronchiolitis obliterans, it has been noted that activation of thrombin receptors may be deleterious. Hauck *et al.* (1999) Am J Physiol 277:L22-L29. Furthermore, mast cells have also been shown to have thrombin receptors. Cirino *et al* (1996) J Exp Med 183:821-827. nGPCR-x may also be useful in remodeling of airway structures in chronic pulmonary inflammation via stimulation of fibroblast procollagen synthesis. See, *e.g.*, Chambers *et al.* (1998) Biochem J 333:121-127; Trejo *et al.* (1996) J Biol Chem 271:21536-21541.

In another example, increased release of sCD40L and expression of CD40L by T cells after activation of thrombin receptors suggests that nGPCR-x may be useful in the treatment of unstable angina due to the role of T cells and inflammation. See Aukrust *et al.* (1999) Circulation 100:614-620.

A further example is the treatment of inflammatory diseases, such as psoriasis, inflammatory bowel disease, multiple sclerosis, rheumatoid arthritis, and thyroiditis. Due to the tissue expression profile of nGPCR-x, inhibition of thrombin receptors may be beneficial for these diseases. See, *e.g.*, Morris *et al.* (1996) *Ann Rheum Dis* 55:841-843.

5 In addition to T cells, NK cells and monocytes are also critical cell types which contribute to the pathogenesis of these diseases. See, *e.g.*, Naldini & Carney (1996) *Cell Immunol* 172:35-42; Hoffman & Cooper (1995) *Blood Cells Mol Dis* 21:156-167; Colotta *et al.* (1994) *Am J Pathol* 144:975-985.

Expression of nGPCR-x in bone marrow and spleen may suggest that it may play a  
10 role in the proliferation of hematopoietic progenitor cells. See DiCuccio *et al.* (1996) *Exp Hematol* 24:914-918.

As another example, nGPCR-x may be useful in the treatment of acute and/or traumatic brain injury. Astrocytes have been demonstrated to express thrombin receptors. Activation of thrombin receptors may be involved in astrogliosis following brain injury.  
15 Therefore, inhibition of receptor activity may be beneficial for limiting neuroinflammation. Scar formation mediated by astrocytes may also be limited by inhibiting thrombin receptors. See, *e.g.*, Pindon *et al.* (1998) *Eur J Biochem* 255:766-774; Ubl & Reiser. (1997) *Glia* 21:361-369; Grabham & Cunningham (1995) *J Neurochem* 64:583-591.

20 nGPCR-x receptor activation may mediate neuronal and astrocyte apoptosis and prevention of neurite outgrowth. Inhibition would be beneficial in both chronic and acute brain injury. See, *e.g.*, Donovan *et al.* (1997) *J Neurosci* 17:5316-5326; Turgeon *et al.* (1998) *J Neurosci* 18:6882-6891; Smith-Swintosky *et al.* (1997) *J Neurochem* 69:1890-1896; Gill *et al.* (1998) *Brain Res* 797:321-327; Suidan *et al.* (1996) *Semin Thromb*  
25 *Hemost* 22:125-133.

The attached Sequence Listing contains the sequences of the polynucleotides and polypeptides of the invention and is incorporated herein by reference in its entirety. As described above and in Example 5 below, the gene encoding nGPCR-74 (nucleic acid sequence SEQ ID NO:134, amino acid sequence SEQ ID NO:268) has been detected in  
30 brain tissue indicating that this nGPCR protein is a neuroreceptor. The identification of modulators such as agonists and antagonists is therefore useful for the identification of

compounds useful to treat neurological diseases and disorders. Such neurological diseases and disorders, including but are not limited to, schizophrenia, affective disorders, ADHD/ADD (*i.e.*, Attention Deficit-Hyperactivity Disorder/Attention Deficit Disorder), and neural disorders such as Alzheimer's disease, Parkinson's disease, migraine, and  
5 senile dementia as well as depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like.

#### Methods of Screening Human Subjects

Thus in yet another embodiment, the invention provides genetic screening procedures that entail analyzing a person's genome -- in particular their alleles for the  
10 nGPCR-x of the invention -- to determine whether the individual possesses a genetic characteristic found in other individuals that are considered to be afflicted with, or at risk for, developing a mental disorder or disease of the brain that is suspected of having a hereditary component. For example, in one embodiment, the invention provides a method for determining a potential for developing a disorder affecting the brain in a human subject  
15 comprising the steps of analyzing the coding sequence of one or more nGPCR-x genes from the human subject; and determining development potential for the disorder in said human subject from the analyzing step.

More particularly, the invention provides a method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the  
20 steps of: (a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering the amino acid sequence, expression, or biological activity of at least one seven transmembrane receptor that is expressed in the brain, wherein the seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or an allelic variant thereof, and wherein  
25 the nucleic acid corresponds to the gene encoding the seven transmembrane receptor; and (b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of allele in the nucleic acid correlates with an increased risk of developing the disorder.

30 By "human subject" is meant any human being, human embryo, or human fetus. It will be apparent that methods of the present invention will be of particular interest to

individuals that have themselves been diagnosed with a disorder affecting the brain or have relatives that have been diagnosed with a disorder affecting the brain.

By "screening for an increased risk" is meant determination of whether a genetic variation exists in the human subject that correlates with a greater likelihood of developing a disorder affecting the brain than exists for the human population as a whole, or for a relevant racial or ethnic human sub-population to which the individual belongs. Both positive and negative determinations (i.e., determinations that a genetic predisposition marker is present or is absent) are intended to fall within the scope of screening methods of the invention. In preferred embodiments, the presence of a mutation altering the sequence or expression of at least one nGPCR-x seven transmembrane receptor allele in the nucleic acid is correlated with an increased risk of developing mental disorder, whereas the absence of such a mutation is reported as a negative determination.

The "assaying" step of the invention may involve any techniques available for analyzing nucleic acid to determine its characteristics, including but not limited to well-known techniques such as single-strand conformation polymorphism analysis (SSCP) [Orita *et al.*, *Proc Natl. Acad. Sci. USA*, 86: 2766-2770 (1989)]; heteroduplex analysis [White *et al.*, *Genomics*, 12: 301-306 (1992)]; denaturing gradient gel electrophoresis analysis [Fischer *et al.*, *Proc. Natl. Acad. Sci. USA*, 80: 1579-1583 (1983); and Riesner *et al.*, *Electrophoresis*, 10: 377-389 (1989)]; DNA sequencing; RNase cleavage [Myers *et al.*, *Science*, 230: 1242-1246 (1985)]; chemical cleavage of mismatch techniques [Rowley *et al.*, *Genomics*, 30: 574-582 (1995); and Roberts *et al.*, *Nucl. Acids Res.*, 25: 3377-3378 (1997)]; restriction fragment length polymorphism analysis; single nucleotide primer extension analysis [Shumaker *et al.*, *Hum. Mutat.*, 7: 346-354 (1996); and Pastinen *et al.*, *Genome Res.*, 7: 606-614 (1997)]; 5' nuclease assays [Pease *et al.*, *Proc. Natl. Acad. Sci. USA*, 91:5022-5026 (1994)]; DNA Microchip analysis [Ramsay, G., *Nature Biotechnology*, 16: 40-48 (1999); and Chee *et al.*, U.S. Patent No. 5,837,832]; and ligase chain reaction [Whiteley *et al.*, U.S. Patent No. 5,521,065]. [See generally, Schafer and Hawkins, *Nature Biotechnology*, 16: 33-39 (1998).] All of the foregoing documents are hereby incorporated by reference in their entirety.

Thus, in one preferred embodiment involving screening nGPCR-x sequences, for example, the assaying step comprises at least one procedure selected from the group

consisting of: (a) determining a nucleotide sequence of at least one codon of at least one nGPCR-x allele of the human subject; (b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; (c) performing a polynucleotide migration  
5 assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and (d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

10 In a highly preferred embodiment, the assaying involves sequencing of nucleic acid to determine nucleotide sequence thereof, using any available sequencing technique. [See, e.g., Sanger *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 74: 5463-5467 (1977) (dideoxy chain termination method); Mirzabekov, *TIBTECH*, 12: 27-32 (1994) (sequencing by hybridization); Drmanac *et al.*, *Nature Biotechnology*, 16: 54-58 (1998); U.S. Patent No.  
15 5,202,231; and *Science*, 260: 1649-1652 (1993) (sequencing by hybridization); Kieleczawa *et al.*, *Science*, 258: 1787-1791 (1992) (sequencing by primer walking); (Douglas *et al.*, *Biotechniques*, 14: 824-828 (1993) (Direct sequencing of PCR products); and Akane *et al.*, *Biotechniques* 16: 238-241 (1994); Maxam and Gilbert, *Meth. Enzymol.*, 65: 499-560 (1977) (chemical termination sequencing), all incorporated herein by  
20 reference.] The analysis may entail sequencing of the entire nGPCR gene genomic DNA sequence, or portions thereof; or sequencing of the entire seven transmembrane receptor coding sequence or portions thereof. In some circumstances, the analysis may involve a determination of whether an individual possesses a particular allelic variant, in which case sequencing of only a small portion of nucleic acid -- enough to determine the sequence of  
25 a particular codon characterizing the allelic variant -- is sufficient. This approach is appropriate, for example, when assaying to determine whether one family member inherited the same allelic variant that has been previously characterized for another family member, or, more generally, whether a person's genome contains an allelic variant that has been previously characterized and correlated with a mental disorder having a heritable  
30 component.

In another highly preferred embodiment, the assaying step comprises performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences. In a preferred embodiment, the hybridization involves a determination of whether nucleic acid  
5 derived from the human subject will hybridize with one or more oligonucleotides, wherein the oligonucleotides have nucleotide sequences that correspond identically to a portion of the nGPCR-x gene sequence taught herein, or that correspond identically except for one mismatch. The hybridization conditions are selected to differentiate between perfect sequence complementarity and imperfect matches differing by one or more bases. Such  
10 hybridization experiments thereby can provide single nucleotide polymorphism sequence information about the nucleic acid from the human subject, by virtue of knowing the sequences of the oligonucleotides used in the experiments.

Several of the techniques outlined above involve an analysis wherein one performs a polynucleotide migration assay, *e.g.*, on a polyacrylamide electrophoresis gel (or in a  
15 capillary electrophoresis system), under denaturing or non-denaturing conditions. Nucleic acid derived from the human subject is subjected to gel electrophoresis, usually adjacent to (or co-loaded with) one or more reference nucleic acids, such as reference GPCR-x encoding sequences having a coding sequence identical to all or a portion of SEQ ID NOS: 1 to 134 (or identical except for one known polymorphism). The nucleic acid from  
20 the human subject and the reference sequence(s) are subjected to similar chemical or enzymatic treatments and then electrophoresed under conditions whereby the polynucleotides will show a differential migration pattern, unless they contain identical sequences. [See generally Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, New York: John Wiley & Sons, Inc. (1987-1999); and Sambrook *et al.*, (eds.), *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor, New York: Cold Spring Harbor  
25 Laboratory Press (1989), both incorporated herein by reference in their entirety.]

In the context of assaying, the term "nucleic acid of a human subject" is intended to include nucleic acid obtained directly from the human subject (*e.g.*, DNA or RNA obtained from a biological sample such as a blood, tissue, or other cell or fluid sample);  
30 and also nucleic acid derived from nucleic acid obtained directly from the human subject. By way of non-limiting examples, well known procedures exist for creating cDNA that is



complementary to RNA derived from a biological sample from a human subject, and for amplifying (e.g., via polymerase chain reaction (PCR)) DNA or RNA derived from a biological sample obtained from a human subject. Any such derived polynucleotide which retains relevant nucleotide sequence information of the human subject's own DNA/RNA is intended to fall within the definition of "nucleic acid of a human subject" for the purposes of the present invention.

In the context of assaying, the term "mutation" includes addition, deletion, and/or substitution of one or more nucleotides in the GPCR gene sequence (e.g., as compared to the seven transmembrane receptor-encoding sequences set forth of SEQ ID NO:1 to SEQ ID NO:134, and other polymorphisms that occur in introns (where introns exist) and that are identifiable via sequencing, restriction fragment length polymorphism, or other techniques. The various activity examples provided herein permit determination of whether a mutation modulates activity of the relevant receptor in the presence or absence of various test substances.

In a related embodiment, the invention provides methods of screening a person's genotype with respect to the nGPCR-x of the invention, and correlating such genotypes with diagnoses for disease or with predisposition for disease (for genetic counseling). For example, the invention provides a method of screening for an nGPCR-x hereditary mental disorder genotype in a human patient, comprising the steps of: (a) providing a biological sample comprising nucleic acid from the patient, the nucleic acid including sequences corresponding to said patient's nGPCR-x alleles; (b) analyzing the nucleic acid for the presence of a mutation or mutations; (c) determining a nGPCR-x genotype from the analyzing step; and (d) correlating the presence of a mutation in an nGPCR-x allele with a hereditary mental disorder genotype. In a preferred embodiment, the biological sample is a cell sample containing human cells that contain genomic DNA of the human subject. The analyzing can be performed analogously to the assaying described in preceding paragraphs. For example, the analyzing comprises sequencing a portion of the nucleic acid (e.g., DNA or RNA), the portion comprising at least one codon of the nGPCR-x alleles.

Although more time consuming and expensive than methods involving nucleic acid analysis, the invention also may be practiced by assaying one or more proteins of a

human subject to determine the presence or absence of an amino acid sequence variation in GPCR protein from the human subject. Such protein analyses may be performed, e.g., by fragmenting GPCR protein via chemical or enzymatic methods and sequencing the resultant peptides; or by Western analyses using an antibody having specificity for a particular allelic variant of the GPCR.

The invention also provides materials that are useful for performing methods of the invention. For example, the present invention provides oligonucleotides useful as probes in the many analyzing techniques described above. In general, such oligonucleotide probes comprise 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides that have a sequence that is identical, or exactly complementary, to a portion of a human GPCR gene sequence taught herein (or allelic variant thereof), or that is identical or exactly complementary except for one nucleotide substitution. In a preferred embodiment, the oligonucleotides have a sequence that corresponds in the foregoing manner to a human GPCR coding sequence taught herein, and in particular, the coding sequences set forth in SEQ ID NO:1 to SEQ ID NO:134. In one variation, an oligonucleotide probe of the invention is purified and isolated. In another variation, the oligonucleotide probe is labeled, e.g., with a radioisotope, chromophore, or fluorophore. In yet another variation, the probe is covalently attached to a solid support. [See generally Ausubel *et al.* and Sambrook *et al.*, *supra*.]

In a related embodiment, the invention provides kits comprising reagents that are useful for practicing methods of the invention. For example, the invention provides a kit for screening a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association: (a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-x seven transmembrane receptor gene, the oligonucleotide comprising 6-50 nucleotides that have a sequence that is identical or exactly complementary to a portion of a human nGPCR-x gene sequence or nGPCR-x coding sequence, except for one sequence difference selected from the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and (b) a media packaged with the oligonucleotide containing information identifying polymorphisms identifiable with the probe that correlate with mental disorder or a genetic predisposition

therefor. Exemplary information-containing media include printed paper package inserts or packaging labels; and magnetic and optical storage media that are readable by computers or machines used by practitioners who perform genetic screening and counseling services. The practitioner uses the information provided in the media to correlate the results of the analysis with the oligonucleotide with a diagnosis. In a preferred variation, the oligonucleotide is labeled.

In still another embodiment, the invention provides methods of identifying those allelic variants of GPCRs of the invention that correlate with mental disorders. For example, the invention provides a method of identifying a seven transmembrane allelic variant that correlates with a mental disorder, comprising steps of: (a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny; (b) analyzing the nucleic acid for the presence of a mutation or mutations in at least one seven transmembrane receptor that is expressed in the brain, wherein the at least one seven transmembrane receptor comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 or an allelic variant thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding the at least one seven transmembrane receptor; (c) determining a genotype for the patient for the at least one seven transmembrane receptor from said analyzing step; and (d) identifying an allelic variant that correlates with the mental disorder from the determining step. To expedite this process, it may be desirable to perform linkage studies in the patients (and possibly their families) to correlate chromosomal markers with disease states. The chromosomal localization data provided herein facilitates identifying an involved nGPCR with a chromosomal marker.

The foregoing method can be performed to correlate the nGPCR-x of the invention to a number of disorders having hereditary components that are causative or that predispose persons to the disorder. For example, in one preferred variation, the disorder is a mental disorder.

Also contemplated as part of the invention are polynucleotides that comprise the allelic variant sequences identified by such methods, and polypeptides encoded by the allelic variant sequences, and oligonucleotide and oligopeptide fragments thereof that

embody the mutations that have been identified. Such materials are useful *in vitro* cell-free and cell-based assays for identifying lead compounds and therapeutics for treatment of the disorders. For example, the variants are used in activity assays, binding assays, and assays to screen for activity modulators described herein. In one preferred embodiment, the invention provides a purified and isolated polynucleotide comprising a nucleotide sequence encoding a nGPCR-x receptor allelic variant identified according to the methods described above; and an oligonucleotide that comprises the sequences that differentiate the allelic variant from the nGPCR-x sequences set forth in SEQ ID NO:1 to SEQ ID NO:134. The invention also provides a vector comprising the polynucleotide (preferably an expression vector); and a host cell transformed or transfected with the polynucleotide or vector. The invention also provides an isolated cell line that is expressing the allelic variant nGPCR-x polypeptide; purified cell membranes from such cells; purified polypeptide; and synthetic peptides that embody the allelic variation amino acid sequence. In one particular embodiment, the invention provides a purified polynucleotide comprising a nucleotide sequence encoding a nGPCR-x seven transmembrane receptor protein of a human that is affected with a mental disorder; wherein said polynucleotide hybridizes to the complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 under the following hybridization conditions: (a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and (b) washing 2 times for 30 minutes at 60°C in a wash solution comprising 0.1x SSC and 1% SDS; and wherein the polynucleotide encodes a nGPCR-x amino acid sequence that differs from a sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, by at least one residue.

An exemplary assay for using the allelic variants is a method for identifying a modulator of nGPCR-x biological activity, comprising the steps of: (a) contacting a cell expressing the allelic variant in the presence and in the absence of a putative modulator compound; (b) measuring nGPCR-x biological activity in the cell; and (c) identifying a putative modulator compound in view of decreased or increased nGPCR-x biological activity in the presence versus absence of the putative modulator.

Additional features of the invention will be apparent from the following Examples. Examples 1, 2, and portions of Examples 3 and 5 are actual, while the remaining

Examples are prophetic. Additional features and variations of the invention will be apparent to those skilled in the art from the entirety of this application, including the detailed description, and all such features are intended as aspects of the invention. Likewise, features of the invention described herein can be re-combined into additional  
5 embodiments that also are intended as aspects of the invention, irrespective of whether the combination of features is specifically mentioned above as an aspect or embodiment of the invention. Also, only such limitations which are described herein as critical to the invention should be viewed as such; variations of the invention lacking limitations which have not been described herein as critical are intended as aspects of the invention.

10

## EXAMPLES

### EXAMPLE 1: IDENTIFICATION OF nGPCR-X

#### A. Database search

The Celera database was searched using known GPCR receptors as query  
15 sequences to find patterns suggestive of novel G protein-coupled receptors. Positive hits were further analyzed with the GCG program BLAST to determine which ones were the most likely candidates to encode G protein-coupled receptors, using the standard (default) alignment produced by BLAST as a guide.

Briefly, the BLAST algorithm, which stands for Basic Local Alignment Search  
20 Tool is suitable for determining sequence similarity (Altschul *et al.*, J. Mol. Biol., 1990, 215, 403-410, which is incorporated herein by reference in its entirety). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in  
25 the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the  
30 cumulative alignment score can be increased. Extension for the word hits in each direction are halted when: 1) the cumulative alignment score falls off by the quantity X

from its maximum achieved value; 2) the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or 3) the end of either sequence is reached. The Blast algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The Blast program uses as defaults a word length  
5 (W) of 11, the BLOSUM62 scoring matrix (see Henikoff et al., Proc. Natl. Acad. Sci. USA, 1992, 89, 10915-10919, which is incorporated herein by reference in its entirety) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm (Karlin *et al.*, Proc. Natl. Acad. Sci. USA, 1993, 90, 5873-5787, which is incorporated herein by reference in its entirety) and Gapped BLAST  
10 perform a statistical analysis of the similarity between two sequences. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a GPCR gene or cDNA if the smallest sum probability in comparison of the test  
15 nucleic acid to a GPCR nucleic acid is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

Homology searches are performed with the program BLAST version 2.08. A collection of 340 query amino acid sequences derived from GPCRs was used to search the genomic DNA sequence using TBLASTN and alignments with an E-value lower than 0.01  
20 were collected from each BLAST search. The amino acid sequences have been edited to remove regions in the sequence that produce non-significant alignments with proteins that are not related to GPCRs.

Multiple query sequences may have a significant alignment to the same genomic region, although each alignment may not cover exactly the same DNA region. A  
25 procedure is used to determine the region of maximum common overlap between the alignments from several query sequences. This region is called the consensus DNA region. The procedure for determining this consensus involves the automatic parsing of the BLAST output files using the program MSPcrunch to produce a tabular report. From this tabular report the start and end of each alignment in the genomic DNA is extracted.  
30 This information is used by a PERL script to derive the maximum common overlap. These regions are reported in the form of a unique sequence identifier, a start and the end

position in the sequence. The sequences defined by these regions were extracted from the original genomic sequence file using the program fetchdb.

The consensus regions are assembled into a non-redundant set by using the program phrap. After assembly with phrap a set of contigs and singletons were defined as candidate DNA regions coding for nGPCRs. These sequences were then submitted for further sequence analysis.

Further sequence analysis involves the removal of sequences previously isolated and removal of sequences that are related to olfactory GPCR's.

nGPCR-x cDNAs were sequenced directly using an ABI377 fluorescence-based sequencer (Perkin-Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM<sup>TM</sup> Ready Dye-Deoxy Terminator kit with Taq FS<sup>TM</sup> polymerase. Each ABI cycle sequencing reaction contained about 0.5 µg of plasmid DNA. Cycle-sequencing was performed using an initial denaturation at 98°C for 1 minute, followed by 50 cycles using the following parameters: 98°C for 30 seconds, annealing at 50°C for 30 seconds, and extension at 60°C for 4 minutes. Temperature cycles and times were controlled by a Perkin-Elmer 9600 thermocycler. Extension products were purified using Centriflex<sup>TM</sup> gel filtration cartridges (Advanced Genetic Technologies Corp., Gaithersburg, MD). Each reaction product was loaded by pipette onto the column, which is then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B tabletop centrifuge) at 1500 x g for 4 minutes at room temperature. Column-purified samples were dried under vacuum for about 40 minutes and then dissolved in 5µl of a DNA loading solution (83% deionized formamide, 8.3mM EDTA, and 1.6 mg/ml Blue Dextran). The samples were then heated to 90°C for three minutes and loaded into the gel sample wells for sequence analysis using the ABI377 sequencer. Sequence analysis was performed by importing ABI377 files into the Sequencer program. (Gene Codes, Ann Arbor, MI). Generally, sequence reads of 700 bp were obtained. Potential sequencing errors were minimized by obtaining sequence information from both DNA strands and by re-sequencing difficult areas using primers annealing at different locations until all sequencing ambiguities were removed.

The following Table 5 contains the sequences of the polynucleotides and polypeptides of the invention. The transmembrane domains within the polypeptide sequence are identified by underlining.

TABLE 5
<p>The following DNA sequence Seq-2356 &lt;SEQ ID NO. 1&gt; was identified in <i>H. sapiens</i>:</p> <p>GGAATTTAGTTGGGCAGAAGGGGAATAAAGTGAGGATGGTTAATGGGTACAAAAATAGT  TAGGAAAAAATGAATAAGATCTAGTATTAGATAGCACACAGGGTGATTGTAGTCAATA  TAATTTAGTTGTACAATTTAAAATAACTAAAAGAATATAACTGGATTGTTGTAACACAA  ATGATAAACGCTTGAGGTAATGGATACGATATTTACCCTGATGTAATTATTACACATTGC  ACGCTGTATTCAAATAACCCATCTAACTCATAAATATTTATATCTACTATCTACACAA  AAAATTAATAAATAAAAAATTTTGCATGATGATCTTAAGTGAATTTTCAATAATAAA  ACATTGTCTGTTTTCATTAAGTTCAATTTAGCAATTTCAATTATGTTTAATTATTTTGC  ATCCTGAATAAAAAATCTTCTTATACTGCAAGATTTGAAGGCAATCTAGACTTACTTCT  AGAATTGTTATGTTCTACCTGTTATAATCAGGCTTACAATTCATGTCCAATTAATTTCA  TATGTAAAGTGAGTTATATTTTTCATGAAGTTGTTGAGTTTTCAGCCCCACTTAAAAAA  ATGTAGAATTGTTTCTGCTCAGTTAAACTGACCTGCTTTTT</p> <p>The following amino acid sequence &lt;SEQ ID NO. 135&gt; is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 1:</p> <p>KKQVSLTEQETILHFFKWKTEQLHEKYNSLYIKLIGHELALQVEHNNRSKSRLPKSCSIRFFIQDAK  IIKHNNCIELNENRQCFIEKFSDDHAKIFLIENFLCRIIFMSMGYFEYRRAMCNYIRVNIIVSITSSVYH  LCYKQSSYILLVILNCTTKLYLQSPCCAIYILFIFFLTIFCTHPSSLYSPSAQLNS</p>
<p>The following DNA sequence Seq-2357 SEQ ID NO. 2&gt; was identified in <i>H. sapiens</i>:</p> <p>CAGGTGCAGCATCGTGTCTCAGTGTCTGCCCCCTGCTTCCACCCGGTGTGACAGCTG  CACGGTCCACCCACGCTGCTTTCCATCGTTCTCATCAGCCCTGTGATCTTTCTCTGT  GGCCCTGCTGTGCTGGTGCCTGTGAGGTCTGTGGACACAAGAGACTGCACGGGCCACA  CCCCCAGCTGGGTGAGTCCTCTCCCTCCTGGGTACTCTGGACAGTAAAGAAAGATGGACA  CGTGGGCTCCGTGGAGCATGAGGTAGTCCAGGACCTCGGCGGCCACAGGTCTGCCTCCC  TGCTTCTCGTGCCTCCCTCCCTTTGGGTCTCTGCTCCACCTCGGTAAACGCTTCGTCC  CACCCCTCGAAGGGTAAATCGAGCTCCTTGGTGGTAAAGCACCCACTGCCCTAGTCAGA  GGGTCCCTCCTCTCTGATGTCTGGTGCCTGGTCTGCTGGTAGAATTTAGCTGCTTT  ATAACCTGGTCTGAAATGAACCACTGGGAAGAAATAGGGTAAATGAACACACAGCTGC  CACACTGCATCCCAACCCTGTGTGACCCTATCACCGCAGACTTTTGTGGCAAGATGACAG  CATCTCAGTTTGCTTGAGAAGCTTATTTTGGCAAGGCTGTTACCACCAGGCAGGCACCA</p> <p>The following amino acid sequence &lt;SEQ ID NO. 136&gt; is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 2:</p> <p>RCSIVSSVSCPLLPPGVDSCTVHPTPAFSPFLISPVIFPVALLCWCVPVRSCHGKRLHGPHPQLGESSPSWV  LWTVKKDGHVGSVEHEVVQDLGGHRSCLPASRALPPFGSLLHLGKRFVPTPRRVNRPWWSTHCPSEGPSS  LMSWCPGLPGRILAALPGPEMNHWEIEGNEHTAATLHPNPVPYHRRLLWQDDSIIVCLRSLFLPRLPPGR  H</p>
<p>The following DNA sequence Seq-2358 &lt;SEQ ID NO. 3&gt; was identified in <i>H. sapiens</i>:</p>



CTATTATTTCTTAACATACTGCATTTTCCGATTCTCTCTAAGTATCTGTTTCTGTAAC  
 CCTATTGGACATTTACTTCTCTTTTCACATTGTCTGCTTTATCTCTTAACTTTGTGTTT  
 CTGTCTCTCACTGCTGTATTGTGAGTTATTACTTAGCTCTCCAGTTAATTCTTTAAGCT  
 TTTTTTGAATCTCTTTATCAGTTCATTGTGTTTATTATTTAGTGACTAAATTTAATTG  
 CTCAAAGTTTTATTTGGTCGTTTAAAATTTGCTTTTGTCTTCATAGTTATTTGTCCTGT  
 TCTCTTTATCTCTTTATTTATTTTGTATGCTTTCATCTGCTTATTTATTTTAAGATATTT  
 ATTTCTTAGCCTCTTTGAGATATTCTATTATCTCTGGTCTAGGATCATTAATCTCCCA  
 CTACGTCTGTAGACTC

The following amino acid sequence <SEQ ID NO. 137> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 3:

IISHTAFFRFSLSICFCNSYWTFTSLSHCLLYLLTFVFSVSHCCIVSYYLALPVNSLSFFCNLFISSLCLL  
 FQLNLIAQSFWSFKICFLHSYFVLFSLSLYLFLMLSSAYYFDIYFLASLRYSIISGPRIIKSPPTSVD

The following DNA sequence Seq-2359 <SEQ ID NO. 4> was identified in *H. sapiens*:

ACTTCTGGGCCACGAAAGCCCTACTGTCTAAATGCTTTTCAGGCCAATTGAAGAAGTA  
 ATTAGACTTACTGGAAGCTTCTGTGAATAATTCTGCAAGTACAATTATGGACTTCCCAGG  
 AAATATTGCCCTCAATATAGAAAAGCTTGTCAAGTGAATCTGATGAGATATATGTAAAT  
 TTGAGATTTTGATATTAGAATGAGTAAAATGATGACATCACGATGTATTAAAGTTGGGGT  
 TTATTTTTTGAATTAATTGTTCATCAGGTAAAAGCCAGCTATAAGTCAAATAAAATATA  
 ATCATGTTCTTCCGTCTTAGCACTCATCTTTCTTGTCTAAATGTTGACAAATGACTG  
 TAAATTTAACAAGCTTATAGATAATAATTGAAAAGTCTTCTAAGAACTGAAAATTGATAA  
 ACACATGGCAATGGCAGGCTATTGCAAGTCAATTATAAGATGTTGTGTGGATGCCCTGA  
 AGTGCCATATAAATGAATGTGACTTCACTACTGCCAAATGAGTCCAATATCCCACAAA  
 TGAAGTGAATAAAGTGCCCTGGAATACTGTGTCTACAGTGTCACTGTAAAGTTACTGTC  
 ATGCTGTATTACTGAAATGATTTGCTGGAAGTAACATGGCACATATATGCACCAAGAGA  
 GTTAAATCTCATCTTATTCTATGAAAATCATGTTAACCATTTCATGA

The following amino acid sequence <SEQ ID NO. 138> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 4:

HEWLTFEIEILSWCIYVPCYFPANHFSNTAQLYSDTVDTVFQALYFQFICGILDSFGSSTEVTFIYRHF  
 RGIHTTSYNCTAIACHCHVFINFQFLEDFSIIYKLVKFTVICQHLEQEKMSAKDGRITLYFILIAGFLPDD  
 NFQKINPNFNFTSCHHFTHSNIKISNFTYISSESTDKLFYIEGNISWEVHNCTCRIIHRSFQVLLQLGLKS  
 ITVGLSVAQK

The following DNA sequence Seq-2360 <SEQ ID NO. 5> was identified in *H. sapiens*:

AACATTATTACTTTCTTTTATGAATATTCTTGGTCTTTCCAAAACAAAACAAGCTATGG  
 TTTAATAAATTATGGTATAATCAAATAATGAACTCTATGCATTGTTAAAGTAACTTTT  
 CAAAAGAATATCTTGTAAACATAGAATAACAGATCCTAGTGCATTACCACTCTTTGGGCT  
 TTATCGCTTTTCCACCATCATTATCTGCATCACTGCCTGCAGGTTTTCTACACGGCCAGG  
 GTTGGTCTCTGCCTGCTCAATAGTCAAGTCAAAGAGGCAGGAAATTAACACCCTCTGGA  
 GGCAGCCTTTGAGGAATGATCCATGGGAGGTGGAGTATAAATACCTCAGCTCTGTTTCCT  
 CTAGAGATATAACTAAGGAATGGGTTTTACATTGTTTCTCAGAGTTTCCTCAAGGTTTAA  
 AACTTCAATCACCCACAGGGGTAGTGGGCTTTATCATAGTATACATCCCTTTGTGGCTT  
 CCCTTCCTTCTGTCTCACTTCTCCATTCCAACTAGGATTTATTCTT

The following amino acid sequence <SEQ ID NO. 139> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 5:

NIITFFYEYSWSFQNKTSYWFNKLWYNQIMKLYAFVKVTFQKNILHRITDPSALPTLWALSFLHHYHLHC  
LQVFY TARVGLCLLNSQVKRGRKLTPSGGSLGMIHGRWSINTSALFPLEILRNGFYIVSQSFLKVLNFNHP  
QGVVGFIIIVYIPLWLPFLVSLHSLKLGFI S

The following DNA sequence Seq-2361 <SEQ ID NO. 6> was identified in *H. sapiens*:

AAGTATTCTTGTCACGGAAGAAGAAAAGGGTTGGGTAGTTACAGGGGGACAACAATGCC  
AGAACTGGGGAGTGTGGACTGGGATACAAGAGAATGAGGGAGCTCAGGATGAGCAGAAGG  
GCGGGGAAGCAATATTCATTAAGCACCTTCTATGTGCCAGTCAATAGGCCAGGCTTCAAA  
TTATTACCTTGCTGAAATCTTCACAGCAGCCCTCTAATAGGTATTTATCCCTGATTCCAT  
ATCCATGCTCTGCTTCCCCTCCTATTACAATGGCTGAAGAATCAAACCCCTTTCAAAGG  
CTAGCACTGTCATTTGCTCTAGATCCCATCCCCTCCATTTTCTTTTATTGAAACAT  
TCTCAATGGTATTCAACATACTCTGCTCTCTCTTCTATTAAATAGGCAATGCAACTCA  
TCAAGCTCTTTTCTCCCTTGGCTACTGCCCATTTCTCTACTTCCCTTTCATGGCAGAAC  
TTCTCGAAAGAGTTTTTCAACATCACTTCATTTCCACACCTCTAACTGACTTTTGAACAC  
AACTAGAGGAGGAGTAGGAGGGGACACTCATTCAAAGTGTCCAATTAAGCCCAATCCTT  
TAAAAGTATTATGTTGTGTCATGATGGCTGTTAAGAGCATGGTGAAAAGATATTAGAATAAG  
ATGTGGGGAATCATGACCGTGAGACAGA

The following amino acid sequence <SEQ ID NO. 140> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 6:

VFLSRKEEGWVVTGGQQCNWGVWTGIQENEGAQDEQKGGEAIFIKHLLCASQARLQIITLLKSSQQPSN  
RYLSLIPYPCASPPITMAEEFKPLSKASTVICPLDPIPSIFLFIETFSMVFKHTLLSLLLNROMQLIKLE  
FSLGYCPI SLLPFMAELLERVFNHFI STPLTDFLTQLEEEEGTLIPKCPKPNPLKVLCCHDGCEHGEKIL  
EDVGNHDRET

The following DNA sequence Seq-2362 <SEQ ID NO. 7> was identified in *H. sapiens*:

AAAGAAAAAGAAAGAGTAGTGTAACAATTCCACTTCTGGATTAAACATTGTAAGGAGACTG  
TGGACCTGTTACAGCAGAAAAACAGATATAATAGGCAAAAATTATTTTTTAAAAAATCTCC  
AGAAATCTGTTCTAAAAACATACAGCAGACTTTTAAAAAACTTGTCTGAGAAAATGTACTA  
AATCTCTGTAAGACAAACAAGAGTCTGTGGCAGCTGAGCAATGTTTGCCCTCACTCTAACC  
TCTCCCTCCAGGTCACTTTCATAAAAGTCAACTCTGGGAAGGTGTGCCCAAATTGAGA  
TTACCTGCCCATTAATTTCCAATCAAAGGATACAGTATATCACCAGGAAGGTAGCCACC  
AGCATTTCTCAGCCCCTCTTACTCCAAGTTGCAGAGGATAAATTCCTGGTGAGTATGGCC  
AGGAGGCCACGTGGCCACCTGGCCACCACTAATAGATCAGAGGATTAATCTCACACATGG  
AAGGATGAGCATACTGGGCCCTGATTGCCCTGACCCAGCTTACTTATAGGATGGAAGT  
TTCACATCAGGA

The following amino acid sequence <SEQ ID NO. 141> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 7:

SCETSILVSWGQGNQGPSMLILPCVRLILSISGGQVATWPPGHTHQEFILCNLEEGLRNAGGYLPGDILYP  
LIGNWGRSQFGHTFPELNFYEGDLGGRGSEANIAHVPTLVCLTEIYIFSDKFFKSLLYVVRTISGDFLKN  
NFCLLYLFSAVTGPQSPYNVNPEVELLHYSFFFF

The following DNA sequence Seq-2363 <SEQ ID NO. 8> was identified in *H. sapiens*:

AGTTAACAAAAAATACTACTTAACCTCTGCTAGAACATAATGTGATACATTTTGTACAC  
CTCTTAGCTTCTTTAGCTGAATTCAGAAATGCAACCATTAGTATTAAAGAGCAGGTACT  
AAGGATTTTCCAAATCATTTTGTATTCTTATCAATATTTCTAGTATTCTTTTAGATCCC

TTCACTCACTTTCTCTATTGCTTTCCATTTCCTGAAGTTTTAAATAAAAATTTCCCTTCTG  
 TTTGTCTTGTAGGAAAAATCATCATGCTTACCACATAGAATGTGAGTTGTAGGAGAGACA  
 CAATGGGAGACATCGGTTAAGGGACAAAAGACATTAAACATTTTAGGTGATTGTGAGTTCA  
 TAATTTTTCCAGAACACAAGCATTGCATGGCTACTCTAATATACTAGATTATTAATAATAG  
 ATATATCTTTGCCCTACCTGATAAACTATTTGTATAAGTGAATATATTTTAATATTA  
 ATCCAATATATTTATAAGAAATATTTGATTGCAAAGTAATCTGAGCATTACGATGATT  
 CCTATCTAAATACTGGCATGGTGAAAATGAGGACAAATCTACCCCTTTCTCTAATGTAGT  
 TACAGGCAAGCTATACTCATATAATAAACATAGAACGTACAATCAAACCAATGCATGAG  
 TGTAGGATGCAACTAAAGTCAAGA

The following amino acid sequence <SEQ ID NO. 142> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 8:

SQKNTTFLLEHNVIHFHLLASLAEFQKCNHYEAGTKDFPNHFVILINISSILLDPFTHFLYCFPPPEVLNK  
 ISLLFVLEKSSCLPHRMVVGETQWETSVKGQKTLTFVIVSSFFQNTSIAWLLYTRLLKIYLCPTTLFVVNI  
 FLIIQYISEIFDLQSNLSITMIPYLNTGMVKMRNLNPLCSYRQAILITNVQSKPMHECRMQLKSR

The following DNA sequence Seq-2364 <SEQ ID NO. 9> was identified in *H. sapiens*:

ATCATTTTGATAACCAGTCTGATCTGAGAAAATTAACCATGTCATTCAAAACATGTCCT  
 CCCCAAATTTAAGAAACATTAGGTCAATCTCCTGGTTAAATAATAGCTGTATGTTTTAGT  
 AGATTTTGAAATATTATGTAATCATTGAAATTATAAGCTTCGGCCCACTTGACTG  
 ACAATACCTGTTTCATTATTTTAACTAGCCTTTGTGGACTACATATCTCAAAGACA  
 AAAGAAAGATAAAAGTTGAAATAATCCAACAGTTATCCTACACAAAAGTATGACAAAATT  
 ACCGTTGCAGAAATTGAACTCATCAAGCCTGAACCTTTGACTTTGAACAATTACATGGAA  
 GAGTGCCACCATGGTGAACGTGTAGACCTGTACAGCATCACAGCCAACCTCTATACACAAA  
 CAAGGGTGGGCTGTATTCTGACCATTATTGGAATAAATTATCCTGATTACCTAATGTCTC  
 TTCACACCCCACTAAATTTATTATTATTATATTATTTTACACTGCCATCAAATTAAGTT  
 GCTAAACACAACTTTGCTCATGTTCAAAATTTCTATAGTGTGCCTCAACAATCACTAAC  
 TAATCCTCAGAATTAATTACCTACTAATTGTTTTTGACAT

The following amino acid sequence <SEQ ID NO. 143> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 9:

SFPVSEKIKPCHSKHVLPKFKKHVNLVLYVLVDFEILCNHLKLASGPQLDQIPVSLFTSLCWTTYLQR  
 QKKDKSNNPVILHKSMTKLPLQKLNSSSLNFLTITWKSATMVNCQTCTASQPTLYTNKGGLYSDHYWNKL  
 SLPNVSSHPLNYLLLLLYFYTAIKLKLKHNFAHVQNFYSVPQQSLTNPQNLPNLPNLT

The following DNA sequence Seq-2365 <SEQ ID NO. 10> was identified in *H. sapiens*:

TCTAGAATCTATACATACTATGTCCAATCCCTGTTCCACAAGTAGTTATTTATATGTGCG  
 AAGGTTCACTCCTGATTTTCCTTTTGCTCCAGGGCAAAGAAAAGATACTGAAATACAA  
 GGTGAGCTTATATCAGCCAGTAGTAAGCCAGTGAGGGCTACCACAGTTTGAAGAAGCA  
 GGGTGAACTTTTACATGAGATTGGGGGAAAAACCATACTGAATAATAAAGGGTTTAA  
 CTGAGATTGAAAGATAGTGCTTTGAGAAGCACACAAAAGATTCAAAATGGGCGTATAAAG  
 AATGACCTGTGCTGAAAAACACATTTTTCGCTACAAGGGACCCAATTGACTAGATGAGA  
 ATTTGTGTGAAAAAGGAGTTGATAAGGCAGGCTGGCACATTGCAGCCAATCTGTGAAAGG  
 CTTTTCATGTCCTGTGAACAGGAAATCACATATCACAAAGAGTGGTCTAGGAATCTGTGTC  
 TGCAACCCCTACAGTGGGGCAGACTGAAGAGGGAATAACG

The following amino acid sequence <SEQ ID NO. 144> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 10:

VIPSSVCPTVGLPDTSTTLVICDFLTGHEKPFDTWLQCASLPYQLLFHTNSHLVNWVPCSAKMCFSAQV  
 ILYTPILNLLCASQSTIFQSQLKPFIIQYGFSPQSHVKVSPCFQTVVALTGLLGYKLTLYFSIFSPLPWS  
 KKRIRSMNLRITYKLLVEQGLDIVCIDS

The following DNA sequence Seq-2366 <SEQ ID NO. 11> was identified in *H. sapiens*:

ATGGGCACCGCTCTCTTTAAAGTACACTTTCCTGACTCAGCTGTCTCTTTTCCTCCTCC  
 ATTCCCACCAATTCTGGGCTACAGGCTTTTCCTCTACTCTCCACAGCATCCTCCCTGAG  
 CCCTCAATCAAAGCACCTACAATACTGCCCTCATAGGGAGGTGCTATCTTTCTGTCTTC  
 CCTGAGCGCTGGGACCCATTGCATTTACCCATTATCCCCAAGGCCTAGCACATGTCTA  
 GCACAACACAGCAATTAAATAAACCCCTGTGGAATAAATAATTGTGGAATAGCCTGTTTT  
 CCATGGATGGTTATACAGGTTGTGCACTGCACAACCATGTGCAACATTCTGGAAAAAGA  
 CAGAAATTTATTGATTGGTTGGGGGTTTTGAATAGCCAAGGAAAACTATTGACCATTGC  
 ATGCCCTCTACCTGGGAAATCACATACCCTAACAACTTCTTAGGCCTTACTGCATGGTC  
 ACATGGGGTAACATTACATAGTTTCTCCAGCTCTCTAGTCTGCCCACAAAGGTGATATT  
 GTTCAAAGGGGCAATCTTTCCTTGCCCTCCACCAGTCTATTCTTAACCTGACCCAAAGTAA  
 TCTCTTTCACTGCTTACCAAAGATATTTAGCTTCAGCTATCCTGTTTGCAGAATGGTGA  
 CGTATTCC

The following amino acid sequence <SEQ ID NO. 145> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 11:

MGTALFKVHFDPDSAVLFSSSIPTNSGLQAFLLSHSILPEPSIKAPTILPSGGAIFLSFPERWDPLHFTHL  
 SPRPSTCLAQHSNINPVEINCGIAWFPMVIQVVECTTMCNIPGKRQKFDWLGLVNSQGLFDHCPSTW  
 ENHIPQLLRPYCMVTWGNHITVSPALSAHKGDIVQRNLSLPSTSLFLTPKSLSLTKDISASAILFAEWR  
 I

The following DNA sequence Seq-2367 SEQ ID NO. 12> was identified in *H. sapiens*:

TCAATAGCAATAAAGCACACCAAGCACACAGATCTCGACTTTTGAATGCCACTTCTCCAT  
 CTTAAAAGACAAAAACAGGACATCTTAGACAAATGGCCAACTCCAGGGTGGTTGGGGCAA  
 GGAAAGAAGACGTGCTTGTGCACATCTTGTACATCAGGTTTAGGAAGCTGTCACTGGTC  
 AAATCTGGGACAACTTGAACATCAAAATAAATAATCATTGTAATGGATTATAACTCATCG  
 ATGTAAGTCTCTAAGTACACACTTATATCAATACATATGTACATATACACATACATACAT  
 CTTTACATACTACTGAATGGCACTAATAATGGCATTGGCAAAGTGTATGCTAACAAT  
 TAACTCAGGCAAGAAACATCAATGGAGGCTAAAGTGGTAGATAAAATTGGGATGAGTAG  
 ATTTTACACAGTCTCCAAGTGACTTCCACAAAATACCCATTATTACAAAGGAAAAGATA  
 GATAGGTTTGCAGCAGAAAAAAAATGTGACACATCATCTTAACTAGGGGATCAGTGTTA  
 ACTTCTCCAGCATGAGACAAGTAGACAAACAAGTCCATCAGAGAGGATGAAGTAAGACA  
 CAGCATCACTTCTGTGAAATTCTGG

The following amino acid sequence <SEQ ID NO. 146> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 12:

RISQKCCVLLHPLWQLFVYLSHAGEVNTDPLVKMMSDIFSAANLSIFSVMGILWKVTRWLCKIYSSQF  
 YLPVLASIDVSCLSLLAQFAKCHYLPFSSMRMYVYMYICIDISVYLETYIDELSIYFYFDVQVVPDLT  
 SDSFLNLMYQDVHKHVFPPCPNHPGVGHLSKMSCFLLRWRSGIQKRSRVCLVCFIAI

The following DNA sequence Seq-2368 <SEQ ID NO. 13> was identified in *H. sapiens*:

TCCGATGATGTTAACACCATATTATTTTAAAGAACATGAAGATTACATAAGAGTAGGCA  
 TTTGCCATTTGTATTTTAAAGAGTCTGCTCAGCTCTTAACAAGGAAGGGCCTATGCAAA  
 ATGAGAAATAAAGTGAAGAACGATTGCTTGTGAGTCTGAAATAACTTAGGTGTCAAAA

CAAGTAACTTTCACCCTCCTTCAACCTGTCTCTTGCCATTTAGCAATCTAAAATAATTA  
 TCCAATGTATGGTTGCACTCCAAAAATCATGTTAACTTGAGATATTCTGAATTTTGTGT  
 ACAATTTTGGTAGAGGGTAAGAGATAGAGAAAAATCTTACATTGTGTTCAAGTGAATTCC  
 CAGACCTCGGGGGTAAAATAAGTGCAGGAAGAATCTCATCAGGATATCTCGGGCAATTTT  
 TCATTAGTACGCATGACAAGCTGTTTACCACAGGCTATTGTTTTATGGAAAGTTCAAA  
 TATAGCAGGATGGGATGTATGGTGTGATATTAACACATATGAACACAATTATTACCTATT  
 TTAGGTATATACGACCTTGTACCTAGAACATTGATACTTTCATTATGATGTACTTT  
 TATAGAATAAGATAAA

The following amino acid sequence <SEQ ID NO. 147> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 13:

YLILKYIIMKSINVSQRQSYIPKIGNNCVHMCYHTIHPILLYLNFQPKQPVVKQLVMRTNEKLPEISDSSCT  
 YTFPEVWEFTEHNVRFSSISYPLPKIVHKIQNISSLTFLECNHTLDNYFRLLNGKRTGRRVKVTCFHLSTYF  
 RLTSKSFFTLELILHRPFLVKSADSKYKANAYSIVIFMFFKNMVLTS

The following DNA sequence Seq-2369 <SEQ ID NO. 14> was identified in *H. sapiens*:

GGCCTCTCTGAAGGGGAAGCAAGCTTGCATCTAGACTTCTTTCTAAAGATAACCTAGACA  
 ATAATGAATACAGCTGCCACCAGCCTCCTATGCACTAGAGGCATTATTCTAGGAGTTTCC  
 GTGTATTAAGCTTATCCTGAAATTAGTTCTTCTTATGACTGAGAGGAGAAGTATTACAT  
 ATTGATTTTCAATTGTTAGAAATGGGAAAATTTTAAACAAGTGTATTAGAGGGCAACCACA  
 TTTTCTGCTCTGCAACCTGCTTCTCCCCCTTACGTCAGGACATCTAGATGAACCCATC  
 TTCGGAAAGGCTGCAGAGAAACATGTCCTACAGACCTACTATCATCTGGTTAAACAATCC  
 CAGTGGACGGACCAAATTCAGACGCTTCCCACTTCTCTCCACTGCACGGATGCTGCC  
 ACACATGCTCATATACCTCTGAACCTTCCAGTGACTACGGCACAGCGACAGCTGAGTTCC  
 TGGGCGCAGAACCACTGGGGCACGTTTTGGCAGCTATGAGCAAATCACTGTGCACAAAGG  
 CAATCCCAGTTTACACTTCCACAGAGAGGAAGTGAATACACTGCCACCCTCACCTGAC

The following amino acid sequence <SEQ ID NO. 148> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 14:

GLSEGEASLHLDFFLKITTIMNTAATSLLCRGIILGVSVYAYPEISSFLLRGEVLHIDFIVRNGKIFNKC  
 IRATTFSAIQPASPPSRQDIMNPLFGKAAEKHVLQTYHVLNNSQWTDQNSRRFPLSLHCTDAATHAHIPL  
 NLPVTTAQRQLSSWAQNHGTFWQLANHCARQSQFTLPQRGTEYTAHPL

The following DNA sequence Seq-2370 <SEQ ID NO. 15> was identified in *H. sapiens*:

AATACTAGATTCTTTCAGGGACTTTTGTAGAACAGGACAAGAATAATCCTTCTCGACAA  
 AGTAAGGAGTGATCTATCTCAAGGCAGAGCATATTCTCTACACCAGGAGGAATTTCCA  
 TTAACATAAGCAATGCCCAAGGATGCTTGTATCAATTTTATTCAATGTTGTTTTCTGT  
 GTTCTGGCCAATATTATGACTTGAGATACAAGTGAGAAAGATGACTAAAGGAATTCATGA  
 GACAAGATGATCACTATTCAAGGATGGTATGATTGTCTATCTAGAAAAAAGAAATGAC  
 TCCATCGTAATTCTGGGAATTTACTAACAGTGGCTGGGTCTGGACAAACATTTAAAAAA  
 TCAATCGTTTCTTGTGTGGCAGCAATAACCATTTAGAAAATGGAGTAAATGCGGAGTTA  
 AGGGGCTGTGAATATATAACAGCAAGAACTCTGATCTGCCGTCCCACAAAGTCGCCTCC  
 GGAGTGGACACCGCCAGGGAAGGCAGGTCTCTGGAGGGAAGGTAGAGAGAAGATACGGA  
 GGATCTGCCCCCTCCCCAGGAAGCTCCCCGAGAAAGGGCCACAACCTGTTTACTCCAGCAG  
 GCTCTGGGGGATTACG

The following amino acid sequence <SEQ ID NO. 149> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 15:

ILDSFRDFLEQQQESFLDKVRS DLSQGRSIFSYTRRN FHHKQCPKDACYHFYSMLFSVFWPILLEIQVRKM  
TKGIHETRSLFRRWYDCLSRKKEMTPSFWEFTNSGWLDKHLKNQSFPCVAAITIKMEMRSGAVNIQQELL  
ICRPDKSPPEWTPAREGRSLEGRREDTEDLPLPQEA PRERATTVYSSRLWGDS

The following DNA sequence Seq-2371 <SEQ ID NO. 16> was identified in *H. sapiens*:

GAAACCTTTGACTACTTCTCTGTCTCACGGTCATGATTCCCCACATCTTATTCTAATA  
TCTTTTCACCATGCTCTTAACAGCCATCATGACAACATAATACTTTAAAGGATTGGGCT  
TAATTGGACACTTTGGAATGAGTGTCCCTCCTACTCCTCCTCTAGTTGTGTTCAAAAGT  
CAGTTAGAGGTGTGGAAATGAAGTATTGTGAAAACTCTTTGAGAAGTTCTGCCATGA  
AAGGAAGTAGAGAAATGGGGCAGTAGCCAAGGGAGAAAAAGAGCTTGATGAGTTGCATTT  
GCCTATTTAATAGAAGAGAGAGCAGAGTATGTTTGAATACCATTGAGAATGTTTCAATAA  
AAAGAAAAATGGAGGGGATGGGATCTAGAGGACAAATGACAGTCTAGCCTTTGAAAGGG  
GTTTGAATTTCTCAGCCATTGTAATAGGAGGGGAAGCAGAGCATGGATATGGAATCAGGG  
ATAAATACCTATTAGAGGGCTGCTGTGAAGATTTCAGC

The following amino acid sequence <SEQ ID NO. 150> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 16:

LKSSQQPSNRYLSLIPYPCASPPITMAEEFKPLSKASTVICPLDPIPSIFLFIETFSMVFKHTLLSLLL  
RQMQLIKLFFSLGYCPISLLPFMAELLERVFHNHFISTPLTDFTLQEEEGTLIPKCPKPNPLKVLCCHD  
GCEHGEKILEDVGNHDTRETEKVVKGF

The following DNA sequence Seq-2372 <SEQ ID NO. 17> was identified in *H. sapiens*:

ACAGGGCATCCTCGCCTTCCACCCACTTTAAACAGCCTGCAAGGCAGTGTGTGACCTAT  
GGCTTTAACTCTGATGAGGAGGATTCTCATGGCATGGGTGCTGAGAACCTGAATCAC  
AAGGTATAAAGCAGGGACCGAAGGACTGTGCCACTGCAGCAACCCCGCTGGGTTTAA  
TGCTCTCCTGTTGCCACCCTGAAATTTTAAAGACTTTTACGGGGTCTTGCTCTGTCTAT  
CTAGGCTGGAGTGCACTGACATGATGCTTATACCTCATCTGGCTGAGACTCACTAGAGA  
AGGTCACTAGTTAGAACTAGAGAGGGGGCTGGGCACAGTGGCTCATGCCAGCACTTTGGG  
AGGCTGAGGCAGGAG

The following amino acid sequence <SEQ ID NO. 151> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 17:

TGHPRLPPTLKQPARQCVTYGFNSDEEDSSWHGLRLTLNHKVSRRRTVPTAATPRWVCSPVATLKFLKTF  
YGVLLCHLGWSAVTCLIPHLAETHRRSLVRTREGAGHSQCHFGRLRQE

The following DNA sequence Seq-2373 <SEQ ID NO. 18> was identified in *H. sapiens*:

CTTGATAGCAATATAAAGCCTTAAATTTTTTTCTGTAGGAAATATCACACAGATGGCTA  
ATTATATGCCATATAAAGCCATTAAGGAAGAAAGGATGGCAAATGCTCCTTTTAGTGAGA  
CTTCTTTGTTATGAGATCTGGGTATAAAAATGTGCAGGTGTGTAAACAGAGGAAGGAGAA  
TTCTGATTAAGTCCCTCAAGAATTGAAGAAAATGGGGTGAGAGACAGAGAACAACTGTGA  
GCTAGGAAAGCTCAAGGAGTAAACCTAACAAGAAAGTTTAAAGCAATGGCTACTTTTATAC  
AGTTTATTTTAGTAAGTGCAAACTCTTAAATGAAGTTATTTATAAAGTTTATTTGAGTT  
GTTTCTGATAATTAAATAGCATGAGAAATGGGAGGAATTTGAGATATTGCAGTTAGAAA  
GGGAGCAGTGCACCAAACTTATTCTTAACTTAAAGTTTCACTCTTACCTAAGGTAAGT  
CCTAATGTGACACCAACTTAAAGCTGAATTAGACAGGAATATTGCAATGAATAAGCAATG  
ACTATTACAATCTACTCAGCATAAAAAGGTTTATTAGAAAGGTTCTGCAATAACACT  
CTATGTAAGAGTTTATGGAACAATTAATAGAATAAAATGATGTACATTTTATGTACTAC  
TGCATTTTACATATTCTAAGGCACGAG

The following amino acid sequence <SEQ ID NO. 152> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 18:

LV AISLKFFCRKISHRWLIICHIKPLRKKGWQMLLLVRLLCYEI WVKAGVTEEGEFLSPSRIEENGVRDR  
EQLARKAQGVNLTRKFKQWLLLYSLFVQILKMKLFIKFIVVFLNSMRNGRNLRYCSKGSSAPNLFITKFI L  
LPKVSPNVTPTSIRQEYCNEAMTIHNLISIKQVHERFCNNTLCKSLWNNNKIDVHFMYYCILHILRHE

The following DNA sequence Seq-2374 <SEQ ID NO. 19> was identified in *H. sapiens*:

CCCTTGAGCACACACAGGGCGATACTTGCCACAGGTGGGACTGAAGGCTTCTTTCTTGG  
CTTATAGTTTGAAGCAATGGGAGTTGGGAGCTCCAAATCATTCATGGGACAAATATCCT  
GTCTTATATTGCTTAAAAAAATCCTATCTAATTTTAAAGACAGGGTGTTTGTCTCTTA  
AAGCACTTTGCATTAAATGTGTAAATTACAGAAATTTCAATGCTCTCTGAAGAGGTAA  
TTGATATTAAACCATGGTAATTCTAATAGCTAACACATATTGGGCATACGGTTTTTTCACAT  
GTCTAAACAGTCCATGTTTCTTAAAAATGCAGATTGCAGGGCCCCACACTGGCTGGGGA  
ATTGCACTTCCAGTAAACACTTCAGATGATTTTCATGATCTTTCAAGTTCGGGGAAAAT  
GGAGCTCGTTTTCCACTAGATTAAAGCAGTATTCCTGATGCGTTCTCAGGCCCTAAA  
AGAATCAACACTCCTCAATAAGTAAACATTCACCTAAACATATCCAGGTGGATCCAATGA  
TCTACC

The following amino acid sequence <SEQ ID NO. 153> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 19:

VDHWIHLDMFKMFTYGVLIILLGPENAYSGILLSSGKRAPFSPNLKDHENHLKCLLEVRI PQPVWGPAICIF  
KETWTVTCEKPYAQYVLAIRITMVNINYLFRHKKLLTQLNAKCFKSKTPCLKNIGFFFKQYKTGYLSHEF  
GAPNSHCFQTISQERSLQSPVASIALCVLK

The following DNA sequence Seq-2375 <SEQ ID NO. 20> was identified in *H. sapiens*:

CTGCTCTATATAAAGATATAGTCCATGTATATGGCTGAGTCTTTTATAGTCCAAAATGTA  
TTTTTCTGTGTACTATGGTTTATTAACCTGACTTATTTTTCTTCTTTCAGATTTAAAAAA  
TGTTAACTAATTAAGTAACTTCCCAAGTACCTACCAATGACATTAATCTTCCTCTTTT  
GTCGTTTGTCTTTTTACCCCCAAATCCTATTAATACAGCAACTTTTTAATATGATTGTC  
TACTTTTCAGAGTACTTCTTAACAACATAGCAAATGCCAAAATGTTAATGGAAGTATTA  
TGAAAACATGCAAAAAATATTTCTTTATGATTCTGATAATTATGAAATGCCTTAGATT  
AAACATGAATAAAATTTAATTATTATATATGTATTCAAATAGTTGGATATATAGTCCTGAG  
AAAGAATCCTTCACTACATATGTTATAAAAAATGGGAATGAACACATTACCTAAGAAGTCT  
GCACTAGAAATAATAAGATACCTTTTCATTCTTGACATCTTCTTCTTTTGAACCAAGT  
ATCTGTA

The following amino acid sequence <SEQ ID NO. 154> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 20:

QILGSKRRKMSRMKRYLISSADFLGNVFIPIFITYVVKDSFSGLYIQLFEYIYNNIYSLIGNFNNYQNH  
KEIFFACFHYFHHFGICYVVKYSEKTIILKSCCINRIWGKEQTTKRGRLMSLVGTWEVTLISHFLNLKEE  
KVKLINHSTQKNTFWTIKDSAIYMDYIFIS

The following DNA sequence Seq-2376 <SEQ ID NO. 21> was identified in *H. sapiens*:

TATCATGCTGCCGCTTCCAATGGGCATCTGTTCCACCATGTTGTGGGCATTCAATGGAGC  
TTTGTGTGCTTCTCACTGTAAACCCTCCTGTATAATTCTGGGGTCCAGCAGAAAACAGT  
ATGTTACCCTAAAATAGGGCAATTGAAGGATCTTTCAAGAAGGGACAAGTTGTAAAGGTG

GGCAGCACAAAGGGAAACCAACAAAAATGAAGACCTGGTGGGACAGGGACAGAGTGA  
 GGATGCTGGAGAGACCCAAAGCTGCAAAGGAAAGGAGCAAGGGGAACAATACCCACCCCT  
 CTCCCCTCCCACCTCCCACCTCCCACCTCCATTCTTCTCCAGTGGTGCCGCCCATTTGGGC  
 AAACCCAGCCAGAAGCCAGGAAGCATGAGAGTTCAGCTGATGCAGCCCATACAGATCAGA  
 CTCCTGGACTTCAGAGTGGGGAGGGTGAGAGGGATGAAGTCTGGAGGCACCAATTGGGA  
 AGGCCATCCAGAATGCTCCTATTCTGTTTGGGAGCTGGGGATGGGAATGTCCCTTCCTGA  
 GGGTATTTATGAATAAATCAAATCAAATCACAGAAATCAAATCACAGAAATCAAAGCT  
 GGAGATTCTCTCCCTCTACTTGCTGGCAGCCAGGATGTGGGCTCATGACCTAAACTCA  
 GTCATTAGAAATTCCTCCGGGAATGCAGTCTTACAGGAGTAGCTCAAGGCCAGGCAGT  
 GGCTCACACCT

The following amino acid sequence <SEQ ID NO. 155> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 21:

RCEPLPGLELLDCIPRGNFMTEFRSAHILAASKRERESPALISVIFLFDLIYSINTPQEGTFPSPAPKQ  
 RSILDGLPNWCLQTSSLSPTLKSRSLICMGCISTLMPLPGFWLGLPNGRHHWRMEVGGGRWEGRGWIV  
 PLAPFLCSFGLQHPVTL~~SLSHQVFIFCWFFVLPFTT~~CPFLKDP~~SIALFGN~~ILFSAGTPELYRRVQEAT  
 KLQMP~~TTWNR~~CPL~~EAAA~~

The following DNA sequence Seq-2377 SEQ ID NO. 22> was identified in *H. sapiens*:

CCCATCTGTCTGAATGCCTCCTGTAGTGGGGGACTCACTCCCTAATGAATCAATCCCTCT  
 TGTCTTTGGAAAGGTCTTCCAAGTGAAGTCAACATCCAGTGAAGCTCCTCCACT  
 CATCCTTTT~~AGCTGGACCTCTGGGGACCAAGACAGCAGACAGCTGCCTCTTCTACAGG~~  
 GCAGCCCTCCAAATGGCTGGGGCCACTGTCTTCTCTGCACTAGAAGACCTTTCTATGGTA  
 GTATCCTTCCACATAAGCTATGACTTCTATCCCAGGAAAGCCTGATTTGTCTCCTCTAA  
 ATGCACTTCCACTTATCTGTGACCTCTTACAATGAAATCAGAGAGAGATAACCTGATC  
 TTCTAACTCAGAGCAAGCAAGCTCCAGGTCTT~~CAGAGGCCCTGCAGGGCACACAGATGA~~  
 CAGCGGATGACCAGAGGGCACATGCCTTGTCTAAAGGGGATG

The following amino acid sequence <SEQ ID NO. 156> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 22:

PICLNASCSSGLTPINPSCLWKGLPTELD~~SNIQSSSTH~~PF~~SWTLWGPRQOTSCLFYRAALQ~~MAGATVFSAL  
 EDLSMVVSF~~HISYDFYSQESLICLLMHF~~LSV~~TLLQ~~NQREITLIFLRASKLPGLQRP~~CAHRQ~~RMTRGHMP  
 CMHF~~LSV~~TLLQANLKGM

The following DNA sequence Seq-2378 <SEQ ID NO. 23> was identified in *H. sapiens*:

TTTCAACACTATGTAGCCTAAAGTTATTCCGTCATCCATGACTATCCTGTCTAAAGAGTC  
 TGAAGATCTTTATTTGGTAGCTATGGCTTCAGCTAGTTCAATTTGCTAAGTTACCTAGAGT  
 GGTGACAGATTTCTAATTATACGTTATGAGAGGTTACTCCCACTATTGCAAGAGACT  
 TCTGCCAAACATAGGCCAAAATTCATCTCCTTGGTTTGCAGGTACAGTTTGTCTAATCCT  
 GGAAAATAATTTCAATGAACTACTTCAGCGTTCAGAAACATTGGAGTTATAAATAGAAG  
 AGGAAGAGCCACATAACCTAATAGACAATTACCTCTCATATGCCAGTGGTCAACACATTC  
 ATAAGCCCATGTGTGCTTGATCCAGGGACCACACAGGTCCTGATGGATTCTGAAATTT  
 AAGGCTTTGGATTACTGGTAACAGAGACATGTTAAAGTACATGTCTTCAGTCTTGAGTAG  
 AGTGAATCAGTCTGATTCTTTTTTTTTTAAATGAGACAAACATCAGGTAAAGACCTTG  
 ACAAGAAGGAAGAGAAATCCCGAGATTCTATAATCATAAATCGAATTGTAATTGCTAG  
 TTTAAGTAGTCCTTCAAAAATACATCTCATTCTGACAGGATAAAACAAGTTTATAAAA  
 TATATTATATTCTGGGTTCACTAGGGGAACAC

The following amino acid sequence <SEQ ID NO. 157> is the predicted amino acid sequence derived from



the DNA sequence of SEQ ID NO. 23:

VPLVNPEYNIFYKTCFILSGMRCIFEGLLKLAITIRLLNLGSLPSCQGLYLMFVSLKKRNQTDYTLK  
TEDMYFNMSLLPVIQSLKFQNPSTLCGPWIKHTWAYECVDHWHMRGNCLLYVALPLSIYNSNVSERSSS  
LKLFSRIRQTVPANQGDEFWPMFGRSLLQWGVTSHERIIRNLSTTLGNLANELAEAIATKRSSDSLDRIVM  
DDGITLGYIVVK

The following DNA sequence Seq-2379 <SEQ ID NO. 24> was identified in *H. sapiens*:

CCTTCTCATCTTTGCTGCTCTCTGCTGACAAATTTAAAAACCCGACATGTGTTAACTCTC  
TCCTTGTCTTCCAACCCACCCACTTATCACCTCAGTGCCATGCTCCAGGTGGCAAGCAG  
AGAGGACTGTGGTTTGATGAGTTTCATTGCGGTGGCTTTAATTACTGATAAGAGCTTG  
ATTATACACATTCTCAAAGGCATTGAAAGTTAAAGAAAGTCCTTTTAGGTAGCAGTCC  
ATGACAAATGCAGTTCATGAAATCTGTGTCCTTTTCATTCCCTTCTGAGTAATTCCTCTC  
TGTCTCTATCAAAGCCTTGGATACTCCATGGTTTACTAGGCAGAAACTTATCCATCCAAC  
ACAGCCACATGGATACAGCTTTGTGCTTTTAGACAATAACCACTTGAGAAAACCTGACCT  
TTTCCCCCACTCTTCATTGAGCTTCTGTCTGCTGCTGAAAACAAGAGGACATCCTGCCACAT  
TGTCTCTGCTCTGCTTACTCTTGAGAAGTCTAGTTGGGAAAACAGGCCCTATAAAGAG  
AGACACTGCAATGCCATGGGGTGGAGACAATAAAGTGATGGCAGCAGAGCACTGGAGAG  
CAGAGGTGGGGTCACCAACTGCCCAATGGCACTGTCCCTCAGAACTCTTGCAATTTGCT  
TTTAACGCA

The following amino acid sequence <SEQ ID NO. 158> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 24:

LPHLCCSLLTIKPDMLSPCLPHTPLITSVPCSQVASREDCGLMSSFMPWLLLIRALYTFSKALESKKVLL  
GSSPQMFMKSVSFSFPSEFLSVSIKALDTPWFTTRQKLIHPTQPHGYSFVLLDNNHLRKPDLFPHSSFSFC  
PAENKRTSCHIVICSALLRLSLVGKTPGIKRDAMPWGEDNKS DGSRALESRGVNTNCPNGTVPSSELLHLL  
LT

The following DNA sequence Seq-2380 <SEQ ID NO. 25> was identified in *H. sapiens*:

AATTATGACATTATGACAGTTTGTCAATTAAGATAACATTCCAAAGAGAAATGGGCATG  
GGCATATATTTACCACTCCCAAGGAAATAGCTAATAAAGTAATAGAGTACAGATTAAAT  
AATAAAATCCAAATTTAATCCATCACATTGACAATGATTAATTAATTTAAAGCAGTG  
TTGGGAAGAATACAGTGAGCTGGTGTCCATACACACTGTGATGAGAGTGTAGAAATCTTA  
CAGTCTTACCAGAAAGCAAATGTATCAAACACTTTCAAATGTTTACACTTCTTAACCTA  
GAAATTCACCTTTTAAGAATTTCTCCTAAGAATATATCTTTGTTTAAAAATATTTACATA  
CAAAGATGTTGATTTTAGTATTATTTGAAAGCAAATAACCCACAGAATCTCAAGTATA  
TGATCCAAACAATGGAATATCTTATAGCCATTAATTTTAGAGATGAATATTTAATAATTT  
AGGAAAATACCTATGATACTTTAAATTTTAAAGGTTACATAGCAGAAAGAGCCATATTT  
CAATTTTGCCTTGGAAAATATGGTATCACTACAGAAATGTTGTAGTGTATCGCTGAC  
AACACTAGTTATCTAGGATAAAGGATATTCTCATTTTCATTTACCTTTAGTA

The following amino acid sequence <SEQ ID NO. 159> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 25:

LKVKEYPFILDNCCQRHYNISVVIPIYFSKAKIEIWPLLNCNFKFKVSVFSI IKYSSSLKLMAIRYSIVWI  
IYLRFCGLFCFQNNTKINIFVCKYFTKIYSEKFLKVEFLGEVTFKCLIHLLSGKTVRFLSHHSVYGHQLT  
VFFPTLLIFSLSMWIKFGFYFNLYSITLLAISLGVNIPCPEFLFGMLSLMTNCHNVIN

The following DNA sequence Seq-2381 <SEQ ID NO. 26> was identified in *H. sapiens*:

CCAATATTGATCTTTCTATCTTTAAAAATGGCAGTTTCATGTGTCTTGATCTAAAATC  
 TTAAAATCAATCTTTCAATTGGATAAGAGGCAGGGAAATTAGCTTGGAAAGGTAAATCTAT  
 TATCCAGAGGCCAAAATTTATGGGCTTTGATAAAGGTGGATATTTTCGATAAGGAGGAA  
 AGAGTAAATTTTACTAACATACTTTGGCTTTTGTTCAGTTTCTTAACCTCTATTTTCGC  
 TTTATTATTTATTTTGTGTTTACTCTTGGGAAAGCAAATTATTTGTTTTCTCACATCT  
 TTTGGGGTCCAATTTTGATGATTCTGATCTTTTTTAGTTGCTTGACCTGTAGACCCCTCTA  
 CAGAACATTGCAGGGCCTCTTCTCAGAGGAGCAGCGGTGATGAGCTTAGTTTCCTAGGCT  
 GGGACTGTTGCGCTGGACTTGACAGGTGAACTGAAAATTGCAGGGATAAGTACACCTATT  
 GAGAACAACATCCCATCTCTTTATCAAAGCTCTTCATTGGCTTTGGAAAAGTGTGTAG  
 GCCTAAGGAACTAACTTTCTAGGGATATTCTAGGTTTTAAACATATGAGAAAGAGAAA  
 GACGTCGGTTCTTATTTAAGAGAGTTTATGAGACCTTATCCTTGAAATAGTCAAATTTAT  
 AATGACATAAGGCTGTATGTGTAGTT

The following amino acid sequence <SEQ ID NO. 160> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 26:

NISFLSLKMAVSCVLINLKNLSIGEAGKLAWKVNLLSRGKISWALIKVDIFRGGKSKFYHTLAEVQFSPL  
FSLYYLFECFTLGKANYLFSHIFWGPILMLIFFSCLTCRPSTEHCRASSQRSSGDELSFLGWDCAGLDR  
TENCRDKYTYEQTSHLEFIKALHWLWKTAVGLRKLNPLGIFVLNIERERRRFLFKRVYETLSLKSNI  
MTGCMCS

The following DNA sequence Seq-2382 <SEQ ID NO. 27> was identified in *H. sapiens*:

ATAAAATACAGATCTGATTGTGTCACTCTCCTGCTTAATATTGTAGTTGACCCTCCAC  
 TGCTCTCATGAAAGTTCATAATCCTTACTGTGGTTGTAAATGCCCTTTTATGATCTGTCC  
 CTGCCCCATTGTGTACACTCATCTGTGCTACTCTCTTCTTCATCAATATGCTCCACC  
 ATACTGTCACTTTCTGCTTATTTTTTTAAAAAAGTATGGAACATCTTTCCCTTAT  
 GTGTCTTATGCAACCTGTCAGACAAAACCATGTTATATTTCTCAACACACAATTTTA  
 TTTCAGGTCTCTGTGCCCTTTACAAATCTACTAATCTTTCTGTCTGGAGTGTTCTTTCTT  
 CTCCTGGCCAAATCTAATCATTTGTCAAGAGTGCAACAGCATCATTTCTTCTGTGACTC  
 AATTCTCCAAGCATCGTATCCTCTGTGTTCTATAGCACTACATTGGATCGGTCCATAAC  
 AATTCTGTGTCAGTGTATTATAAGAACTTATTACAGGTTTTGTCTCTTCTACTATGGCGTG  
 AGCCTTTTAGTCATATGAATTGTGATTTTGTATATTTAGCGCCTACCATGGTGCTTAATT  
 CGTGGTAGGTGCTCGGTAAATG

The following amino acid sequence <SEQ ID NO. 161> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 27:

KIQILCHSPAYLLTLPLLSKFIIILTVVNNLLSVPCPFVYTHLVLLSFFINMLHHTVIFLLIFFKKVWNIS  
FPLCVLNLSDKTTTCYIFSTHNFISGLCALYKSTNLSVWSVLSSPGQILICQECNSIISSVTQFSKHRIL  
CVPIALHWIGPQFCQCIIRTYLQVLSLLWREPFSHMNCDFVYLAPTMVLNSWVLGK

The following DNA sequence Seq-2383 <SEQ ID NO. 28> was identified in *H. sapiens*:

CTATTGGTTTAATAAATTATGGTATAATCAAATAATGAACTCTATGCATTGTAAAGT  
 AACTTTTCAAAGAATATCTTGTAACATAGAATAACAGATCCTAGTGCAATACCCACTCT  
 TTGGGCTTTATCGCTTTTCCACCATCATTATCTGCATCACTGCCTGCAGGTTTTCTACAC  
 GGCCAGGGTTGGTCTCTGCCTGCTCAATAGTCAAGTCAAAGAGGCAGGAAATTAACACC  
 CTCTGGAGGCAGCCTTTGAGGAATGATCCATGGGAGGTGGAGTATAAATACCTCAGCTCT  
 GTTTCCTCTAGAGATATACTAAGGAATGGGTTTTACATTGTTTCTCAGAGTTTCCTCAA  
 GGTTTTAACTTCAATCAACCCACAGGGGTAGTGGGCTTATCATAGTATACATCCTTTGT  
 GCCTTCCCTTCCCTTCTGTCTCACTTCTCCATTCCAACTAGGATTTATTTCTTTTCCCT  
 AAAACAAAACAAAATGTTTAACTGAAACCTTACAAAACACGTAAATTTATATTTAAA

AAATCTAAATATTTGAGGAGAGAACGAAACCTAAGTATATGCCAGGTATAACACGATTG  
GTGGAGATAGCTTTAAAAAAGTTCTTGAAAAATTTAGTTTTTTAAAAGGGTACCCTAGTAG  
AAGGTGACTTAACTGCCTAATTC

The following amino acid sequence <SEQ ID NO. 162> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 28:

YWFNKLWYNQIMKLYAFVKVTFQKNILHRITDPSALPTLWALSFLFHHHYLHHCLQVFFYTARVGLCLLNSQV  
KRCGRKLTPSGGSLGMIHGRWSINTSALFPLEILRNGFYIVSQSFLKVLNFNHPQGWALSYSFVASLPSCL  
TSPFQTRIYFFSLKQNKMFNLKPLQNTNLYLKNLNIGENETVYAQVHDWRLKSSKIFLKGYPSSRLNCLI

The following DNA sequence Seq-2384 <SEQ ID NO. 29> was identified in *H. sapiens*:

CTGGCTTCTGAGAGCCTCCTGGTTAGGAAGGAAGTTGTTCTCTTTCCACTGCAAGCTTAG  
AAAGCCTTCCAAGTTCTCTCTCTGTCAGCATAAAGAGACAATAACTCAGAGGAAGGTAT  
CCCCAGGAGTTTCCAGACAGCTGCACAGATTTAAGTGCAGAAATCTGAGCAGAGGTATAG  
TCTTGGCATTACATGAACACCTTTCAGTAGCAGGAAGAATAAATGGAAAGAGAGCTACA  
GAAATACCAGGGGCGAAGTCTTCATCTGAAAGTCCAATCTTTGATCAAGAGCTGGTAGGA  
AGTCTGAGAATTTGTATCAGCAGTGATTCTAGGCTGTCTGGTCTGAGTAATTGGGATCAG  
AGCAACAGCTGATATCATGCTTACCTTGTGCCAGGCTCCCTTCTAAGGGCTTCCTGGACA  
CCTGCTCGTGTGAGTCTCACAGCAATCACATGAGGTATGTTCTGTTGTTGTCTCCTTGT  
GCGGATGAAGACACTAGGCACAGAGAAAAGTGGCCACAGGTGTACAGCTGGGGAGGCCAG  
AGCCAGAATTCAGACCTGGGGTGTCTTGGCTGATGTGAGCTAGTGTGGGCCAGCATGGGA  
CACAGAGGGAGGATTAGCTGGAGAAGCAGGACAGAGGGCAAGAGAGACGAGATCTCCGAC  
AGTGCTGGGTGAGAGACACTTTCCTGAGCCATGATTAAACCTGATTATGGGACATGTTTT  
AGCCTGTGACA

The following amino acid sequence <SEQ ID NO. 163> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 29:

LASELLVRKEVVLFPQLQAKAFQVLSFCSIQRQLRGYPQEFPSCTDLSAEIAEVSWHLHEHLSVAGRIN  
GKRATEIPGAKSSSESPIFDQELVGLSLRICISSDRLSGLSNWDQSNZYHAYLVPGLLRASWTPARVSPH  
SNHMYRVLLLSPCAEDTRHRENWPQVYSWGGSQNSDLGCLGCELVWASMGHRGRISWRSRTEGKRDEIS  
DSAGSETLSAMIKPDYGTCTFSL

The following DNA sequence Seq-2385 <SEQ ID NO. 30> was identified in *H. sapiens*:

ACAGTGAGCAGAGATGGAGTCACACCTTTTCAAAAATTTAACAATCATCATCGATATGC  
ACAGCCTTCATGTGTAGTGTATGCTCCAGCTACAGCTGTAGTTACCCAATCTCAAAGCA  
AGTAAACAGCAAGATTCACACTAGCTCTTAAGTGGCCAAGCTATATTTCTATAACTAGA  
ATTGCTATTGTGGATTTCATAGTTATAATAACAGGATAAGACCACTTTATCCATGTA  
TTCTAGTGACTTTTTCTTCTATAGCAAAAAGAAAAATACATCTTTCACCATTTACAAGT  
ACAAATTTCAAGGAGAAATTTTAAAAGGAGAGTAACAACTGTCTGAGTTGCAGCAAGA  
CTCCTGAGAGTTCATTTCTGGGCCCTCTGCTGCCTGTTTTTGGCATTGAACCCAGGAA  
TCTTTTCTAAAGCACACAGAAATCTTGCAAAAGAGGCCATTTCTAGTTAGGCTTTTGTCC  
AACTGTCTAGTTAAATAAAATTAATTTCTAGATTACAAATGTGCTTCAAAGGTTTAAACA  
AATTGAAATGTCCTTAAGTATTTCAAATAAATTAAGGAAGAATTCCTATCCCATAGTCT  
TCTACTTTCTCTTCCACACCTATGATGAATGTCTGAAAAG

The following amino acid sequence <SEQ ID NO. 164> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 30:

FQDIHHRGCRGKKTMGMGILPFINTGHFNLLNLSTFCNLRIILDSTWKALEMASFARFLCALEKIPGFNA

KNRQQRQEMELSGVLLQLRTVCYSPFKISPNLYLMVKDVFFLLLEEKVTRIHGSGLIVLLLMEIHKQFLK  
YSLASELVWNLAVYLDDWVTTAVAGSIHYTRLCISMMIVKFCEKVLHLCSL

The following DNA sequence Seq-2386 <SEQ ID NO. 31> was identified in *H. sapiens*:

CCCTTTTTTCTGCTTTCAGTTTGATTGATTACACCTTACAGGCTTGGTATGATAAGTTT  
AAAACATATTGAAGGTTTATGTACTTATAAAAACCTCATCATTCCCTAAAGAAAAAAAT  
CTCAATTTGGTTTAGTGTCTTGTAGTCTTGCTTTCTACATCTTACTAATGTCTCATTTA  
TTTATTCATTTTGCTCTGTACATTTAGAATGATTTTGATGGGCAAAAATCATGGTAGTT  
ACAAACAGCCCTTTAAACTATTGTTATACTTTGTTTCAGTGGATTCTGGTAGAGGCTTTA  
AGGTAATTATTTCTTTAAAGCATTGTGTAAATATACCTCCTACTGTAGTGCCCTTGGGAA  
CAGGCAAAATTCAGAACTGGCCTGCTAGCAGTCTTACCAGGGTTATAAAAGTAAGATTAT  
TATATATAAAACAGCATTAACTCAATGCGTGGTGTGTTGCAGCTGGCAACAACCTCGCT  
CCCCAAGCTGCTAAATTCGTGGTCTTATGAATGTCTCCATTGCTGTGTTTGCTGTAAACA  
GAAGTGGGAGGGTGTTCCTCAGTAGCCTTGACTGTTTACCAATGCACACTCC

The following amino acid sequence <SEQ ID NO. 165> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 31:

LFSAFSLILHLTGLVNIILKVYVLIKTSSFPKEKKSQFGLVLSLSCFLHLTNVSEFIYSECSVTFRMILMGKN  
HGSYKQPFKTIIVILCSVDSGRGFKVIIISLKHCVNIPTTVVPLGTGKIQNPASSLTRVIKVRLLYIKQHLN  
AWCVAAGKQPRSPSCIRGLMNVSIADFVAVTRSGRVFPSSLDCLPMHTGVCIGKQSR

The following DNA sequence Seq-2387 SEQ ID NO. 32> was identified in *H. sapiens*:

TTATTTGGTGTTAATTTATAGGCTCAAAGGTCTAAGGTGCCCCCTGTGCGGTTGCCT  
GTGGTTCTCTTTGCTCCTGTCTGCCCTCTTGGGCCCAATACCTAGTATTGTGCTTAGGAT  
TCACAAACGCAACAAATACTTACTGAGCACCTACTCTGTGCCAGGTGCTGTGCTATATGC  
TGAGAAAACAAATGTTAAACAAGATGGATAAGGTTTCTTCCCTATGGTGTCATAGTCTA  
GTGGCAAGACAGGTAATAATGACTCAGTGTATTCTACTAAGGACAAGCATATCGTGCTA  
AGAAAACCTGTGTGGGAATGGGTGAGGGAAGGTATCCTTGGAGTAGCCCCGTTTGAAGTG  
GGATCTGAAGACTGAGAGTTATCTAAGTGGGGAGAGCATTGCAGGCAGGGGGATCAGCAT  
GTGCAAGGGTTCTCAGAAAGGAGGGAGAACATGTGTAAGAAATATCACTGTAGTTGCAA  
CCAG

The following amino acid sequence <SEQ ID NO. 166> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 32:

IWCFHRLKGLRCPVAVACGSLCSCLPSWAQYLVLCIGFTNATNTYAPTLCQVLCYMLRKQCTRWIRFSSL  
WCPSSGKDRLSVFYQAYRAKKTGVMGQGRYPWSSPVTGIRLRVIVGRALQAGGSACARVLRKEGEQCVR  
NITVVATQ

The following DNA sequence Seq-2388 <SEQ ID NO. 33> was identified in *H. sapiens*:

TCATTATTATAAGAATTATAAGAATTCTGAAATATTAGCCTTAAATAACCAAGTTAATA  
AAGCTTAAACTTTTTATGGAATTATCCATTCTGTTTGAATAACTGAACTCTTTCA  
ATACTATTGCTTGTTCACTTAACAATGATTACTTGAACATAGTTTCAGCTAAAGCTTTA  
TGATATTCCTAATCTAGCATTATTTTCGCATTGCTTTCCACCATCACTAAAGTAATTA  
CTACATGTTTACCAACTAATTATCTGATGGTGCAATTAAGAATTGATCTTACCTTAATA  
TTTTATGGTATCAAGTGTTTTTCATTTCATCAAGAATATTCCATTTTGCTTATATTTAA  
TGATGAGCTCTAGAATATCATCACTAACATATCTAGCAAATTATAAATATGTCATTTT  
AGGTAAATATTTAAGAGTATGTAGTGCTATATTTAGTTATTTTAAATCAAATACTTA  
ATGTTTATACTTTTTAATTGATGTACAATTTCAATCTTTAGAATGCGCTTATGAAATA  
ATTGCCCTTATTATAGTTTATAACAACCTTAATATATCTTCTGTATCTATAGCAGATGA

TTTATAAAATGCTTTTCTTTATTAATAACTGTCTCTATCTCAAGTTCTTCATAGTGAGC  
TATTTTCTTTTGTATTCTGTAGAGATACATA

The following amino acid sequence <SEQ ID NO. 167> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 33:

IIIRIIRILKYPNNQVNKATFYGIIHFCFEKYTLFKYYCLFTQLEHSSAKAFMIFTNLAFIFALLSTITK  
VITTCSPNTNYSYGALRIDLYLNILWYQVFLHSSRIFHFAYILMSSSRISLTYLANYKYVIFVKYLRVCSA  
IYLVILNQILNVYTFMLYNFQFFMRMLNNCPYYSFITTLLIYLLYLQMIYKNAFLYLSLSQVLHSELFLEFV  
FLRYI

The following DNA sequence Seq-2389 <SEQ ID NO. 34> was identified in *H. sapiens*:

AGGCAGTAATTCAGTAATGTGATGAAGTAGCAAGAGATAAGTAAGTCCAGGTCAGTGAA  
GACTTCGTGGGGCTGACATATGAAGTGAAGAAATGCCACTTTTGGACTTTCAGTTAAGA  
CAAAAATAAACTTACCTCTTTTTTTTTTCCAGGTATCTGTACTTCCCTATTTTGCAA  
TACTTAATGGATACATACAATCTGTCAACTCTTCTCTGACCTGCGCATACACTGCTC  
CATCTGCCTGAAACAATCTTCCCTGGTCAACCGCTACCACTGCCACCTTGAGAACA  
GCTACTCATAGTCACCTCAGATTATATCGTTTCTCACCCTCTCATCTCTTCTTCC  
CGTTTCACCACCTCCCTTCAACCTTGGTGGGCTTGGCCATCTGTCTGCTTGACAGGACA  
CCCCATTGTTACCTTTGACTGGACTATTAGATGACATCTCAGTTACTTACCTTTTATGT  
GCTAGAATTAATTTCTAGCTGGAGTTGTCCCATGACCTGAAGCTGAGTGCTGCTCTA  
CCATGCAAGAAGCTCTATTGCCGAGGCTAGGCTGTTTGGGGGCTTCTCTAGCCAATG  
TGCAATGTCCCATTCCTAGTTGCATTCTGAAATATAACATCTGAGTTCACAGTAT

The following amino acid sequence <SEQ ID NO. 168> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 34:

YCELCRYISECNEWDIAHWLEKPPKQAASAIELLAWSRHSASGHGDNSSSEINSSTKVSNDVISSQRQGPCV  
KQTDGQSPRLKGGGETGRKMRWVRKRYNLRVTMSSCSPRWQVWGGPGKDCFRQMEQCMRRSREKSQIVC  
IHVLQNRESNRYLGKKKEVSLFLSLKVQKWAFFQFICQPEHVFTDLDLLISCYFITLLELLP

The following DNA sequence Seq-2390 <SEQ ID NO. 35> was identified in *H. sapiens*:

TTTCGAAAAAACGTATATGAAAGATTAAATATGAGTTATGATGTCTTTTTTTATCCCA  
AATCTGCTTTAATTATCATCTATGAGAACATTTTGGACATGCATGAACATACAAGTGT  
TCTATGTACCCTTCCACAGGAAGTATTAGAGGTTAAGCATCATTAGCCAAAAATGACTA  
GACAACTTCAATGAGAGGACTGATGTGAACATTTAAATATATATCAAGATAGATCTAAG  
GTTAAAAATTATGAGAATAAAATTGGAAGAACAATGTATCAACGTTATGCTATTCAAAA  
CTAGAAATAATGCATGTAACAATGGGAGAAGAAGGGAAGTAAAAAGACAATTGTAAA  
AGCAGCTTATTTGGATAGCAAATGTATGGGAAGTAAAGTACACACATTAACTTGGCAAAC  
CAGCAGATAAGAAGTTACATAAGAATATAGATGGCTAATGACATTATACGTATAAATAG  
GCCTTAAAAACAAATATTAACCTTT

The following amino acid sequence <SEQ ID NO. 169> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 35:

KVLIFVLRPIYTYKCHPSIFLCNLSAGLPSLMCVLYFPYICYPICTFYNCLEFYFPFFSHCLHALFLVLNS  
ITLIHCSSNFILNFPYIYLDIYLVHISPLIEVCLVIFGMMLNLFWKGTNTCMFMHVQKCSHRMIKADL  
GKKTSLIFIFHIRFFE

The following DNA sequence Seq-2391 <SEQ ID NO. 36> was identified in *H. sapiens*:

GGCCGCCCAGGTGAGGAACCGTGGTCTAAGTCCAGCTTTATTCTTAGTTGGAGGAGTG

GCCTTAGGTATGTACAGGGCCCCCTTAGGCCTTTTGGTTGTCGTTTTCATAAAAGGCAGC  
 TTGTCTTGCTGCTGACAATCATCTTTGAGAGTGTTAGACTTAAATGAGATCCTGCAGTAG  
 TTTTCACCCCTCCACAGGTAGCAAAATCTTTACTCTAAACAAATGTACTTGATTCCCTGA  
 TGCTAAAACAAAAGAAAACCTGGAATTTTATTACTACAAACATATTCTATAAGCCCTCA  
 TGTATATTTTTTACTTTTCTTGGAGCCCCCTCAGTAAGAAAAACAAAACAGCTTTTAATAC  
 AATGTTTTTACAATGGCAAAGTTCAAACACAGACAAAGGTAGAGGCAATGGTATGATAAA  
 GCCCCAGGCATTTCATCACCAGATTCAATAATTACCAATTCAATCAACCCAATTTAG  
 CTCTCCACCTCACACCTCACTTTTAAAAGACAGATCCTCCCTCATTAGATTAGTTCATT  
 CACAAATATTTTATATGATCTTGAAAATATAAGTGCTCCTTTAATCATTGTGATATCAAA  
 TTCAAAATTAACATTAATTCTCAAATAAATAGGGCTATTTTGATG

The following amino acid sequence <SEQ ID NO. 170> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 36:

HQNSPIYLRLNVNFEFDITMIKGALIFSRSYKIFVNELIGRICLLKSEVGGELKLGLIGNYIWMNAWGFI  
IPLPPLSVFELCHCENIVLKAVLFFLLRGSKSKKYTGLEIYVCSNKIPGFSFVLASRNQVQFVSKDFAT  
 CGGKLLQDLIVHSQRLSAAARQAAYENDNQKAGALHTGHSSNESWDLDHGSLTWAA

The following DNA sequence Seq-2392 <SEQ ID NO. 37> was identified in *H. sapiens*:

TTTGAAAGTACATGTATACTAATCTACATCTAGCATCAAATAACTACCACTTCTCTCCTT  
 CCTGTTTATATCATTACTGCCTTTTATTTTCATTTATCCACATGCTATAATCACTCAATAC  
 TTTGTTACTATTATTGTAAACAGTTATCTTTCAGATCAGTTAAGAAAAATAAACTTAAT  
 TTTACCTTAATATAGTACTTTTCTAATGCTCTTCCCTTTTTTATGCAGTTCTTTTTGACAT  
 TTCTCATAGGGCAGGTGAGCTGGCAATGAATTATCCAGTTTGTGTTGTGAGAAAATATC  
 CTTATTTCTTTGAATTTGAAGGATAATTTTGCTGAATGCAGAATAATAGTTTGGTAGCTT  
 TTTTGTGCAACACTTCATGTATTCTCCTTTCTTTGTGTTGTCATGGTTTCTGAAGAGAA  
 AGATAATGTAATTCTTATCCTTTTCTCTATGGATAAGGTGTTGGCTTTTCCCCCTCTC  
 TAGCTTCTTTCAAGATTTTCTCTTCTTTGGTTTTTTCAGTTTAAATATGATATGCCT  
 GGGTGGAGATTGGAATTATTAT

The following amino acid sequence <SEQ ID NO. 171> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 37:

LKVHVLIIYIHQITTTSSFLFISLLPFISFIHMLS~~LN~~LTLLLTIVIFQISEKNLILPYSTFLMLFLFYAVLF  
DISHRAGQLAMNYSSFVCQKISLFLIRIILLN~~AE~~FGSFFVATLHVFSFLCVMVSEEKDNVILILFPLWIR  
 CWLFPLSSFFQDFLFSLVFCSLNMICLGGDLDDL

The following DNA sequence Seq-2393 <SEQ ID NO. 38> was identified in *H. sapiens*:

ACTTCTAACTGCTGGCTTTAATTTAATTTAATTTAATTACAGCATTTTCCACACATGCCC  
 ACAGGCTCTTGGTAATAGTTGCATTTTAAATAATCTAATATATAATAATGACTTTGTTT  
 TTAATTTTCCACTGAGAGTTGGATCCTGAGTTGAACACAGAGCTCCAGACAGGGGCGTCT  
 GGTTCACTCCATGTGATTGGATTTAGGGAACCAAGGGGCTCCTAATTGGAATAAGCTG  
 TGCTTTACCCCCCTATCCCCACACACCTGTGTTAATGTCTCAGCAAGCATCCCATAGG  
 ACATGAAATGACCGCTTGTTCAGTCAAAATGATCAAACAGTTGAGCAGGCATTCTCA  
 GGCTGGACTGTGAAAGGAAAATGAGGTAAGCGAGCAATGCCTGGCCAAGACCATTATAC  
 AAAGAGACTCTATGGACAGCACTCTGGTGGTGGCCTTTACGGAGTGACCCACTGCTCTCT  
 GCCTTTATCCACAAGTCACTGGGCCAACTAGAACTGTAATCAAACATAGTTCAACCAAA  
 GGATGAATTTTATGACTACTGATTCTCCTTTGCAAAGACCGTGGTTGATATTCATCGGT  
 AGGC

The following amino acid sequence <SEQ ID NO. 172> is the predicted amino acid sequence derived from

the DNA sequence of SEQ ID NO. 38:

AYRISTTVFAKEKSVVIKFILWLNVLQFVGPVTCGRQRAVGHSVKATTRVLSIESLCIMVLARHCSLTSI  
FLSQSSSLRNACSTGLIILTETSGHFMSYGMLAEDIKHRCVIGGESTAIFQLGAPWFPBIIQSHGVNQTPLS  
GALCSTQDPTLSGKLKTKSLLYIRFIKNATITKSLWACVENAVIKLNIKASSK

The following DNA sequence Seq-2394 <SEQ ID NO. 39> was identified in *H. sapiens*:

CTCGAGCAGTAACCTGTGCTTCTACAATTATGACACCCACTCCAGGGATAGTCACTGCCA  
AAGGGTAGAAGCTGCTGGGGGCTCATTGCACTCACACAGACTAAGAGTGTAGCATCTCCCA  
GTTATGCGGGCATCAGGGCAACATGGGGAGAACAGTGGCAGGCACATAAGGCCACCCCCA  
GGTACAATGTCCAGTGCAGTTCACGGGTAGGTAAATCTACTCTGTGTCCCCACAGACCCA  
TAGACTCCCAGGGGGCACAAGTCAATCAGGGCCTGACCTTGGTAGTGACATGTGTTATG  
TTTGCAAAGGCTGTGACAGGTACCCATCCACAGTGGTGTACCCCAATGTTGCTCTATG  
CACTGTGGCACTTGGGCTGGGAGTACTACATGTTCCCACTAGCCAGCCCCATCATAAAC  
GCTATGGGCCAGCCAGGGGTTGGGCACACCATGTGTCTTGCGAGCATCCTTTGTCCAAAGC  
TGCCATGTTGCATTCCAGGCATCAGCCATGGGACCCCAAGTCTCCAACCATGTCCAGTT  
CTCTGCAGACACAAGATGTATGTGCCAAGGCAAGCCATCCGAGCCCTGCTGGAAGGGCA  
GTGCATATCCAATAGTTGGAAACATTGGTCACCTAGTGTAAAGGTGTGGGCCAGTCCACA  
ATGCAATTGGAGTATGTTAACCTCTGG

The following amino acid sequence <SEQ ID NO. 173> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 39:

QRLTYSNCIVDWAHTLHVTVNSNYWICTALPAGLRMACLGTYILCLQRTGHGWRLGGPMADAWNATWQLWT  
KDAARHMCPTPGWPIAFMMGLASGEHVLPQVPCIEQHWGNTTVGWVPVTAFAFANITHVTTKVRPLTLC  
PLGVYGSVGTQSRFTYPTALDIVPGGLMCLPLFSPCCPDARITGRCTLSLCECNEPPAVLPFGSDYPWS  
GCHNCRSTGYCS

The following DNA sequence Seq-2395 <SEQ ID NO. 40> was identified in *H. sapiens*:

AATTTTTTTTCACTACGGAACCTCGTTTGCTAATATAAATGCAGACTTTTTTTAAAAAAA  
AGCTTTATTGGAAACATGATGAAAAATGTGATGTATTAATACTTACTGATACTCCAAGA  
AAAAAATAATAAATATTTAGAAAGCTCCTCCCATCATTTTCTTTGGCTTTTAACTCTA  
CCAGATCTTTGAGAATGCATATTTGCTGGTTAACCAGATGAACCACCTTTCTCTACT  
AGTTCTGCAAGATTCAATATCATTTCATAGTCTCCAGCACTCTAGAGTAATCATTACTAGC  
TGTTAGGAAAAATTATGGTATTTCTTAAAAACTTTCTTTGTGACAAGTGAATAAACCAAAA  
GGATTAATAAAGATGTTCCAGTTTGGGAAAAATAATGCAATGAATACTGCATCTGATG  
CACCATTTAAGAAAGAGAGAAAAATAAATGCTCATTTCTAATTGTCCTCATTTCAGCAG  
CTTCCAAATATTCTTCTATTTCTTTCTTTTAAGTAATTACCACATTTTCATATTGCT  
GAATCATGAA

The following amino acid sequence <SEQ ID NO. 174> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 40:

FMIQQIKCGNYLKRKKKNIWEAAEMRTIRNEHFYFLSFLNGASDAVFIALFFPNWNIFFLILLVYSLVTKK  
VFRKYHNFNSLLSAGDYEYILQNGKGGSSGPATICILKDLVELKSQRKWEELSKYFIIFLEYQVLIHHI  
FHHVSKSFFLKKVCIYISKRVSVVKKK

The following DNA sequence Seq-2396 <SEQ ID NO. 41> was identified in *H. sapiens*:

CCCGAGTGACAGAAGCCATTTCACTGCCAGAGACTCTAGCGGCCTTCAGTTCTCTTGAG  
CTGGAGCCACTGGGTCTTGATGAAAGCTCACCAGGACATCTCATGTGGACCTCGGGCAT

CTGAGCCGGGACCATCTATTACAAGTGCAGAAACCAGATCATTAAATGCAGAGCTGAATT  
 CAAATTGTTACTTGCTAGCTTAGGAAAGAATCCTTGGAATCCAACATATTGTCTAAATG  
 GATCAGTTAATCTTACTATGTGCATTCTACATACCCTTTCATTGTTTGGGCTTAAATAAC  
 TTTTCTGCTTTGTCTGGTTAAATTTATCCAATGTGGATCGCTGGAAGAATATGATGTAT  
 GTTTTAGAATAGAAACAGTTCTGAGATGAAGTTGAGCACAATTCCTGTTCTAGTTGCAA  
 TTAAATATAAATATAGCATTTGACATAAATAGCTGGCCCGATATATTTAGAGTACAAGT  
 TAAGTGTATCCCTTAGAATTGGGCATTGACTCCGTAGAATTCCTTTGTACAAGGTG  
 AGCAAATGTATATTTTGTAAAAATAAGTATCTGACTGCCAAACGGACAGAAAGCTCTT  
 TGCCATATGTGTTTTCA

The following amino acid sequence <SEQ ID NO. 175> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 41:

ENTYGKELSVRFGSQLIFNKIYICSPCTKGNSTESMPNSKGMTLNLYSKYIGPAILCQMLYLYLIATRTG  
 NCAQLHLRTVSILKHTSYSSSDPHWMKLNQTKQKSYLSPNNERNVCRMHIVRLTDPFRQYVGFPRILSASKQ  
 FEFSSALMIWFPHLDGPGSDARGPHEMSWAFIQDFVAPAQENRPLRVSGSEMASVTR

The following DNA sequence Seq-2397 SEQ ID NO. 42> was identified in *H. sapiens*:

CTTTTAAATTTTGTGTTTGTAGCAGTTGTTGTATCCATGTGTGTTGGTGCCCATATGTA  
 TTGTTTGGGGTTTGGTTATTCTCTCAAACCAAGTACCGTAAAAAGTTGAATTTTGT  
 ATTTCTTTATTGAGTAGTGGGACCGTCTAGACTGTGTGCTGACTCTTACTAAAGTCATTT  
 GTTTTTCTTACCCGTGGAGAGGTGTATTCTTGAACCCCTTAAACGGGTCTCTACTTTGGC  
 CTAAGACCATATTAGAAAACCTTTTTGAAGTCACTTATTATATGCCATATAATTAAAAAG  
 TTATATGGTATATTCTCCATTACATTTTAGCCACAATGCCGTATATTAAATAAGCAAA  
 CAAACTATATGTGGCAATTAACCTTAAAAAAGCCTGAATTGGCTCTTAGAAATAT  
 TTAATCAAGTAGTATCCACTAGAACTTAACATTTTATCCTGTGGATCATCACACAAAA  
 TACCAACCCCTGCTGTCACTCAGGGTCCTAGCAGGAACAGGTAGCATCAATAGGATAAT  
 TGATGAGAGCTTAAGAAAGGAATATTTACAAATATGTGGCCAGATTAGGGGAAACAGT  
 AAGGTTGGGAATGCCGCCAGGATTCTAACAAGAGTGAGAATCTATTTCTACT

The following amino acid sequence <SEQ ID NO. 176> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 42:

LFNFVFVAVVCIHVCWCFYVLFVWLFSONQVTVKSLNFSISLLSSGTVTVCLLLKSFVFLTRGEVYSTLT  
 GLYFGLRPYKTFKSLIICHIIKKLYGIFSHYILATMPVYISKQTCGNNLKKKAIGSKYLIKYPLELNIS  
 SCGSSHTKYPTLLSFRVLAGTGSIKDNELKKGTIYKYVARLGETSKVGNAQDSNKENLFL

The following DNA sequence Seq-2398 <SEQ ID NO. 43> was identified in *H. sapiens*:

TCCATGTAACATTGATGAGCACAGTTTTCTCTTCTGTAGCAAGCACTCCTCTGCCTAATT  
 CATATGACTAAAAACAGTGCTTCTCAAACATATGGTCTCAGGAACCCCTTAAATCTTAAC  
 AACTAGTAAATGACCCCAAAAGGTTTTTATAATATGAATTTATATATAAATATTTTAT  
 TGAAGTCCACTTTTATGAAAATAACCTTTTTCAAAAATTTATAAGAAAAAATAGTA  
 TTATTTTACATATTTGAGGCATCTTTTAAATGCCTGGTTTAAATAGAAGACAATTGAATAT  
 TCATGTCAACTCTGGATTGATCTGTTTCAATATGTGTCTTTGGTTGAAATACATGAAG  
 GAAACTTGGGATCATCAGACATATAGTTAGAAAAGGGTGGAGTATTTAACAGCCTTTTT  
 GGACAACTGTGACATTGTGCTTTGATATTACAACAAACTGGAGAAGTGGTAGGTTCTA  
 AATGATTAGTTGCAACATGGAATCTGAAACCATCATGAACATTTGTAACTGGCATA  
 TTAAGATCTATTTATCTATCTTGCACTTTGAATGGGATCCTTTGCTCATGCATCTTTTG  
 TAACATGAATCATCTCAAACACGTTGGTTTATTGAGTTATGC

The following amino acid sequence <SEQ ID NO. 177> is the predicted amino acid sequence derived from



the DNA sequence of SEQ ID NO. 43:

HVTLMSTVFSSVASTPLPNSYDNSASQTYGLRNPLKSQLVMTPKRFFIIILYINILLEVHFYENNLFISKIS  
EKNSIILHIGIFLMPGLIEDNIFMSTSGFDLFQYVSLVEIHEGNLGSSDILEKGGVFQPFWTTVDIVLYYN  
KTGEVVGSKLVATWNLKPHHELFIWIKIYLSILHFEWDFLLMHLEVTIISNTLVHVM

The following DNA sequence Seq-2399 <SEQ ID NO. 44> was identified in *H. sapiens*:

AATTAATAATCCCTGCAGTCAAATTAGACTCTGCATGTCTGGGGATATTTAAAAGGATAAT  
GTATAGGGGTGCGCATGGTAACTCATCAAGTGGTAATTCTGTACCTTTCTGAGTGAAAAC  
CTTGAAAGGAGAAGACAAGCAATTGGGGAGATAACAGCACCAGAAATTGAGTTCATCTG  
TAACCTAGGCTCTCTGTGAGTTTGTACCAGCTATTACCATGTGGATGAAAAACAGTA  
AAAAGACAAAAAGATTACATTTCAAGGCTCCCTAAAATTGCCAATTCCTACTCTATAGC  
TGATTCTCAGCACAGGAGGAAATGGGACTAGAATGCTGGGAGATGACACTATCATCGAAC  
AGTCAGCTCCAAGGAGAAGCCTAATTGTTACTTCTCAATGGCAGAAGGCGGGTGCTTCCC  
CCGGGCGAGGATTCTGTTAATCCTTAGGTTAGAGCCAGCTTCAACCCAGTGTACAGG  
TCAATTACCACCTCCAACCTGAGGGGCGACATGAACCATACTCAGCAGCGGCGCATG  
CTCCCTCCTCAGCACCTCTGTACATTGAGAGCTCCTGCATGGGATGCCGAGAACTCACA  
CCCTTCAGGGCTGCTGAAGATCATATGACTGATCATCACTTTGATTTTGAACCATCT  
GTCAACAACGACAC

The following amino acid sequence <SEQ ID NO. 178> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 44:

IKIPAVKLDLSACLGIKFRIMYRGCHGNSSSGNSVFPVKTLKGEDKQFGEITAFIEIFCNLGLSLVCLPAIH  
HVDEKQKDKKDSHFKAAPNCQFHSIADSQHRRKWDNAGRHYHRTVSSKEKPNCFYSMAEGGCFPRGRILFNP  
VRAQLQPSVTGQLPPSNPEGRHEPYRSTGACSLSTCTFRAPAWDAENSHPSRAAEDHMTDHLFLTHLS  
TTT

The following DNA sequence Seq-2400 <SEQ ID NO. 45> was identified in *H. sapiens*:

GCCTAACTGAATTATAACCGCGAGTTGACAGTGGTGAGCATAGCTGATGAGATGCAAG  
CAAAAAAGAGTATTGCTGACCTAGGACCATGAGGAAAAACCAATCCAATTAGTCAAG  
TTGGAGGACATTTGTTGAAAACCCACACTTCCATGAGGTCTGTAGCCTTGAGCCTATCA  
GTGCCGACACAGAATCTGAATAGTTCAATGCCTCTTTCTGTTAAAGAGGAGACGCCT  
CACTCTGCCGCTCAATCTTGGACTTGTGTTGTGACAGAGGTCTTGCTTATGTAACACTC  
GCTTTTAATACTATAATTCACAGAGTCCTTTGAACACATAAAGGGAAAGCCACTTTCGCTCC  
TGTTAAGGATGTATAAGCACAAAAATGAACAGTGAATTAATCCTAGTGTATTATACATT  
TTTTTTTAAAAAAGAATCTAAGCCAGAATGAGGTACTGCCTAGGCAAAGAAGAAGACA  
GCTCATCACAGGTGAGTGTAAACAGTTTTTCATATGTACAAATTAAGCAGCCTGAAACAA  
AAGGCACTCAAAAGGTAAAGAATACAGTCCACCCCTCTGATTTGTCAAATCAAAGTTC  
TGTCAACTG

The following amino acid sequence <SEQ ID NO. 179> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 45:

SQNFDLTNQRGGLVFFYLLSAFCFRLLNLYIKTCYTHLAVFFFAAVTSFWLRFFFKMYKTLGLIHCSFFV  
LIHPQERKWSLYVFKGLCELLKASVTARTSVHKQVQDAAGVSSILTERGIELFRMFCVGTDRLKATDLME  
VWSFQMQSSNLNLDLVFPHGPRSAIFFCLHLISYAHHCANSRLFS

The following DNA sequence Seq-2401 <SEQ ID NO. 46> was identified in *H. sapiens*:

AAAAAAAAAAAAATTCAGGGGAAAAAAGCAATTAAAAAACATACTATAAAAAATAATAC

AAATTACAAAACAACCATTTACATAGCATTACATTATATTAGTTATAAGTAATCTAGAG  
 ATGATTAAAGTGACGGAGGAATGTGCATAGGTTATATGCCAATACTGCCTCATTTTATA  
 TGAGGGACTTGAACATAGAAGGGTTTTGGAGTCCACAGAGGTCCTGAAACCAATTTCCCC  
 TTCCCATGCCTGGGATGACTGAATTATACAGCAGCAAAAATGAATATACTCAAGCTATAT  
 GCATGAGTCTCATAAATATAATGCTCACAGAAAAAGCAAGTTGCAGAAGGGTAAATACG  
 GTTGATATATAAAGGTGCTAAACACAGAACTATTTAATGATATACGGATGCAGTAAAGT  
 ATAAGAAATGTATGCAAACTTACTTAAATTCAGGGTGTGGTTACTTGGAGTAAGGCGAA  
 TGTTTGGGATGTCAGTAGGTACCTGACAAATGGCAACTTAAC

The following amino acid sequence <SEQ ID NO. 180> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 46:

VAICQVPTDIPNIRLTPSNQHPEFKVCIHFLYFYCIRISLNSSVFSTFIYQPYLPFCNLLFSVSIIFMRLM  
 HIAVYSFLLLYNSVIPGMGRGNWFQDLCLQNPSMFKSLINEAVLAYNLCTFLRTLKCYVNGCFVICIIF  
 IVMFFLLFSPEFFFF

The following DNA sequence Seq-75 <SEQ ID NO. 47> was identified in *H. sapiens*:

AGCTAGGGTGGGCAGGAGTGGTCTCTGAGAGGTGACATTGAGCTGAGACCTGAATGACA  
 AGAGACCAATGTCAGCTCTCTTTAAGAAAGTTTCTTTGTTTTAGTGGCTCTCTCCATA  
 CTCTTATTTTAAACTCACTTAACATCAATATAAAAGTGCCTTTGCAGCAGGACACTTTT  
 AGGAGGTCTTGAGCCCCCTCTCCACAGCACTCATCTGTGTACAAACAAGTTGTTGCTAG  
 TGGTGTGGAGCTCGTTTTTCCCAAGCTTCACCTGGCATTACCCAGATCTGTTCAACCC  
 TGGGCATCTCTCTCCAGCTGGATGCTCACCAACTTGTTCGCCTCAGTTTCTGGAG  
 GAGCCTGACTCTATTTTGGCCCCCTTGAAAGAAAGTACAGGACTGGGTTGAGGCAGCTG  
 CTCACACTCACCAGAGGCTCCATATCTTGTAGGCCACACTGGCTGCCATCAAGAGCTGG  
 CAGTCTTGAGAAAGCAGAAAGCAGATGGTGGGTAGAGGAGCGAGTGATATGGAAGGGC  
 ACAAACAGAGGGTGAAGAGGCCACACACCACTAGGATGGTCCGATGGACCTGGCTCGG  
 GCTGTGTGCCTGTCTCTCATGAGGTTCTCTCTGGCTTGATCAGGCTCCTGACCATCAGT  
 GAATAGCACACCAAAGTGACC

The following amino acid sequence <SEQ ID NO. 181> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 47:

VTLCVCSLMVRSLIKPEENLMRTGNTARARSIRTILLVCGLETLCFVPFHITRSFYLTICFLLSQDCQLLM  
 AASVAYKIWRPLVSVSSCLNPVLYFLSRGAKIESGSSRNGRTSWVSIQLGGRDAQGTDLGNAKVKLGKNEI  
 QHHQQLVCTQMSAGGRGAQDLLKVSCKGHFYIDVKVNKSMERATKTENFLKESHWLSLVIQVSAQMSPLR  
 DHSCPP

The following DNA sequence Seq-76 SEQ ID NO. 48> was identified in *H. sapiens*:

CCAGGGGGAGGGGGGCACGGGCTATAAACGCTCGGCCGAGCGGCGCCGGCAGAGAGCCG  
 CCGAGCCCAGCACAGCTGCCCTCTGGACCTGCGGACCCAGCCGAGCCCCCTTCTGAGT  
 TCCACAGGCGCAGCCCCCGGGCGGTGCGGCGGAGGGTCCCCGGGGCGGTGCCAGGGCGC  
 AATCTGGAGGGCGGCCGGGAGGAGGAGGTGCGCGCGGCCATGCACACCGTGGCTACGTC  
 CGGACCCAACGCGTCTGGGGGGCACCGGCCAACGCTCCGGCTGCCCGGGCTGTGGCGC  
 CAACGCTCGGACGGCCAGTCCCTTCGCCGCGGGCCGTGGACGCTGGCTCGTGCCGCT  
 CTTCTTCGCGGCGCTGATGCTGCTGGGCTGGTGGGGAACCTCGCTGGTCATCTACGTCAT  
 CTGCCGCCACAAGCCGATGCGGACCGTGACCAACTTCTACATCGGTGAGTGCGSGCCGCT  
 GCGCCGCACCTGCTGCCGTCCCGGGGGGCTCCGAGGGCCGAGCGGCTGGGGCGCCCTCT  
 CGCGACGC

The following amino acid sequence <SEQ ID NO. 182> is the predicted amino acid sequence derived from

the DNA sequence of SEQ ID NO. 48:

QGEAGTGYKRSAAAAPAESRRQHSCLDPADPSRAPSVPAQPPGGRAEGSPGRCOGAILEGGREEVRA  
AMHTVATSGPNASWGAPANASGCPGCGANASDGPVPSRAVD~~AWLVPLFFAALMLLGLVGN~~SLVIYVICRH  
KPMRTVTNFIYIGECGPLRRRTCCRPGLRGPSSGLGRPLAT

The following DNA sequence Seq-77 <SEQ ID NO. 49> was identified in *H. sapiens*:

AAGTCGCCTGTCTTTGATCTGGTAGCCAGGCTGTGATGGCTAGCTTTAGGATATTTCCC  
TATATTTCTCTTGCTGTCAGGTTACCCCTTGGTATACCTGTAATTGATTTCCCAGTTAG  
AGAGTTTAGATGTGGACAGGGGAAGTACAACTACAGCTTAGTGCAAGATAAACCAAGGG  
TGTAAATATCAAGTTGTAAGTGAACAGAAATATTACCAATAGGATTTCCAAATGAACAG  
GATGGCAAAGAGTTCTGGGGTGTGGAAGTCAGAGTAGGTGCCAAAGGATCTAGATCAAAG  
GGGTTGGTAGATGAGCAGGGATGGGTGAGAGAAATCTAGGACTGTTAAAGCAAGCATGAC  
CCAGGCCATGTTCTGAGGTTGGTAAAGTGAATTATAGAAGGTGAGACCAATGTGAGATT  
GTGAGATTTTAACACCCCAAGAGGGAGTATGTGCCTCAGGCAAAGAAATGGGAAAAA  
AAAAACATGGTATATGGCATATTTGAGGAGCAAAGATAAGTTCATTGTCACTAGGGCAGA  
GCAAGGGATAAGTGAATGGTGTGAGACAAGATTGGAGAGGTTAACAGTGGCCAATAACAA  
GTGATAAAATAATTTTCAAATGAGAGCAGCCAGCACTTATAAAGTGGTTAATGTGCAC  
CAAGTACTGCTTTAAGTTATCTGCAGTATTATTG

The following amino acid sequence <SEQ ID NO. 183> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 49:

IILQDNLKQYLVHINHFISAGLLSFENYFYHLLLATVNLNSLVSHSLIPCSALVTMNLSELLKYAIYHVF  
FFPFSLPEAHTPSLGLKSHNLTFGLTFYNSLYQPONMAWVMLALTVLDFSDPSLLYQPLSRSGFTYSDF  
HTPELFAILFIWKS~~YV~~IFLFKYNLIITPLVYLALSCSLYFPCPHLNSLTGEINRYTKGPD~~SKRN~~IGKIS  
SPSQPGYQIKDRRL

The following DNA sequence Seq-78 <SEQ ID NO. 50> was identified in *H. sapiens*:

GCCTCCAACCGATATTTCTGTCTGTGCTCTGACCAGGTAAGTGGCCATCACCAATGCCC  
TGTAGTATAGTAAATGGGCCATCTCAAATGTATCTCTATCCAGTGCTCTTCTCCTAGA  
CCTCTTGACACCTACTCCACATGTAAGACCTTCTACATTTGGTTGTTGTTGTTATCA  
TCTTCACACATTGCCCAACAAGAACATCCAGAAGCCATCATCAGCACCACCTGCCAGG  
TCATCAGCTCACTCTTCTTCTCAACCCAGCCTCCATGAGAGGCAAAGGCGCTTAAC  
TGGCTCTCCTCTGCTTGTAAATCACATGAAATCAAGCATGCTTATAGTGTCTTAGTACA  
ACAGGAAATTTACTTTCAAACAAGGAAAGCCACAGAAACCTGGGGATCATTTTAGGGGC  
TTTATCATCTGCTGGCTGCCTCTCTTTATTGTTTCTCTGCCAGCCAAGATACCACCATA  
TTAAGACATCTTCATCTGCTGAGCTTTTTTTTTTTTCTTTTGATACCAAGTCTCAC  
TCTTGTCTCCAGGCTAGAATGCAATGGTACAATCTCAGCTCACTG

The following amino acid sequence <SEQ ID NO. 184> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 50:

PPTDISVCCSDQVLGHHCQPVVMGHLKLYLPSALLDLLHLLHMDLLHFGCVVHHLHTLPNKNIQKPSS  
QHCPGHSSSLFFLNPSLHERQRLTGSPLLVNHMKIKHAYSVLVQGEIYFQTRKATETLGIILGAFIICW  
LPLFIVSLPAKIPPYDIFILLSSFFFFFLIPSLTLVSQARMQWYNLSSL

The following DNA sequence Seq-79 <SEQ ID NO. 51> was identified in *H. sapiens*:

CAGGCGCCTCAACTGTTCCACAAACCAAGCCTGAAACCAGAACTCCAACCTCTAGTCTGA  
AAAGCAAAGTGGCACCTCGCAAACCCCTGTGGCCCAAGTAGTCTCACCAACCTTGGG  
GAAGAAGCAGAATTCAAGCTGTAAGTGCCTGTTGGAGAGAGCAACCCCTCGGCCTCTGTC

CTCGAAAGGCAGCACCAAAGTTTCCAAGTGAATCAAATGTGCAGGGAGGATC

The following amino acid sequence <SEQ ID NO. 185> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 51:

ILPAHLIPLGKLWCCLSRTEAEGWLSPTGSYSLNSASSPRLGETTWGHRVFARCHFAFQTRSWSSGFRLGL  
WNSGA

The following DNA sequence Seq-80 <SEQ ID NO. 52> was identified in *H. sapiens*:

CTGTACCTGTCACAGTTATCAAAAATTTATTTCATTGAGAAGTCTTTGTTGAACACCTGTT  
ACGTGTACTGAGCATTGTCCTAGGTATTTGAGATACATCAGTGAACAGAGGATCCTTAAC  
AGACAATATACATAATAAGTTATGTAATAGCTTACAAAGTGACAGTACCTTTGGGAAAAA  
GGAAAGGTATTATAGGATAAAGATGATCAATGAACAGGAAGTTGTCAGTTTAAATTGAG  
TGGTCTGGGTAAGGAAGATCATACCTGAACCAAGACACAAAGGAGGTTAGGGAATGATGA  
GCCCTGCA

The following amino acid sequence <SEQ ID NO. 186> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 52:

CRAHSLTSFVSWFRYDLPYPDHSINCKLPVHSSLSYNTFPFSQRYCHFVSYYITYYVYCLLRILCSIMYL  
KYLGCQSVHVTGVQRLNEIFDNCDRY

The following DNA sequence Seq-81 <SEQ ID NO. 53> was identified in *H. sapiens*:

TAGCAGAGCAGGTGCTAGTGATATTGTCAGAACAGGTGCTGAATGAATGCATGAACAAAT  
GCATGAATGTGGAAATGAAAGGGGATGCAGATGGAGATGATGCAGATGGAGATGATGATG  
CAGATGGAGATGATGCAGATGGAGATGATGCAGATGGAGAGCAGTGGCCATGCAGAGTCT  
TTGCAGACCTTGGCTTGGCTTCAGGCTGTGGGGGCTCTGCAAGCCAAGGGTTTGAGTTCC  
ACCTCCAGTGCTTGCCAGCAATGCCACCTTGGGTGACCTTTATCTTGCTACCTGGAAAGT  
GGGGATGCTGGCAGCCCTCCCTCCTGGCATCACTGACACTGCATGGTCAGGGTGTGATC  
CCTTTGGGTACAGCGGGGGTGGTGGACCTCCAGGTGGGAGGTCCAGTTTGGATGAAA  
GGCCAAGGACGATTATAGGAGAGCACAGGAGTCTTGCTTAGCCCCAGCAATTCCACAG  
AACCTGCTGTGAACTGCTGGCTGCTGCCGTAACCTTTCCCTGCTCCCTATTTCCACTCCT  
TGGAGGCCGCAAGAACAACCTGCTGGCTGGCCTTGGCCACTGCCT

The following amino acid sequence <SEQ ID NO. 187> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 53:

AEQVLVIFAEQVLNECMNKMCMNEMKGDADGDDADGDDADGDDADGDDADGEQWPCRVPFADLGLASGCGG  
SASQGFEFHLQCLPAMPWPVTFILLPGKWGCWQPLPPGITDTAWSGCDPFGYRRGWTSQVGRSSLDERPR  
TIHRAQESLLSPSNSTEPAVNCWLLPVTFPCPYFHSLEAARTTAGWPWLP

The following DNA sequence Seq-82 <SEQ ID NO. 54> was identified in *H. sapiens*:

AGTCTTTTTCTTAGGGAACCTTGTGTTGCTTCACTATATAGTTGTTGTTTCAACAATT  
TTGTGTTGTTTACAGTTTCACTGTGACAGTTTGATGTTAGGTTGATTCTTTTCTCCT  
CTGTATAAAAGATTATGTCACCAGAATCTTCTTCACTTTGGATAGGACCTAAAGGA  
CCCTCTCAATCTGAAAATCTATGCTATTTGTTATCACAGAGCAGTTTTCTGCTGTCAATT  
CTTTGATTGTTACTTTTCTATTTATTCCTTTTCTTTTCTAAAATGCCATTATTTGTAT  
ATTGGAGTCATAGATCTGAGATCTGTGAATTGCTATTGATGCTCATATCTTTTGTCAA  
ATGTTTCCATGTCTCCAAGTCTTTGTTCTCTATTGTGAGATATTATTTGTATTGTTTG  
TCCAGAATATTAATTTAGTTCTATTGATTGACTATTCTTTGGTTTTGCTGTTGAATTTT

AAATTCAGGAATAGTGTGTTTTCTTTTCAGATTATTTTTCTGTGACCTAATTGCATCT  
TCTTACGGGGTCTTATTATA

The following amino acid sequence <SEQ ID NO. 188> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 54:

SFSLGNFVVASLYSCCFNNFVLPHSFTVTVCVDSFSSSVKIMSPESSFITLDRTRLISIKSMLFVITEQFS  
AVISLIVTFLFIPFSLSKMPLFVYWSHRSEICEFAIHVSYLFANGFHVSKSLFSIVRYYLYCFVQNINLVL  
FIDYSLVLLNFIQECVFLSDYFFLPNCIFLRGLII

The following DNA sequence Seq-83 <SEQ ID NO. 55> was identified in *H. sapiens*:

GCCCAGGGAAGCCAAAAGATTGGACATCCATGCTCCCCTCCTCTCCCTCCCGACTGCCA  
TCTCTTGATGGCGGCCAGTGTGGCCTACAAGATATGGAGGCCTCTGGGGAGTGTGAGCAA  
CTGCCTAAACCCACTCCTGTACTTTCTTTCAAGGGGGGCAAAATTTGAGTCAGGCTCCTC  
CAGAACTGAGGCAGAACAAAGTTGGGTGAGCATCCAGCTGGGAGGAAGAGATGC

The following amino acid sequence <SEQ ID NO. 189> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 55:

PREAKRLDIHAPLLSLPDCHLLMAASVAYKIWRPLGSVSNCLNPLLYFLSRGAKFESGSSRNGRTSWVSIQ  
LGGRD

The following DNA sequence Seq-84 <SEQ ID NO. 56> was identified in *H. sapiens*:

TCCTTGGTCATTTTGGTGTGCTATTCACTGATGGTCAGGAGCCTGATCAAGCCAGAGGAG  
TAACCTCATGAGGTACAGGCAACACAGCCCGAGCCAGGTCCATCCGGGACCATCCTACTG  
GTGTGTGGCCTCTTCACCTCTGTTTTGTGCCCTTCCATATCACTCGCTCCTTCTACCTC  
ACCATCTGCTTTCTGCTTTCTCAGGACTGCCAGCTCTTGATGGCAGCCAGTGTGGCCTAC  
AAGATATGGAGGCCTCTGGTGAGTGTGAGCAGCTGCCTCAACCCAGTCTGTACTTTCTT  
TCAAGGGGGGCAAAATAGAGTCAGGCTCCTCCAGAACTGAGGCAGAACAAAGTTGGGTG  
AGCATCCAGCTGGGAGGAAGAGATGCCAGGGTTGAACAGATCTGGGTAATGCCAAGGTG  
AAGCTTGGGAAAAACGAGCTCCAACACCACTAGCAACAACCTTGTGTACACAGATGAGT  
GCTGGTGGGAGAGGGGCTCAAGACCTCCTAAAAGTGTCTGTGCAAAGGACACTTTTAT  
ATTGATGTTAAGTGAGTTAAATAAGAGTATGGAGAGAGCCACT

The following amino acid sequence <SEQ ID NO. 190> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 56:

SLVILVCYSLMVRSLIKPEEPHEVQATQPEPGPSGILLVCGLETLCFVPFHITRSFYLTICFLLSQDCQL  
LMAASVAYKIWRPLVSVSSCLNPVLYFLSRGAKIESGSSRNGRTSWVSIQLGGRDAQGTDLGNKVKLGKN  
ELQHHQQLVCTQMSAGGRGAQDLLKVSCKGHFYIDVKVNKSMERAT

The following DNA sequence Seq-85 <SEQ ID NO. 57> was identified in *H. sapiens*:

GTCACACTGAATTAGGGACCACCCTTGTAACCTCATTTTAACTCGATTGTCTCTGTAAAG  
GCCAGTCTCCAAGTACAGTCACATTTGAGGTACTGAGGGTTAGGACTCCAATGTATCT  
TTTTGAGGGGACACAATTAAACCTAATAGACCACAATTAAATGGAATGCAATAATAAAA  
AACTAACTTTTATTGAGCATTCTGAGTCTGAGTTTGGCATTGCTCAAGAGTGCCTTACAT  
TAATTAATGTAATCTTCACAATCCTATGAACTCAGTATCATTATTACCCACATCTTACAA  
ATGAGTGGTTGGAGTCCATGGCAAGAGTAACCTGCCCAAGGTACGCTGCTGGTAAGATC  
AGAACCAGACTCAAAAACAGTAGTCTAATTCACAGCAGATTCCGTCAACAACTATTCTA  
CACAGTCTCTACTTTATGGGGTTCAACATAGAGACTATTTTGATGTCTGCGGTAGCTGTG

AGAATGTGGCTCAGAGACTTCCATCTATGGGGAACCAACCAAGGCCCCAGCTCC  
TGCACCTTTGAGACCTGTCACTATGTTATCACCGAGCCACATTTCCCATGGGCTGCTTCC  
AGCCAATGCCCAAACAATGGCAGGGAGACTAAGGCATCCTGTTCTGGGGAGATGTGGGA

The following amino acid sequence <SEQ ID NO. 191> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 57:

SHISPGTGCLSLPAIVWALAGSSPWEMWARHSDRSQSAGAGAFGLSSPMEVSEPHSHSYRRHQNSLYVEPH  
KVETVNSCRNLLWNTTVFESGDLTSSVTLGKLLPWPPTTHLDVGNNDEFIGLRLHLMGTLEQCQTQT  
NAQKLVFIIAFHFNCGLLGLNCVPSKRYIGVLTLTSTSECDCWRLGLYRDNRVKMELQGWSLIQCD

The following DNA sequence Seq-2337 SEQ ID NO. 58> was identified in *H. sapiens*:

ATTCTGTCTCTTCTCTCTGCGGCCCCATCTCCTGAGCCCAGCGAGCTCAGTGCT  
AGTTCACCTGTTTGTCTCCTTGTCTGCAGACACAGAAGATTGGGAGCGTTCCTGCCGAG  
GTTGGTAAGGATACCTGGAACAGTGGGCGCCTCTTGTCTCCCACTTGCTAGGAGTAAA  
GCCGTTTAAAAAGACACCTGAGCCTCTCCGGGTTCTGCTCCTCACTCAACCCACAGTA  
GATCTGGTGGGAGGTTGAGGGCTCAGTGAATCTGCAGGTGCAGCATCGTGTCTCAGTG  
TCCTGCCCCCTGCTTCCACCCGGTGTGACAGCTGCACGGTCCACCCACGCCTGCCTTT  
CCATCGTTCCTCATCAGCCCTGTGATCTTCTCTGTGGCCCTGCTGTGCTGGTGCCTGTG  
AGGTCTGTGGACACAAGAGACTGCACGGGCCACACCCAGCTGGGTGAGTCTCTCCC  
TCCTGGGTACTCTGGACAGTAAAGAAAGATGGACAGTGGGCTCCGTGGAGCATGAGGTA  
GTCCAGGACCTCGGCGGCCACAGGTCTGCCTCCCTGCTTCTCGTGCCTCCCTCCCTTT  
GGGTCTCTGCTCCACCTCGGTAAACGCTTCGTTCCACCCCTC

The following amino acid sequence <SEQ ID NO. 192> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 58:

ILSSSLCLRPSPPEPSELSSASSLFAPPCCRHRRFSGVPAEVGKDTWNSGRPLCSPLARSKAVKDTASPGSC  
SSLNPTVDLVGRLRAQICRCSIIVSSVSCPLLPVDSCTVHPTPAFSPFLISPVIFPVALLCWCVPVRSCGH  
KRLHGPHPQLGESSPSWVLWTVKKDGHVGSVEHEVVQDLGGHRSCLPASRALPPFGSLHLGKRFVPTP

The following DNA sequence Seq-2338 <SEQ ID NO. 59> was identified in *H. sapiens*:

AATATGTCTTAATATTCTAGTAGGGTTAATCTTTATTGCTTTTTTCTTTCTAGAATTTT  
TCTTATATTATTTTCATATAAATTTTGAATAAGTCTGGTTTGGGGGTCATATAGCAA  
TAGGTAAATTGATTAATAAAGTGATTTGGTGAAGTTTCACAATACATTATGAATCAAC  
TTCGGGAGAGTGGTTATGCTTATGTTTAGTCATTATATTTTAAAATGTGACATATCTTTC  
CATTTGTTTTAAGTCCTTGATCAAGCATTAGTTGCCTCCTCTGAGAATCTATAATTAAAT  
TCAAGATAAAATAATTTTTCCATTTATTGACCCATTTTTAGCTTACAATTTGTTTCTA  
CCCTTGTAAGTATTATGTTTGGTAAATTATTTTTATTAATATCTCCCTTACAGATATTA  
TACGCCATAAGGAAAGGAGTCACAGATTGGTAATAGAGACTCAATACAGTTTGTGGA  
ATGATGAAAGCATTATGAGGCATATTTCTTACTATGTTACCTAATAATCTTAAAGTTA  
TCAAGTTATTAAGTAGAGCCCATTCACAAGTCCAGATCTTTTGATTTTAAATCCTGTATT  
TTTCCATATTTTCAATATTTAATAGGGGAAGTAACATGCTAAATGCTATAGTTTGTCAA  
TTTTATATCT

The following amino acid sequence <SEQ ID NO. 193> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 59:

NMSYSSRVNSLLLFSENFYSYIIHFHINFRISLVWGVQVNLKIFGEGFTIHLINFGRVVLMFESHYILKCDI  
SFHLFVLDQALVASSENLLNSRNFFHLLTHFLTICFLPLVLCVNYFLNISPLQIYAIRKGVTDLVET  
QYTFVGMKALGIFSYYVHLIILKLSSYVEPIHKRSFDFKSCIFPYFYQLIGEVTCNAIVLQFYI

The following DNA sequence Seq-2339 <SEQ ID NO. 60> was identified in *H. sapiens*:

AAGGAAAATGGAACTAGATGAACGTGACAATATAAGACTTCCAAATCCACGTGGTTCCA  
TGAAAATAGGAAAACCGAATGCCAAAGGGCAGGCCACAGAAGGAGGAAGACCAGCGCTA  
TGAGCAGGATGGTCACGTACAGCCTGGTCAGTGGCATCTTCCGGGACCCACAAAGGATCC  
TGACCAGCAGGACCAGGCTGGAACACAGAGAGCCACACATAAAAAAATCAGCCCCCTA  
CTATGATGAAATCTGATGTTTAAACACCAACAGAATCAGCACCCTAAACAGGAAGTCAC  
AGAACTCCACTCCAGGACGCTCCGCAGCAGAGACAGGGCCAGAGCATGACACACACGA  
CCGCTGACAGGTGTAGGGGCGGGGCGGCAGCGCTACCAGATGGGCCACAGGACGTACAG  
GCAGCGCTCGGTGCTCATGGCGCTTAGAAAGCTCAGGCTTGCAAGGTAGGAAAACATCAT  
CACAGGGCTGAGGATTTTGAAGATGGGATGGAGGATATTGATGAGGCTTAACGGAAAACG  
TATAATGTGGCTTCTGAGAAAGAGGAAGTCGGCCATGGACAGGTTGAAGATGTAGATGGA  
GAAAGCGTTCCTGCGCATGCGGAAGCCCAGGAGCCAGAGCACGACTGCATTTCTGTCTAT  
CC

The following amino acid sequence <SEQ ID NO. 194> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 60:

MTGNAVVLWLLGFRMRNFAFSIYIFNLSMADFLFLRSHIIRFPLSLINILHPIFKILSPVMMFSYLASLSF  
LSAMSTERCLYVLWPIWRCRPRPYTCQRSCVSCSGPCLCCGASWSGVSVTSCLVVLILFGVKHQISSGGFF  
YVWLSVVPAWSWSGSFVGPGRCHPGCTPSCSRWSSSFGLPFGIRFFLFSWNHVDLEVLYCHVHLVSIFL

The following DNA sequence Seq-2340 <SEQ ID NO. 61> was identified in *H. sapiens*:

CACGACGCACCCCATTA  
ATGGGTTCCCTGGGGGCAGGGCATCAGTCCCACTCACTGCTGGGCCTCCAGGGCCTGCCA  
AAGGGGCAAAGTCACACTCAGACATAAACTCTTGTTTGTAGCAATCCAATAAACAGTCAT  
GAAACTAAGTGAGGAAAGTTATTAGATTGAAGGATTGAGGGAAAGTCCCATCAAAAAG  
TAAACTTGATCCCACCTCCACTTCTTGATGAGTTACTTAATCTCTCTGCCTCAGTTT  
TTTCACTATAAAATAGAAACCATGAGAGGACCTACCTCACCAGGCTGTTCTAAAGTTAA  
ATGAGTTAATTCTGTACAAGCTGAGAACAGCATCTGATACAGTATCTAATAAAGTCAGT  
TATTATTACTTTTATTATTATTATGTACTTGGTTATCATTATTTTCATTATCAATTATT  
ATTCTCTTCACTCTTTGCTGCCACCTGGAGTTCTCTGGAACCCCTTACGGCGTACAGCA  
GGGAGACAGGGGAGGGCAGATGCCATTTGCACAGCCATTGGGACTAATAAGCCCCAGCAC  
CCC

The following amino acid sequence <SEQ ID NO. 195> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 61:

HHTHTHTHTHTHTRTHPINGFPGRASVPLTAGPPGPAKGAKSHSDINSWFQSNKQSNVRKVIRLKGFE  
KSHQKVCLDPTSTSWMSYLISLASVFSPIKKPEDLPHQAVLKLNELIPVQAENSIYSISQLLLLLLLCTW  
LSLFSFINYYSLHLFAATWSSWNPFTAYSRETGEGRCHLHSHWDAPAP

The following DNA sequence Seq-2341 <SEQ ID NO. 62> was identified in *H. sapiens*:

TAATGTGGGACTAAAAAATATTAAAAAATAATGACTTCAACCTTCCCAAATTAGGATGG  
AAGAACATAAACCTAAATATTCAAGGAAACAGGAGCAACCCCTAAATAGAAATACACCCAA  
ATACATTCAATTTCTGGAATGAAAAAATAAATTAATCTTGAAGCAACAGAGGA  
AAAATGGCACATTTCTTACAGAAAAACAATAATGTAAACCACAGCAGATTTCCATCTGA  
AACCATGAAGGTTGGAAGGAAACAGATAATATTTTGAAGTACTGAAAGAACAGAACTGT  
GAACTGTAAATTCAATACCCAGCAATAATATTTCTCAGGCACTAAAGTGACATAGAAAAC  
ATTGTCTAATGAAAGATGCTAAGGTAATGTGTTGCTAACAACTTACCTTTAAAGAATA





GGAAGGATGGAGCTCCGACCCCTTTGCTTTCTGAACTCCTGCTGAGAGAGTTGGCTCCA  
 CAGCCCTGGTAGGGCTCGGGTAGCTGCTGTGGCTGAATCAGTCCTCTGTTATCACCCGCT  
 CGGTGCCATGAAGTGAAAAGCAGTCTCTGCCCTCCTCGTTCCTCCAATAAGCCCATCCT  
 AATCACCTTATCATGCTCCTTCCACACCCTGAGAAAAATGGCCTCGCAGCAGACGTTT  
 GAAGTCACCGGGACTGGAAGTCTTTCAAATGGCACCTGATTTGGCTACATGCCTGCAG  
 ACAGGTGAAAGTTAGTGCCCCCATTTACAGGTGAGGCCACTGAGGTTGAGAGAAGTCAA  
 TCAATGATGTGATCATGCTCACACATCCCAGCAGTGACCAAATATGTAACATTCATACAC

The following amino acid sequence <SEQ ID NO. 199> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 65:

VYECYIFGHCDVASHHLTSLNLSGLTCEMGALTFTCLQACSQIRCHLKDFSSPGDFKRLLRGHFFSGCGR  
 SMIRVIRMGLLBERGGQRLLFHFMPSGQRTDSATAATRALPGLWSQLSQEFQKAKGSELHPSFLADCHP  
 ASSHSPQGYVMLALKASLGRGCICHPLPCKIFEVQRALQAEHPHLLHSPVGMHSPSVGM

The following DNA sequence Seq-2345 <SEQ ID NO. 66> was identified in *H. sapiens*:

CCTGCCCCACCACCAATACTGGTGCCACGTAAGTTGTCTAGTGAAGTGAAGGAAATATT  
 CTCCTCATCAACTGCCACTCTCAAGGGCCCAAGTGTACCATTTGGAGGCTTAGGTATTGA  
 TCTGCCCCACCGTCATCACTGGCACCCTATGCACACCTTCAGGGACCTAAGGACAGGCC  
 ACTCTGCCTGCCACTGTCTACTGTGACGCAAGGACTGGCCTGCCTAGTGTCTCCATCC  
 ACAGCAAAGCATTGCCACAGCCCCTAGTTGTTAAGCCACTGAGGAGCTCACAGACACCAC  
 TCACACTGTTTACAGCAGGAGAAATCCTATGGGGCCTATAATACTGTGCCACCTTGGAT  
 CAAAACCAAAGTACTCTATGCAACTAACACTACAGCTATATCTACAGGAAAAAGCCTCTC  
 CCTACAAAAGCCAATCCAAAAACCTAGGAGAAGCAACTGTCACACCAAATACACAGATAC  
 CAACTTAAGAACATAAGAAACATGAGAAAACAAGGAAACATGGCATTCTTCTAAAGGAGCA  
 CAATAACTC

The following amino acid sequence <SEQ ID NO. 200> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 66:

LPPPPILVPTVVTEEIFSSSTATLKGPSVPFGLGIDLPHRSSLPMTFRDLRTGPLCLPLSLLVKDW  
 ACLHPQQSIATAPSCATEELDTTHTVYSRRNPMGPPIILCPPWIKTKVLYATNTTAISTGKSLSLQKPIQK  
 PRRSNCHTKYTDNLRTETENKETWHFLKEHNN

The following DNA sequence Seq-2402 <SEQ ID NO. 67> was identified in *H. sapiens*:

AGCTGGGATTTCTGCTAACTGATGTCCAGTCGGTATTTGGATATCTCCAATGACATGAAA  
 CTCACTACTGCTCAGCAACCATAGGAAGACACTGGCCAGCCCATCCACTCATGCGGTGCT  
 GGAACCCCTTTTTTATTTTAAAAATATTAAATTGACAAAAATTGCGTCTGTTCAAGGTGCG  
 ATGTGATGCTTCGATCTAGATATATACAGGTATATTGATTACCACAGTCAAATTAATAA  
 CAAATCTATCACCACCCATGATTACCATCATGTTGAGGGGATGAGGCAGTGAAGACACTA  
 AAGATCTGCTGTCTTATCAAATTTCAAGTCAACAATACAGTATTATTAACACAGTCACCA  
 TGCTGTGCATTAGGTCCCCAGAACATGTAAGTGAAGGTTGTATCTTTTGACCAACATCT  
 CCCAGCTCTGCATGAGTGGATGGTCAGCATTTCCTCAAACCCACTCTGAAGACTTTGCCT  
 GGTGGCTACATCAATATCTCCTGAGAAAGTACAAAAGTCCAGGCCAGTCACAGAAATT  
 CTGATGCATA

The following amino acid sequence <SEQ ID NO. 201> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 67:

LGFLLTDVQSVFGYLQHEHYCSATIGRHWPAPHPLMRCWNPFFILKYLIDKNCVCSRCDVMLRSRYIQVYL  
 PQSNLTNLSPPMITIMLRGGSEDTKDLSYQISSQQYSIINTVTMLCIRSPHEHVTEGLYLLTNISPALHEW  
 MVSIFQTHSEDFAWLATSISPEKVQKSRPSHRNSDA

The following DNA sequence Seq-2403 SEQ ID NO. 68> was identified in *H. sapiens*:

CAAAATATACATGCATGTACATACTATGAAATATGTATTATGTAATTTTGTGATTCTAT  
 GTATAAGTTAAATGCTTTTATATTTGCATTTTAAATTGATACTGCACACATAAAAATGA  
 ATGTGAAAATTTATTGTGGTAATTTAGATTTTAAATTTTTTACATAAAAGGACATAGAA  
 TAGCAAAGGAAAAACAAAACAACTGAAAGACGTAACAAGTTGAAAAATAGATCAC  
 AGATAAAGGAAACATTTTATACTTTGATACACTTAATAGAACCTTTTGCTTATATTTGA  
 ACTAGAGCCCCACACTTTCATTTGCCTAGACCTTACAAATATATAATCAACCTGGGA  
 CACTGAATTAAGACAAAAGCCATATTTACAAAATGGGCACCATAGCCCAAGCTATTGC  
 TTTGAAGCTACATTAGTTCCTGTTTCCAGCTGTGAGCCTGAATCCATTTAGGAAGTGA  
 GACTGGCCAGGGTTTCTGTGTAGAGTTTGGCATTTTTATTCTCTAGGACCTGCAAGAGT  
 CTACAGTAATTGTAGACTCAAAATGTGAGAGATTGCTGCTTGTATTATATAATGCCCC  
 ATACT

The following amino acid sequence <SEQ ID NO. 202> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 68:

YGALYKYKQQLTFLSLQLLTLAGSRIKMPNSTQKPWPVSLPKMEFRLTAGNRNCSFKAIAMVPIFVNI  
GFCLNSVSRVDYIIICKVKMKVWGSSSKYKQKVLLSVSKYKMFPLSVIYFSTCYVFQVCFVPLLFYVLL  
CKRIKNLNYHNKFSHSFLCCAUSINANIKAFNLYIESQKLHNTYFIVCTCMYIL

The following DNA sequence Seq-2404 <SEQ ID NO. 69> was identified in *H. sapiens*:

TATTTTCTATCTACCACATGGAATCAGAACTGTCTTGAGATTTATGCATCTGAACAAT  
 AATATTTAGAACATCATCTCGTCTTTGACACCCTTTGTTCAACACAAAATGGCTATTCA  
 AACTACTCTGGAACCTGTCTTGTCAACCAATGCAGGAATCTTAGTTAATGTATTCATA  
 AACACACGCAGGTTTCCCTTAAGCACAGACTCCATGTAAGACAAGTTTCATACTTTTCA  
 TTGTGAAAGATGCAGGTACTATTGGATGGATCTGAAGAGTTGGCAAAATGACAGGAAGAT  
 CAGGCAGGCTGCCTGTTTAACTTTATGAAATTTTTCATGTTTTATTATCTATCTACTC  
 AGATAAAATTAGGTGGGACACATTTTAAATGCTTCCAATAAATAAGAAAAATGTGCCTGC  
 AGCATGAAAAATCCTTTGACTGCCTTGTGTTATTTGCAACAGATGAATCTAATTTGTATT  
 CAGACATCAGTGCTATAACTAAGTAGAGAAATAAATGGATGTCTATGATCTCTCTCAA  
 TTATTTAGTAAGGATGAAGTGTCAATTGGCTAAAAGTAATAACACCATGGCTGTACTTAG  
 TGTTACACCTATTAGGTAGAAATATACACATACACGCATATATACAACAGATTAATAA  
 CACCAGAAG

The following amino acid sequence <SEQ ID NO. 203> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 69:

SGVINLLYICVYVCIFLPNRCNTKYSHGVITFSQLTLHPYIIERSTSLFLLVIALMSEYKLDSSVANNT  
 RQSKDFSCCRHIFLIYWKHKCVPPNFIVDRNMKNFIKLKTGSLPDLPLVILPTLQIHPVPSFTMKKYETC  
 LTWSLCLRETCLWNTLTKIPALVDKTFQSSLSNHFVLNVKSKTRCSKYCYSDAISKTVLIPCGREN

The following DNA sequence Seq-2405 <SEQ ID NO. 70> was identified in *H. sapiens*:

TCCTGAAGTCAGATAGTAGGAGTCTTCTAAATTTGTTCTCTTTCAGAAGTATTTTGGCTT

TTTATTCTTATGAATTTTCGTGTGAATTTAGAAACAGCTTGTGGATTTTAAAAGGAAAT  
 GTCTGCTTGGATTTTGAATGGAATTGCGTTGCATCCAGATCACTTTGAGGAAATTTGTATC  
 TTAATTCTATTGAATTTTCCAACAATAGACATGATGTAGCTCTCTGTTCACTCTCTCTT  
 GATTTTTTAAATAGACATTTACAGTTTTTGGCACAGAATCTGTATATGTTTGTAGATT  
 TATAGCTAAGCATTTTATGTTTTTGTATGCTGTTTTAAATTTTAAATTTCCAACCTGGTCAT  
 TGCTGCCATACAGAAATAAAACAGAAATACAGAAATACAGGGTACAAAATAAACTTGACC  
 TTGTTTCTTTCACTCTAGATAGTATTGCTTATTAGTTCTACTAAGTTTTTGGTAAGTTCT  
 TTGAGATTTTCTCCACAAGCAATCATGCTAACTAAAAATAAAACAATTTGTTTT

The following amino acid sequence <SEQ ID NO. 204> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 70:

NKIVFIFSHDCLWRKISKNLPKTNAILSRVKETRSSLFCTLYFCISVLFYGSNDQLEIKILKQHQKHKML  
 SYKSNKTYTDSVPKTVNVYLKNQRRAEQRATSCLLLENSIELRYKFPQSDLDATQFHSNPSRHFLKSTSC  
 FIHTKIHKNNKAKILLKENKFRRLLSDFR

The following DNA sequence Seq-2406 <SEQ ID NO. 71> was identified in *H. sapiens*:

AAAAATAAAAGTTATGGATCACAGCAGATCATAATAGAGAATAGTCCATCTCTCTCTAGAA  
 AATTTTTTAAAAATAAATCTTAGAACTGCATGGGAAATACTGTAAAAACAAAGGTTATTG  
 TCCTCAGCTATGAATTAGAATAAATTTGGCACTAGATTATGGGGTATTTCCACAGGAAAG  
 TACCTTACTGATTTTCCCTCTATCCTTCTTGATACATTATGGTTGAACCCACTGTTATGC  
 AACACCTGCTTACTTTGGCCTTAAGGGTCATAGTGACAAAAGAGAAACCTTTAAAGAAGT  
 CATAGTAAATGTTAGGGAAAGGGATTTTCAATGCATGGATATATTTGGCAAGGTAAACAA  
 AAAGTTGCCTGATAGCAAGGGAGGAGGCAGGCCACTGTGAATAGCAACTTATACTAGTCA  
 ATATTGAAAAGTAAAAGCAGTTGAATGGTTTCAAAGTATATAAGAATACAACTGATTGC  
 TTATAAAATGTTTTTTAAGTAGAGACTGCACCTTAAATGTGAGATGAGGCGGATCTATACA  
 TTAATTTTATATACGCAAATGATCCTACTTACATTCTTGAAAATAATTTGACTCTTTAGG  
 TGAACCAACTGAAATCTCATTTCACACTGTTGATTGCTAGTAAATAATTCTCTTTAGTA  
 TGAGAAAATCAAAGAAGTTTGAAGTGAACAAATTCTAAATTACTAGAATATGATTTAAA  
 TGGCTAGGAGAATATTATAAGGGGTATAAACAGAAATATTAATCCAAATATTTAAGATGC  
 TAATCTGGGTAAAAGCTATTTTGTAGATGACATGAATTTTCAAATACTAAAATTTTAA  
 AAATAATCATTCCACAACTTATTAAAGCTGTGTGAATGTATGTAAATACTAAGTAAT  
 ATGTTATTCATTTTAGGAACCTTTATGTATGTTTTCATACTAGTATTAGAAAATAATTCT  
 GAAAGGAAGATGAAAATGAAAATATTCATTTAGGTTAAAC

The following amino acid sequence <SEQ ID NO. 205> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 71:

VPKIFSFSFQNYFLILVKHTSSNITYYLVFTYITHSLNKFVEMIILKILVFKFMSSQKLLPRISILNIW  
 INILFYTPYNILLAIIFFRICSTSNFFDFLILKRIIYANQCKDFSWFTRVKLFSRMVGSFAYIKLMYRS  
 ASSHIKVQSLLKKHFISNQFVFLYTLKPFNCFYFSILTSISCYSQWPASSLAIRQLFVYLAKYIHALKIPF  
 PNIYYDFKGFSEVMTLKAKVSRCCITVGSTIMYQEGRENQGTFLWEYPIICQIYNSLRITTFVFTVFP  
 MQFLRFIFKNFLGEMDYSLLSAVIHNFYF

The following DNA sequence Seq-2407 <SEQ ID NO. 72> was identified in *H. sapiens*:

ATGATATTCCTATTGGATGGTGCTAATCTGGTGCAGGGTTTCTTAACCTCAGGACTACTG  
 GCATTTTGGGTGAGGTCACTCTTTATTGTGTAGGGCTGTTCTGTGGATTGTAGAATGGTA  
 AGCAGCCTCCCTGGCCTCTATCCACTGGATGCCAGTTATACCCGCTCCAGTTGTGACCAT  
 CAGAAATATCTCCAGATAAAATACCAAATGTCCTTGGGGGAGAAATCGCCCCAGTTGG

GAACCGCTAGTCTGGAGAACTCCAAGATTTAAAGGTTGTAGAAGAGAAAGAGCTGCCAG  
 AGAAGACTGAAAGGGCAGTGGAGGAGAGTGGGGTGTGTGGGGGGGTGTGGGCAGGAGC  
 CAAAAGAGTGTTCAGGACTTGGTCATGATCCTTTTAAATGCCAGTCAGATCATGTCA  
 CTTCTGCTCAAAACCATCCACACGCTTCACATCCCATTGAAATAAAATGCCAACTGCT  
 TACCATGCCCTATACACAGAACAACCTGTAATAACCTGGGCACCTTTGAGAGTGAAAGGAG  
 GCAATACTAATAATCATGCCAGGGCAGTTCAGGGCACACTGGAGGTACCATCTCCTAAGC  
 TCAGGCCCCCTGCCATCTCTCCAGCTTCATCCCCAACCACTTTCTGCCTTGTCCACTCAC  
 CCACGACAGCCTTCTTGCCATTTGTATTGGGCCATTCTCACATTGCAGGGGCCAGAGCTT  
 AGGATGACAAACATATAGCAACACATATAATGTAATGTCAGTGATATTAATAGATGCTGT  
 GAAATAAGATAAAGTGAGGTGGAGACATAGGGTGAAGTGGGGATTGGTGGCTATTTTACT  
 TAGGGGTGAGGAGATCGTCTCTGAGGATGAATCACTTATGCAGAGACCCGAATGGAGAGA  
 GGGAACTCTAAGAAGATCTGGGAAGAGGATTCCAGGCAGAAGGAACAGCAAGTGGAAAGC  
 CCTGAGGTAGGAACAAGCATGGAATATCAATAGAATGGT

The following amino acid sequence <SEQ ID NO. 206> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 72:

PFYYSMLVPTSGSLSTCCSFCLESSSPDLLRFPLSIRVSAVIHPQRRSPDPVKPPIQSPYVSTSLYLISQH  
 LLISLTLHYMCCYMFVILSSGPCNVRMAQYKQWQEGCRGVDKAESGWGSRDQGPPELRRWYLQCALNCPGM  
 IISIASFHSQRCPGYYSVYRAWAVGILFQMGCEACGWFGAGSDMILAFKDHQVLETLFWLLPTPPHTP  
 TLLHCPFSLWQLFLFYNLILEFLQTSQSGLAISPPRDIWYFIWRYFWSQLERVLASSGRPRGRLITILQS  
 TEQPYTIKNDLTQNASSPEVKKPCTRLAPSNRI

The following DNA sequence Seq-2408 <SEQ ID NO. 73> was identified in *H. sapiens*:

TTCCTATTGGATGGTGCTAATCTGGTGCAGGGTTTCTTAACCTCAGGACTACTGGCATT  
 TGGGTCAGGTCATTCTTTATTGTGTAGGGCTGTTCTGTGGATTGTAGAATGGTAAGCAGC  
 CTCCTTGGCCTCTATCCACTGGATGCCAGTTATACCCGCTCCAGTTGTGACCATCAGAAA  
 TATCTCCAGATAAAATACCAATGTCCCTTGGGGGAGAAATCGCCCCAGTTGGGAACCG  
 CTAGTCTGGAGAACTCCAAGATTTAAAGGTTGTAGAAGAGAAAGAGCTGCCAGAGAAGA  
 CTGAAAGGGCAGTGGAGGAGAGTGGGGTGTGTGTGGGGGGGTGTGGGCAGGAGCCAAAAG  
 AGTGTTCAGGACTTGGTCATGATCCTTTTAAATGCCAGTCAGATCATGTCACTTCCT  
 GCTCAAAACCATCCACACGCTTCACATCCCATTGAAATAAAATGCCAACTGCTTACCAT  
 GCCCTATACACAGAACAACCTGTAATAACCTGGGCACCTTTGAGAGTGAAAGGAGGCAATA  
 CTAATAATCATGCCAGGGCAGTTCAGGGCACACTGGAGGTACCATCTCCTAAGCTCAGGC  
 CCCTGCCCCTCTCTCAGCTTCATCCCCAACCACTTTCTGCCTTGTCCACTCACCCACGA  
 CAGCCTTCTTGCCATTTGTATTGGGCCATTCTCACATTGCAGGGGCCAGAGCTTAGGATG  
 ACAACATATAGCAACACATATAATGTAATGTCAGTGATATTAATAGATGCTGTGAAATA  
 AGATAAAGTGAGGTGGAGACATAGGGTGAAGTGGGGATTGGTGGCTATTTTACTTAGGGG  
 TCAGGAGATCGTCTCTGAGGATGAATCACTTATGCAGAGACCCGAATGGAGAGAGGGGAAT  
 CTAAGAAGATCTGGGAAGAGGATTCCAGGCAGAAGGAACAGCAAGTGGAAAGCCCTGAG  
 GTAGGAACAAGCATGGAATATCAATAGAATGGTGATATGG

The following amino acid sequence <SEQ ID NO. 207> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 73:

ISPFYYSMLVPTSGSLSTCCSFCLESSSPDLLRFPLSIRVSAVIHPQRRSPDPVKPPIQSPYVSTSLYLIS  
 QHLLISLTLHYMCCYMFVILSSGPCNVRMAQYKQWQEGCRGVDKAESGWGSRDQGPPELRRWYLQCALNCP  
 GMIISIASFHSQRCPGYYSVYRAWAVGILFQMGCEACGWFGAGSDMILAFKDHQVLETLFWLLPTPPHT  
 HPTLLHCPFSLWQLFLFYNLILEFLQTSQSGLAISPPRDIWYFIWRYFWSQLERVLASSGRPRGRLITIL  
 QSTEQPYTIKNDLTQNASSPEVKKPCTRLAPSNRI

The following DNA sequence Seq-2409 <SEQ ID NO. 74> was identified in *H. sapiens*:

AAGCTTACCCTGGCTTACACTCTTATCCAATGCCATTTACCTTGTGTGATACATAAT  
ATCTTGTATGAATCCTATTTTCTGTGTTGTGTACCTTTCTTTGAAGAATATGACCTG  
TCTCAATAATTCTTTTATGTTTTCTCTTAGTCCTTTTAACATCAGCAGGGCATTGTGA  
GTGGTGACAGGAGAAACATAAACATATACCTCTTTTCTATTGCTTTTCTGCTATTTACAA  
TAATTCGTATGACTCTGAAACAAAAGAACAATTACCTGACAAATTTCTTCTGAGTCCTA  
TATCTGGCTTTTATATCCAATCTCCTTTTATCATGCTATTACCTCTCTTTTCTTCTGTC  
TTTGAGGATGGGAAAATTCATCAACACCCTAAATACCAGCCAGAGAGGAAAAAAGAGTCT  
GGATGGAGGCAGGACTCCTTTCAAAGCTGAATCTCAAGCACTGATCACGGAGCAGCAGCA  
AAGAGACACTCAAAAAGAGTGGAGAGAGGAAAAACTAGCTGATCTCTAAGGTGTCTTCCA  
TTCAAATCACTATAATTATAAGAATGTGATTACTGGAGGAAGAACAAGGGCAGGGGCAT  
TTCTGCAACATGACGCAAAAAAATATTGACCTTAAATTTGATACATATGAACCTTTCTAAA  
TGTAGAGAGAAGCTACCTCCTTGCTGCACTTGTATGTGTGCCATTCAATTCATTTTAATA  
AAAGTTTGTAAACATGAATGAATGCAGGGGACAGACCCTCTTATGAGAATGCAGCAT  
AGTTTCAGAGAAAGTCTATTTACCAAAAACCTGAATACATGTTTATACTGAAATTTTAATTT  
TTTCTATTTTATTTTAAATTTGTGATAAAATATAAATAACATAAAATTTACCATCTTAATC  
ATTTTAAAGTATACAGTTCAATAGTATTAAGTCCATTGCGATTATTGTGCAACCAATTC  
CAGAAGCTCTTTTATCTTGCAAAAATGAACTCTATACCC

The following amino acid sequence <SEQ ID NO. 208> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 74:

KLTLAAYTLIQCHLPCVIHNILYESYFLCVCVPFFEEYDLSQFFCFSLSPFNISRAFVVVTGETTYTSFLL  
LFCYLQFCMTLQKNNYLTISFVLYSGFHIQSPFIMLLPLFSSVFEDGKIHQHPKYQPERKKESGWRQDSF  
QSISSDHDGAAAKRHSKRVERGKTSSLRCLPFKFTIIIRMLLEEEQGGHFCNMTQKNIDLKFDTYELSKC  
REKLPPCCTCMCAIHFILIKVCKHEMQGTDHLMFMQHSSEKVYLPKTEYMFILKFFFLFLFLIVIKYKHK  
FTILIIIFKYTVQYVHSHYCATNFQNSFYLAQMKLYT

The following DNA sequence Seq-2410 <SEQ ID NO. 75> was identified in *H. sapiens*:

ACCACAAAGGCTAGAGGCATGGATTATTGGAACTCTCTTCTGAAAAATTTTTACTAAT  
TTGGGAGATTACAGTCAGAATCAATGGGTGATGGTTTATAGAGTGATACCAACCTTGTC  
CAGTCCTGCTCATCATTTCATCAACAAAATGAATAAGATGAAGAGAGTATGCTTATG  
ACATCAGTGAATAGTACAGATCTCAGACTGCTGAAGAATGTACAAGATGACTTAGCCTGG  
ATCCAAAAAGCCAAGCTGGAGAGGTAGGGTGGTTCCAACAAGACAAAATGTAAAAACGAA  
GACCAATACTTAAGACCAAAAAGTCAAGCCAAACAAAACATGCTGATGTGGCTAAACAGC  
AAGTTGTGCTAAAAAATAAGACTCAAGAAGTCAAAGGTCAGTTTATATGAATCCAAAA  
GCCAATGCAATTTTAATTTGCTTTAATAAATATGTATTATCTGGAAAAAACACATACTA  
CAGTGAGTTTCTGTGGAATGAAATACTAAAGCATGTTTCTTGGAGAAAGAGTTTCCAT  
GACCAAAATAAGTTGGGGGATACTCCAAGTTGATATAAACAGGTTTATTTTCTACAGGAAT  
ACTCAAAGTCGATATGGTGACTATTGCTTCTCAAAGTTATTTGAACATGGAACTTCTT  
TTTGTAGTACCTCTTGAGGCTGGTGTTAAAGAGAACACTCTTGAGAAAACACTGAACAAG  
GGCTGTCTCAGGAGGCAGTTCTCTGTAAGTGGGACTCTTTTTAAAAACAGAAGAGATCCA  
AACATCAGATGAGTGTGGTCTAAATGACCATAAGGTTTCCTCCTACCCCTCGAAGTCTGT  
AATACTTGGTTATCCAGACCTAACAAACAATCCTAATTCCTCATGACACCTGGACCAGAG  
TTTCTGATGAGAGAACTCTAGAGAAATACTAGTAGCAGAGTAATGATTTAAAAA  
AAAACCTTTTCTCCAATGAGTGATGCTTCAAAGGGCTG

The following amino acid sequence <SEQ ID NO. 209> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 75:

QPFMSHSLEEKFFFEELNHYSATSSISLEFLSSETLVQVSWGIRIVCVWITKYYRLRGEETLWSFRPTLICLD  
 LFCFKESHLQRTASDSPCSVFSQECSLHQPEVLQKEVFHVQITLRSNSHHIDFEYSCRKTCLYQLGVSPN  
 LFGHGNSFSKKTCSISFHRKLTVCVVFQIIHIYSKLLHLWLFGBFINPLTSVLFSTTCCLATSA CFVWL  
 DFLVLSIGLRFYILSCWNHPTSPA WLFGRSLSHLVHSSAVDLYYSLMSAYSLHLYSFCLEMMSRTGQGWYH  
 SINHHPLILT VNL PNKIFQKRVSNNPCLEPLW

The following DNA sequence Seq-2411 <SEQ ID NO. 76> was identified in  
*H. sapiens*:

CTCCAGGATGCGCCCTTCCCGGCACAGCCCACTGCCATATCTTGCTGGAACCTGGGTCA  
 TCGTCCATCGTCTATCACAGGCTCCCGCAGCCTTCGTGGATGCCATCTATGTCCGTGGGT  
 CTCACCCGTCTCGCCACCAGCTTCCACTACGACGCTGGACAGTACACAGGGAGCAGACGG  
 GGATTCCAGGAGGAAGCCACTGCAAATAGGGCCTGCAGCTGCCCTCTCTCCTTCTGAAAT  
 CCTAGCATAGTCCAGGACACAGCACCTCCCTGGCTGAGCAGCTGAACTGCCAAGCTCAAC  
 TCCCTGATTGAGCAGATATTCTGCAGAAATAGAAAAGGATGGAGGGAAGGCTTCTTCCCA  
 CACAATGAACATCAAACCCACCCAAGGGGCAGTGGCTGGGGCCTCCCTTCCCAAACAGCT  
 GGCTCAAACATGCACAAATTTTCCCAAAGTGGGCTGGGAGCAGGGCAGCTGGCTTCCA  
 CTTTCATATTACTGATGCATCCAGACATACTTCCATAGTGTTAAAAATTTTGGATGTA  
 TGTCAAATGCTCTTAAGAGTGCATCTTAGGCATGTGGTAAATAAATATGATGTAATCCT  
 CCCGCTCTCAAGGGTGCTGCTGCCCTCTCCCTCCCTCCCTCACTGGTCTGGGCAAGCCC  
 TTGACCTCCACGATCTCTCTGCGCCTCTCGTGACGCCACACAAGGGGCTGTGCCAAAG  
 GGAAAGGTAGAAAAGAAAAGAGGATGTGCTGTGTGCTGTATCATCCCTGTGCCAGAGACA  
 GGGCACAGGGTGGTGGCCTTGCAACCCGGCGCATCCCCACATGGGGAAGCTGGGGTCA  
 CCCTGCACCACAGGCATCCCATCAGCCTCTGTGACACTGACAATGATTCTCGTGAATGGA  
 CAGGCTGAATGGTCTCAGCCCTCTCTTCTATGCTGGCTGAACTCTGAGGCGGGAACAG  
 GACAGACAGTGGCTGGAGGCCCTGGCAGGGAGGGCACCT

The following amino acid sequence <SEQ ID NO. 210> is the predicted  
 amino acid sequence derived from the DNA sequence of SEQ ID NO. 76:

RVPSLPGPATVCPVPASEFSQHRKRLRTIQPVHSRESLSVSQRLMGCLWCRVTPASPCGGCAGGARPPP  
 CALSLAQGQHTAHPLFFPLAQPLVGVTRGAERSWRSRACPGPVREGGRGQHPWRREDYIIIFIYHMP  
 KIALLRAPDIHPKIFKHYSMSGCISNMKVEASCAPSPLENFVHLSQLFGKGGPSHCPLGGFDVHCVG  
 RSLPSILFYFCRISAQSGSAWQFSCSAREVLCPLGCDFFRREGSCRPLYQLWLPPIVCSLCTVQRRSGSW  
 WRDGDPRIMASTKAGGACDRRWMTQVPARYGSGLCREGAHPG

The following DNA sequence Seq-2412 <SEQ ID NO. 77> was identified in  
*H. sapiens*:

CTGTCAGTTTGGTGCCCTCGGCTACGCAGGGCCTGTAGAAAGGTGCCCTCCCTGCCAGG  
 GCCTCCAGCCACTGTCTGCTCTTCCCGCCTCAGAGTTCAGCCAGCATAGAAAGAGAGG  
 GCTGAGGACCAATTCAGCCTGTCCATTACGAGAATCATTGTGAGTGTACAGAGGCTGAT  
 GGGATGCTGTGGTGCAGGGTGACCCAGCTTCCCCATGTGGGGGATGCGCCGGTGGTGC  
 AAGGCCACCACCTGTGCCCTGTCTGTGGCACAGGGATGATGACAGCACACAGCACATCC  
 TCTTTTCTTTCTACCTTTCCCTTTGGCACAGCCCTTGTGTGGGCGTCACGAGAGGCGC  
 AGAGAGATCGTGGAGGTCAAGGGCTTGCCACAGACCAGTGAGGGAGGGAGGGAGAGGGCA  
 GCAGCACCTTGGAGACGGGAGGATTACATCATATTTATTTACCACATGCCTAAGATCGC  
 ACTCTTAAGAGCATTGTACATACATCCAAAATTTTAAACACTATGGAAGTATGTCTGG  
 ATGCATCAGTAATATGAAAGTGAAGCCAGCTGCCCTGCTCCAGCCCACTTTGGGAAAA  
 TTTGTGATGTTTGGAGCCAGCTGTTGGGAAGGGAGGCCCAAGCCACTGCCCCCTGGG  
 TGGGTTTGATGTTTATTGTGTGGGAAGAAGCCTTCCCTCCATCTTTTCTATTTCTGCAG  
 AATATCTGCTCAATCAGGGAGTTGAGCTTGGCAGTTCAGCTGCTCAGCCAGGGAGGTGCT

GTGTCCTGGACTATGCTAGGATTTTCTGAGAGAGAGGGCAGCTGCAGGCCCTATTGCA  
 GTGGCTTCTCCTGGAATCCCCGTCTGCTCCTGTGTACTGTCCAGCGTCGTAGTGAAG  
 CTGGTGGCGAGACGGGTGAGACCCACGGACATAGATGGCATCCACGAAGGCTGGCGGAGC  
 CTGTGATAGACGATGGACGATGACCCAGGTTCCAGCAAGA

The following amino acid sequence <SEQ ID NO. 211> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 77:

CQFGALGYAGPVRRVPSLPGPPATVCPVPASEFSQHRKRLRTIQPVHSRESLSVSQRLMGCLWCRVTPAS  
 PCGGCAGGARPPPCALSLAQGHHTAHPLFLPFLAQPLVVGVTGGAERSWRSRACPGPVREGGRGQQHPW  
 RREDYIIIFIYHMPKIALLRAFDIHPKIFKHYGSMSCISNMKVEASCPAPSPLWENFVHVLSQLFGKGGPS  
 HCPLGGFDVHCVGRSLPSILFYFCRISAQSGSAWQFSCSAREVLCPLGCDFFRRREGSCRPYLQWLPPGIPV  
 CSLCTVQRRSGSWWRDGDPTMASTKAGGACDRRWMTQVTPAR

The following DNA sequence Seq-2413 SEQ ID NO. 78> was identified in *H. sapiens*:

TATATTTTCTGGATTACATGCCAGGTTACAAAAGGAGACCCACACGAAATCCCTGAACT  
 CCTGTGCCCACCCAGAGATTAACATGGAGAGGTGAGGGGCTGTTTCTCTCCATAGGCTT  
 CAGTGGCCTGGATGTCTGAGTTTTCAGAGACAGGATAAGTCCACATATTATTTTAAACA  
 AATTTCTTCAAACTCAAAAGCTTTCATATCTTACTTTCTTGGTAAGAGTCAAGTTTATTA  
 TCCACGTCCATACAAACACAGCTGGCTACACAACTGATCTAGGACAAAAGTCAGAAAC  
 ATGGGGCCATAGGATTCTGGGTAATGTGCTTTCTAACAAAACATATCATTTTACAGAA  
 AAGCAGACAAAAGTGATGAGAGTCTTCTGCCTTTAGAATTAGCTGACTTTAAAAATTAATT  
 TAACTCTGACATGTGACAAGAATTTTATACATCATTGCAAAATTAAGGCACTTTTGA  
 GTGGAAGTACTGATTACAGCATATTTTGTAGAGATAATGGACTTTATTTAAACACAT  
 TCTACCATTTTCTCCTGTGTTTTCTTTGAGTCCACAGAGGAAAGTTACTACACAAATTC  
 AGGTTATTTTATTGACGGTTATGTTATGGTGAAGCTAGATGAATAGAGTTTAAAGTTAA  
 GTTTTGTGGGTATTTCCAGGCCACTTGGCACATCAAACAGGTAAGCACTTTTCTCAAA  
 GAAAGTGTGTGTATTGATCTTCTTGTCTCTAGTATTGACAATTATATGAAATTTTAA  
 GCATCTCCTTAGAATCCAGCTTTTGGAGGCAATTTCTATTGAGTTTATGGCTA  
 ATCTCTTATGACATCTGTCATTCCAAGTATTTAACTCTCATATGTTTCTTTGGTGTGCA  
 TTTTTCATTGTTTAAAGCTCGTTTCTTAGGTGAGGTGTGTGTTCTTTCTTTAT  
 ATCACAGGGCTTTGTCCACAGGGTAGACTCAGCTCATGTT

The following amino acid sequence <SEQ ID NO. 212> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 78:

HELSPCGQSPVIKKEHTPSLTETSLNKKNAHQRNIEFKYLEQMSEISHKNLNRNWPSKSWFEFGDANFILS  
 ILEQSKINTTHFSLRKSAYLFDVPSGLEIPNKTLLFILHNNITVNKNNLNLCSNPLWTQKQEKMVEC  
 VLNKVHYLYQYAVISTSTPKCLNFAMMYKILVTCQSINFSQLILKAEDSHHFVCFVNMIVFVRKHIYP  
 ESYGPMFLTFPCRSVCVASCVMVDNKLDSYQESKIKLSCKKFVKYVDLSCLKLRHPGHSLSWRENSPPL  
 HVNLWVGTGVQGFVGLLLPGMIQKI

The following DNA sequence Seq-2414 <SEQ ID NO. 79> was identified in *H. sapiens*:

GAAAGCTGACAAAATTACATTTCTTGAGTCCAGTATCTATTCTTTAATTGTCTTCCTTTA  
 TATTTGAACTCTTAGTCAACTGTGGTCCAAAGAGCATTCAACTGAGGAGGGAGGCTCGCT  
 AATTTTCCCTCACCTAGTGACGCCCATGCTTGAGCTTCATGAAATTTAAGATAATTATTA  
 TTATATAGTTATATAATCATTTTCTGTAATCTTTTCTTCTTCTTTACTTTTATTTT  
 TAAAAGCAGAAAACAATAAAATGGCCATCAATTGCATGAACACTGCTCTAAAAGATAAC  
 AGTAAGACCGAACCTGAACTGTTGGCTACCTGGCCGTGCCATATTAATAGCTTACAAGGA  
 TCAGATATAGAAATATCAATCACAGGTTGTGTAGAGGTGTCATGTACAGAGCACAAACAT

TGTATATTAAAGGATGTTGAGCTTTTATAATTATTGCTATGGTTTATACAGTGTAAATA  
 AGCCCATGATAAATAGGAGCTCATATTTTATCTTAATGAAGTGCTATTTTATATTACTTA  
 TTGATTTATGTTTTTCCCCCAAGAAAGTTTAACTTCTGAGACTTAGAGACTCATTTAA  
 ATGCTTTGACCCCATACCTCTTTGTCAGGGTGCAGGAGGATGTGTATGATCTTAACCTT  
 TACAGCAAATCTCTTCTTTTGGATGGGGTATTGCAATTTTCTTTAGAGGATCACACTTA  
 GTCCAGTTCAATGTAGTTTAGAAGGGCTGACTTCATCTCTGGTTCCATGGGTGGACGCT  
 TGATCCACTCTGGTTAAGCAAATACTGCATCAGTGTAACCTCATTTGTGAATGGGTACAT  
 GATCCAAGCTGGACCAATAAGAGCCCTACCTAGAGTTTTGCTTGAATTGTTAGGATAAAG  
 GGAAATCTTTCTGAAGCACCAAGGTTATTTCTGGAGAAATCATGACCAAGAGTGAAG  
 CCAATGCATGAAAAACAAAGCCGTGAGTAAAAAAAAG

The following amino acid sequence <SEQ ID NO. 213> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 79:

KADKITFLESSIYSLIVFLYITLSQLWSKEHSTEEGSLIFPHLVTPMLELHEIDNYYYIVISFHVLSFSS  
 LLLFFKSRKQNGHQLHEHCSKKITVRPNLNCWLPGRAILIAYKDQIKYQSVVRCPCTEHNIVYKDVELL  
 LLLWFTYVAHDKELIFYLNEVLFIYITYFMFFPQESFNLLRLRDSFKCFDPHTLFAGCRRMCMILTFTANLF  
 FWMGYCNFLEDHTSSSMFRGLHLWFHGWTLDPWLKILHQCNFVNGYMIQAGPIRALPRVLELLGR  
 EILSSTKVIFWRNHDQESQCMENKSREKKK

The following DNA sequence Seq-2415 <SEQ ID NO. 80> was identified in *H. sapiens*:

ATGCATCATGTCTTCATTTTGTGGCCTCTAATAGATTCTTGGGATGTAAAAGAACTCATT  
 TTATATACATATGCAAAATTTAAACCTTCTATAATAAGTCTGACATCACCTGTGCTCTCT  
 CTGTGTTTGTGTTATCAGCAAGTGAATTTCTCAGTACTCCACATCACAAACCCCAATTA  
 CCACTCCATATGTTTCCCAAATTAGTAGCTAATAGCGTTTTCCAGGCGAATGTATCTAG  
 AAATACCCAGGGATTCACTGCTATACCTAAGTCAGCAATGGTTCATCTTTCTCCTTGCTG  
 TGGAGGAGAACTTGACCAGAGGAGTCCACTTCCCCTGGCCCGGAGCTTCTTGCAATGGGA  
 AACTAGCTGCTCCTGCTGCTACTTGGCTGATGATTTACCCTATAGCACATTTTATCTTTA  
 CGTAAACACACAAAGTCCTTTACGCTCTTTGTTCTGTTCCCATGCCATGACTCCTTCCT  
 GGAATACCATTCCTTTATTCTTACTCACTAAATAAGCTCTTCTACTCCTTTTCTTCGGGC  
 CCCCTTCCTCTGATTGAGTCTGAGAAACAATACTGTCTGTCTCCATCAAAGCTAATTTTC  
 TGCTCCTGTTTTCCACCATACTTTGCCATTCTAGACATCTGTTGCATATCATTTTTTC  
 TGTTACTTAACATGATCAGTCTTCATTCTCCTCCCAGACTATACTCCTCCTGG  
 GTTCAGAGCATATCTCATTCATTTCTGTGTACCTTTGCTTATCTCAGTCTGGCTTCAG  
 AGTAGATACTTCAGAGATGCTATTTAAATCAGAGTTAGGGTAGTTAGAATAGGAGAGAAT  
 GAGGACTCTATGGTGCTCAGGTGCCATGCATCCTGCAAAGAGAACATGAAAGGACATTTT  
 TTTTTCCTTCAATAATTACATGGAATCCTTCAGTGATCCCTGTGTCTGTTGGGCCTTGAG  
 TAATTACCTGCAATCTCTGTCTTTGTGAGGCTATTAATTA

The following amino acid sequence <SEQ ID NO. 214> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 80:

MHHVFILWPLIDSWDVKELILYTYANLKPSIISLTPVSSLCLCYQQVNFVLPVPHHKPQLPLHMFPKLVAN  
 SVFPGECIKYPGIHCYTVSNGSSSLLWRRTPPEESTSPGPAASCMGNLLLLLGLFTLHILSLRKHTKSFHV  
FVPVMPPLPGIPFFYSYSLNKLFSFSSGPLPLIQLRNNYCLSPSKLIFCLLFSHHTLPFTSVAYHFFCY  
 LTNASVFIHSPRLYSSWVQSISHSFLCYLCLSQCWLSQRYFRDAIRVRVVRIGENEDSMVLCHASCKE  
 NMKGHFFFLQLHGLLQSLCLLGLLPAISVFVRLLI

The following DNA sequence Seq-2416 <SEQ ID NO. 81> was identified in *H. sapiens*:



GCCAGTCAATGCCAAAGACATTCTGTTCCGGTTTGGGAATGAATAAACTTCTGATGCCCAT  
 ATGGTAACCTTATGCTTTGAGAACTCTTCTATAGCACAATAAAATCTGAGCCGTCAGAGT  
 AACTAAGTGATGGAAAATGAATAACTAAATGTATAGGGAAAGAATCCAGAAAAGAAATTT  
 GTATTTTATTTTTTCTAAGTAACTCCACAGATATGTTTGAGAAAAGTGTATGATCTAGT  
 GAATAGAATACTCAAACTCTAATATACAAGTCACAGGTATGGGCCCTAGTTACTTCACT  
 AAATGACTGGCTTTAGGCAGATAACTTGTCTGGTTCAGTTACTAAGTATGAGAAATAGA  
 AAATACATCATTAACCTTTCTATAATAGTCCACAACATTTTCAGCACACCCAATGTGACAA  
 AAAACCGTCTCAAGCCCACTTCAGTAACAAGTGAAGATTTGTGGGTTCAATTAAATGTCA  
 AGGCCAGCAGTAAGTGAGGGCTGGTCTGAGGCTGACATATTCTGAGGAGAACATGGTCT  
 TGTCTTCTCTTTCTGGGCACCTTTGTCTCTGGATGGAATCCATTCTTGGGCAGGCTGA  
 AGTCCTTCTCTCATGGTGGCAAGATGGATATGCCAGGCAACCATCCTGTCTGCAGAGAGC  
 CTGCTTAGTGAGAAGTTTGGGATTAGTTCTGACTTGATGAATTTGGGTCTCATGTTTAT  
 CCCTGGATATATCTCTTTGCTCAGGTGAATGGATATGTTGACTGCCACACCTGGGTTTC  
 TGTGACTACTCCTGGATTCAAGTATGGAGTCAGCCCCAAGTAAGGCCCATAAACAAGGGT  
 GGAGGAGAGTGGTTCCTGGAAAGAAAGTCAGGGTAAAGGCAAGGGGACAAATGCCAGATG  
 GGCAGTAAATGGCAGCTGTCCAAATTTATGCCTGAACCACTGAAAGGAATCTTCACTCT  
 CACTGTGGGTATTAACATAGGACGCGGTGATGCTTAATGG

The following amino acid sequence <SEQ ID NO. 215> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 81:

PVNAKDILFGLLEIKLLMPIWPYALRTLHKNIAVRVTKWKMNNMYRERIQRNLYFIFSKLPQICLRKLYD  
 LVNRILKTLIYKSQVWALVTSLNDWLADNLGSSSYLEIENTSLFFYNPQLFQHTQCDKKPSQAHFSNNEF  
 VGSEFKCQQQVRAGSEADIFGEHGLAFSLGTFLVWMSILGQAEVLLSWWDGYARQPSCLQRACLVRSF  
 GISSDLMLNLGMLFIPGYISFAQVNGYVDCHTWVSVTTPGFSQVSPKGPTRVEESGSWKESQKKGKGTNAR  
 WAVNGSCPNEFMPEPLKGIFFLTVGINIGRDAW

The following DNA sequence Seq-2417 <SEQ ID NO. 82> was identified in *H. sapiens*:

ACTAGCTTGGATGCACAAGGATTCAAGGATGCATAGTTAGCAAGTAGCAAAGTAGTTATC  
 AAGCCTAGGCGGGCGCTGACTCCAGAATTCAAGCCCAAGGTCACTTCTCTATACTATTTT  
 ACATTGTATTTAAGAACTACATGAACATGAATGCATGGTGTGATGCTTATAGTTTCCTGA  
 TGCTTATAGTGTCTGATCCTACTTCTGCATAAGCCATGCAAAGGTAGTGACCCAGACTG  
 TAGAAATGCGTCAGAGTGAGATATACCAACAAAATGAAACGAGTGAAAGTAGTATAATTT  
 TCCAACATGTATACACTCTCTCACACACATACAGTGAGAGGAGAACTAAAGATTAGT  
 GACAGGGGATTTATAACATTATAAAATCTGAGAGCCTGAAAACAAAGATCCAAGGCAGAG  
 CTAGAGGAACACAGGTATGGGTCAAGTCAAGTTGAGAACACAGTGATAGGGTTCA  
 GAATGGTTAAGTATAAACAGAACTAGTGTGACAGAAGTCATTCTTACATAATATTTTTTT  
 AGTTGGTACCAAGATGGAGTAGATGCAGTATGTGGTAGTAAATCACAGGTAATTAACATA  
 AATTGTTAAAAAATTGAAATATTGTGCTCATTACTGATTTGTCTCCAATATTTATCTCTGA  
 TAGTCAATAAATCAAAATATATCAAAGCTTAAATTGTGAGAATAAAACCATGTTTGTAT  
 AATTGCAGAAAAATTATTGAAAAGCAAACTTGTGAGGAATCCACGTGTTATCATTGCA  
 CAGCTCATATGAATCTGAAAAGTCACAAATAAATTAGCAACATGGAGTTAATTGGTTTTT  
 CTTTTTTTGTCTTACTGTTATTTTTCTTTACCATGCAATTTCTTTTCTGGTTTTTGT  
 TTATTATGGAACAATACACTCTTTTTCTTAATATTATGCTTCTGCATCCTTGCTTAT  
 GAGTTTCTTCTTACATGAATGCTGTCCTTCTTCTCC

The following amino acid sequence <SEQ ID NO. 216> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 82:

RKKDDSIHVRNRNSARMQKHKEKRVYCFHNKTKRKEIACGKEKQSKKRKTNLHVANLFTVFQIHMSCAMI  
 TRGFDPDKFCFSIIFLQLYKHGFYSDNLSFDIFFIDYQRILETNQAQYFNFQFSLPVILLPHTASTPSWYQL

KKYYVRMTSVTLVLFILNHSEPYHCVLNLHLTDPYLCSSSSALDLCFOALRFYNVINPLSLIFSSPLTCMC  
VESVYMLENYTTFTFRILLVYLTLTHFYSLGHYLCMAYAEVSGSHYKHQETISITPCIHVHVVLKYNVKYR  
EVTLGLNSGV SARLGLITTTLLANYASLNP CASKL

The following DNA sequence Seq-2418 <SEQ ID NO. 83> was identified in  
*H. sapiens*:

CATGGCCCCAAATTAGTTTCCACCTTATGTTCCACTAGTTTCATAGACAAACCTCTTCC  
TGCCATACTGGTCTGGTCAGTGCCTCCAGACACTGCAGTACTGCCTTGAAGTGGTTGC  
TGTCATCTTTTCTCTCTGTCTATCTAAATTCTAGCCTGTCTTTGATGGCTAAAAGCCTAAC  
ATCTCTGTGGGCCTCAGAGAAATTATCTTCTCTGCATTCTCCAGTTGGCATCTCTCAC  
TAATGGATTAATCATATTACCCTCTCCTATTGTTATGTGCTTTTATGCATATAATCTTAG  
CCCCCATAGGACCAACTGTAATCCCTTTGAGGACAGGGGTTTGATCTGTACCTATTT  
ATAGTTCCCCACGTGCCTAGAGCCTCTTGCACTGTAGGCTGGGGGAAAATATTTGCTT  
ATGCTGATGATCTGAGAAAGATAATACTGCAAACAGGAGAAGTAAAGATTCTTTGTCTT  
GTTCCATTTTGGAATGAATTAGTGGCAGGTAATCAGTTAGAGGTCAGTTCAGAAGGTTAA  
ATACGTGGACTTATCCCTGTTACAGGTCTCTTATCTTTACAAAGATTGTGTTCCGTGTA  
CTAACCTCTTTCTAAATCATTGGTGTGTTATTTACAAGAAGGACTGGGCCAAATATGTG  
AGGAAACATCAATGTATACTCATCCCTACCATTGAAAAACAAGTTTTAAGTGTGTGTAC  
CACTGATGAAGTATGAAGAATAACGTTCCATTCTCCAGAGTACTCAGGCCCTTTGCC  
TGGGACTGCTAGCTACACATGCAAAGTGAATTCTATATCAGCATTGTTGTAAAGCCCACTA  
TTCTCACCCTACCAGCTTAACGTCAACCAGTTATTTAATAGGATTCTAATTAATTTAATT  
CTCCACTGGTAGCAATTTCTGATGCACAATGTCTGTGCCTTTTACCTCTTGCATCCCTT  
CCCCAGCACTTAACCTCAGCAGGTTGCATATAGCAGGAACC

The following amino acid sequence <SEQ ID NO. 217> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 83:

WPQISFPPYVELVSTNLEFLPYWSGQCPPDTAVLPTGLLSSFLSVIILACLWLKAHLCPQRYNPLPHSSSW  
HLSLMDSYYP LLLLCAFMHII LAPPDQLSLGQGF DLVPIYSSPRASLLHTVWGWKIFAYADDLRKIILQTG  
EVKISLSCSIWNELVAGNQLEVSSEGNWTYPLQLQVS YLYKDCVPVTNLEFLNHWC CYLQEG LGQICEETSM  
YTHPYHLKNKFVCPVPLMKYEERSHSFQSTQALCLGLLATHAKILYQHFKPTILTVPALQPVIDSNFNSPL  
VAISDAQCLCLPLCIPSPALNSAGCIQE

The following DNA sequence Seq-2419 <SEQ ID NO. 84> was identified in  
*H. sapiens*:

TAACTTGTTCCAGCACAGATTCAAAAGTCTAAATTCTGAAGTCTCAACTAAATGTCATCT  
AAACCAGATGTAGGTGAGACTCAAGGTATGTTTATTCTGAGAGAAATTGCTCTCCATCTG  
TGATTCTGTGAATCAATAGGTAAAGAGCTTCCAAATGCAATGGTGGGACAGACATAGA  
ATCGACATTCCCATTCAAAAGGGAGAGTAGGAAGGAATACTACAACAACAACAAAGTA  
AACGATAAATCTTAAGGCTCCAGAATAATCTCCTTTTGATGCCCCATCTTCCAATCTTCC  
AGGCACACTTGGGCAGGCGTTGGGCCCCCAAGGCTCTGGGTGTCCAGTCCCAGCCCACA  
TGACAGCACTTACATATTAGAGCCACATGCCAGGCTGGAATGCCCTCTAGTGGCTCTAC  
TGGTCTATGGTCAGAGGGTAGGCTGTCTCTATGACTCTGCCAAGCACAGCCTTAGTGGA  
GGCTTTTGTGGTGGCCCCACCCCTATGTCAATTCTTTGCTGAGCCTCAAGACTTTCCA  
GGGCATCCTTTGAAATCTGTGTGGAGTCAGCTTTCCCTCTATGGTATTGCACTGTGTGTC  
CTGGTGGAGATGATACCTAGAGAACATTACCAACGTTTATCATCTGTGCCCTCCAGAAAG  
GTGGCCACTGGAGCCCACACCACACTTGGACCCTCTGGAGCCATGCCTGGAATGACTGAG  
CAGTCTGTGTGAGAAAGCAGGGAGCAGAGATGAGGTAGCATAGGGCAGGAAGTGTGAG  
CTCGCTGGGCATCCTGGGCCCTCTTTGACCTTGTCTGTCCCCCTAGGCCCTTGGCAGC  
CTGGGCCTGTGATGGGAGCAGCAGCCGTCTGATGTCTGAAATGCTTTTAGTGGGGGTCA  
TTCCTCCATTGCCCTTGATGAAAAGCACCTGGCTTCTGCAGTTCATGTTAATCTGATCAA

ATGGTTGCTGGGCCACATCCTTGGTATTCTCTCCCAAACA

The following amino acid sequence <SEQ ID NO. 218> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 84:

TCSSTDSKVLKSQLNVITRCRDSRYVYSEARNCSPSVILIKVKSFNAMVGQTNRHSKREKEGILQOQQQ  
SKRILRLQNLLMLPHLPIFQAHLGRRWAPKALGVVPAHMTALTYSHMPGWKCPLVALLVYGQRVGLLLL  
CQAQPWRLFVVAPPLCQFFAASRLSRASFEICVESAPFLWYCTVCPGGDDTRTLPTFIICALQKGGHWSPH  
HTWTLWSHAWNDAVLCQKAGSRDEVAGRKCAPVGILGPSFDLVLSRPWHAGPVMGAAVMMSEMLLVGVI  
PPLPKAPGFCSSMLISNGCWATSLVFSFK

The following DNA sequence Seq-2420 <SEQ ID NO. 85> was identified in *H. sapiens*:

CCACAGAAACATTCTTCAGTAGAACTTTAATATTACTGTCTTATAAAATTCTGTCAAATG  
AACAAAAGATAACCCATAATTACACCCTAATATGACTGCTTTTAACATTTTACTGTATTT  
CAGCCTTTTGTCTATGTATATAATTTTACAGAGTTGTAATCATAACCCAGTATATGATTTT  
ATCATGTTTTCCCACTTACCATTATAGGTATTTTAAATATGCTACATAGTCTTCATGGT  
TGTCAATTGTAATAGCTATGCTGTAATAGTTCCTGAATTGAAGTGCCTTTATTTACTTAG  
CTACCCTATTATCTTTAAACAATTTCTAATTTCTTTTATAATAAACATGGACATATTTT  
TGACAGGGGTGTTCTTTTTCACATCTTGACCTACTTTTCACATAGTGTACAAATACCTG  
ACCAAAGAATACAACTTTTTGTCTCTTGACGTATATTTCCAAAAGATTTTAAAAGGTG  
CATTAAATTTACTCTGCAGCTGGTGTAATGAAGACCATTTTGTCAATTGTTTCTTGAGAG  
TAGAGCTTCCAAAAGTAGGGATATGTGGCTAGGAGGAAGAAATCCAGCCTGGGGCAGGCA  
TTCTGTAAAGAACTCCAGTTCTCACTGGTACACTGGTTTTATTTTCTCTGTTTCTTGCA  
GACTGAGCAATTGATAACTCTGTGGTCTCTTTGTTTTTACCATTGTTGGAACCTCCGT  
TGTGCTTTTTTCCACATGGAGGAGAAAGAAGTCAAGAATGACCTCTTTGTGACTCA  
GCTGGCCATCACAGGTAAGTAATATGCAAGTGAGAGGCAGGAAGCTATATGTGAAGTCC  
CTATGGCTTCTGCTTTTAAATGAATTTTATCAAAAAAAAAAATGTAAACGCATCGGTCA  
ATTTGGGAATAATTTCTGAAAGAATATAAACCTATATTGAAATATTTCTCTGGCATAC  
TTAACACATATGAATGCCTCTAAGATTTTATTATAAAAGT

The following amino acid sequence <SEQ ID NO. 219> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 85:

HRNILQNFNITVLNSVKTCDNPLHPNMTAFNILLYFSLFAMYIILQSCNHTQYMLSCFPYHYRYFYCYI  
VFMVVIVNSYAVIVHIEVLYLLSYPIIPKQFLISFYNKHGHISDRGVLFHILTYFSHSVTITPKNTNFLSL  
DVYFQKIFKRCINLLCSWCKRPFCHCFLESRAKSRDMWLGGRNPAWGRHSVKNSSSHWTGFIPLCFLQT  
EQLITLWVLFVFTIVGNSVVLFTWRRKKKSRMTFFVTQLAITGKLCKEAGSYMSPYGFLLLMNFIXKKKM  
RIGQFGNNFKNIKPIFEYFLWHTHIMPLRFHYKS

The following DNA sequence Seq-2421 <SEQ ID NO. 86> was identified in *H. sapiens*:

AATAAGCAAATCTATTTTGACAGAAAGATTCATGATTGCTCCTGGCAGCAGGGGTGAGG  
AAGTTGGTGGGAAATGGGTACAGAGATTCTTTGGCGATGATGAAGACGTTGTAACAGCT  
TTTGAATTTTACAATCCAGAATTCTATTCTCTGCTAATTAGTCAAATAAAGGGCAGAAAA  
TATACATTTTAAAACACAAAGATGCAGACATTACATTCCACATACAAGAGGATGTACCCC  
AGCAAAACAGGTGATAAACCAAGAAAGAGAAAGATGGGATCCAGGAACAACAGCTTCA  
ACCCAGGATAACAACAAGGGAAGTACTCCAGTGTAAACAGCTGGGCAGCCAGAGAGACA  
GCATGTAGTCTCATTGAAGCAGAAAGACAGAGGTTCTGAGACAGAGGTCTCCAGGAAA  
AAAAAAGAACCTGACTTACTGGATAAACAAGTCTTTAGTTTAAAAACAACAAAAA

TGTATACACATATATATATAAAATCAGGTAGTATAAAGAAAAACAGAACTCCAGAGATTCTGGGTCACAGAAGGGGAAAGGGCTGTTCAAGAAAGTGAAATTGAACTAACTGAAATACAGCTATCTTTATATTGGAAGGACAGTCAGGAAGTCAACAGATAAGGCCCTAACTGCATAAAGCAGGAAACAGCAGACTAAAGACATTATTAAGAAATATGGAACACAACCAAAAGAAATAGCAAAAACAAATGAAAAGTGACTGTTTTTCATAAGTGAGGCAGGGGAAGAGAAGGGGTTATTTTTTCCCCATTATATGTCTTTAAGAACTACTTGCTAAAAATATTGGGCACATATGAATTGATAAAAGCGAAAAACTTTTTACTTCACAAGTGCAGCTTTAACATACGTTGATTACAGTGAAGTTTTTGTCTGTTAACCACTTTAGTAGGATTGTCTAAATTTAGTGATTTACAATGCCTGCAGTAGAATCAGAAGATTTACACTGAAGGGATTAT

The following amino acid sequence <SEQ ID NO. 220> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 86:

IIPSVIFFYCRHCKSLNLDKSYSGQNKNFVINVCSCTCEVKSFSLLSNSYVFNIFSKFLKTYNGEKNPFSSPASLMKNSHFSLLFLLVVFHISCLSAVSCFMQFRPYLLTSLSFQYKDSICFSFNFTFLNSPFPFCDPGISGVLEFFILPDEFIYICVYSFLLFFLKTKCLSSKSGSFFFSWRPLSQNPLSFCFNEDYMLSLWLPSCHWSSSLCCYPGLKLLFLDPLSLSWFITLFCWGTSSCMWNVMSASLCFKMYIFCPLFDLAENRILDCKIQKLLQRLHHRQKNLCTHFPPTSSPPAARSNHESFCQNRFAV

The following DNA sequence Seq-2422 <SEQ ID NO. 87> was identified in *H. sapiens*:

CCTTCTCTTTCGGGTATTTTAGTCAGCCTCTTTTTATCGCTGTTATCACAGATATCCCCAGAGACCCTTGTTATCATAATTGCTAATGTTTCACAAAAGATGACCATTAGTTTAAATTAATCTTATAGGACTTACACTCTCAATTGTTAGGCAAGGAAATTGAGCCAGGTCAAATTAAGTAAATGCCCCAAATCTCACTGTTTTCCAGTAATTTAAAGAGTGACATCCAGAAAATCTGTGACTTCTAGGAATACATTTAGAAAAACATATACCAGAGGGTTAATTGCAGCATTGTTTTTAACAGCAAAAATTGGAATAAATACACATCAATTGGATACAGATAAATAAGTATGAGATATTCAGACCAAGATCCTGTGCTGTAATTGAAGTGAATGAAGTGGCAATGTGTGCACCAAGTATCCCCAAATATAATTTACTAAAAAAGCAAAATGCTGAATGATTATGCTGTATGATAACATTATATAAAGTCTGAGAACATGAAAAGCAACTGCAACATAGATTATAGCTGCATAAATAAATAAATAGTATAATAACATTTGTAGGAATGGAATAGAGAAAACATTATGAGATCCAGAGTGCCCCAAAAAACCTGCCCCCATATTTTAAATCAACCATTTTCTCATTTAACCCCATTTTTCTCATCACTTACTATGTGACTAGATGTTCTTTGGTTTGTTAAAAAACATTTCCGATTCCTTAACATACCTAAAAATATAATAAATTATTCTCTCATTATTTTCTTCTACATAATATACAAATTACTTCAAATACGTACACAACCTTACTTTACATAAATAATCTAACACAGTGGCTTTTCTTAGGTATGCATTCTACTAAATCATATATTCCTTCTCTAATAATAAAAGATTATATGACTTATAATTATATACTACCATAGCTGGGCTATCATAGTAGCCTTTCTTTAATATAAATACTTTGATACA

The following amino acid sequence <SEQ ID NO. 221> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 87:

CIKVFILK GKATMIAQLWYIIISHIIFLLLEKGIYDFSRMHTKPLCIILCESKLCITYFEVICILCRRKENNLLYFVCGIGNVFLTKPKNISHSKGKMGLEKMGVLDLYGGRFFWGTLDLIMFFSIPFLQMFIIILLFIYAAIIYVCSCFSCSQTLYNVIIQHEFSILLSLVNIIWGYWCTHCQFIHFNYSTGFWSMNISYFIYLYPIDVYLVEIFAVKNNAAIKPSGICFSKCI PRSHRFSGCHSLKLLGKTVRILGNLNLTLWNFLAQMRVVDLIKNMVIFCETLANYDNKWSLGISVITAIKRGKLPKEK

The following DNA sequence Seq-2423 SEQ ID NO. 88> was identified in *H. sapiens*:

GGGACATTTTATGCTGGGGAACATTTTAGGCAATGGTCCCCAAGACCTTTTCGATAAGG

ATACTCCAGCGAAACAAATGAGACTGTTACAGGAGGCAGCACTGAGGCAGGGCAGGTGGC  
 ATTGGAGAACATGCACACCACCCCATGGGCACCGTGCAACACCACCCACCCCATGGAA  
 GTGGTGACAACAGTGGGGAGGGGAAGCCTGTCAAGCAGATGTACCAGGTGCTTCAAGCA  
 GTGTTGTAGGTCCCTGCTTATAGGTGCCAGGCCAACTCACCACCTTCCTTCGACTCTTG  
 GAAAGAAAATAGTGGAGGTCTTTCTAAATCATGTGAGACAATAACTCCCCAGAGGTGCC  
 ATCCTCTAGATTCCAGGGGATAAAGACGAGCACAAGAAGTACTGCTGAGCACTTTGTGTG  
 GGATGTGTGTCTAAACACGACAATCTGAAGACAGAGGTGTAGAAATTGGCAAGTTTCCTA  
 AAGCATGACAACACACACCCAAAACCTTCCATAATGATTCCCTTTTCCCTGTATTTT  
 CCTGGATGCACCATCACTATGGGAACAGGATGGTTACTCCCAATCCCTGTCAACCACC  
 GCTTATTTAATAAACGATTTCTACTTTACTGAAATTGATGCTTCGTTTTCTTCTAATTCC  
 ATTTCTATACTTTACCTCTGCTCTGAGTTACTGAATTTATAACCCCTCTTTTAAACAGA  
 AGTCTTGCAAGAACAACTACAGCAGTATCAGCAACCAACAATGCCACCAATACAGATTA  
 AAAAAACATTCTTATCTGAGGCCAGGTAACCAATTTATGCAAAATAACTCAACAGATGC  
 TGGTCAGTACTAGCTGACCCATGAATTTAAGCTCTTACTTGGAGAAATACAACCCAAAG  
 AGGAGAGAAAGGAAAAAATGAGTCTCATATTAACATACAATAAAACCTTATTAAGTAT  
 AACTCCATAAATTATGAGTGGCAATCAGATAGATAATTCA

The following amino acid sequence <SEQ ID NO. 222> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 88:

NYLSDCHSFMELSVNKVLLYVNMRLIFFLSLLFGLYFFQVRAIHGSASTDQHLLSYFAIWLPLGRECFNL  
 YWWHCWLLILLEFVLARLLFKRRVINSVLRAEVKYRMELENEASISVKKSFIKAVGDRELGVITLPIVMV  
 HPGKIQGKRESLWKSFGCVLSFCRKLNFYTSVFRLSCLDTHPTQSAQQYFLCSSLSPGIRMAPLGELLSH  
 MIKDLHYFLSKSRKVGELAWHLAGTYNTASTWHLLDRLPLPTVVTTSMGGGWCCTVPMGWCACSPMPAL  
 PQCLLQSHLFRWSILIEKVLGTICLKCS PANV

The following DNA sequence Seq-2424 <SEQ ID NO. 89> was identified in *H. sapiens*:

TATTATGTTATTGTGTAATTATTTGAATTATTGTCCCTTTCCATCAATCCCCAAACAC  
 ACACATATTAGGTGGAATCCTTAGGGGCTGAGATGATGTTTTATTACATCTGCATGCC  
 TGATGTTAAGCCAGTGTCTGGGCACGAATGCGATTTAGTGAGTGTTCCTGAACATGAAT  
 AATGAATTCACCAAGTGAAGCATGAGTGGATCTGGTGGGGGCACAAAAGCTGACTCCAG  
 GTTCCAGGAATCTGGGTGGAGAACTTCTGGGCTGGAGGGAGCAGAGGACCACTGTGTGA  
 GGTCTACGTGGTCTGGCTGGCAGGGTTAGCAAGGATGCAGAGGAGTTTCTGGGTCTGTC  
 TCAATGATAATTTAAAACAACAATAATAATTAACATTCATTTAGTTCTTACTATGTGTC  
 AGTCCCTTATTGCCTTCTATGTATTAGCCACTAATCCTCAAAATTCAGGGGTTAGATA  
 TTTTTCGGTCTATACTATACATATGAGAAAAAGGTAGAACAGGGAGGTGCAGAACTT  
 GCCCAGGATACAGCAAGTAAATGGGAACCTGGGATGGTCCCTAGGGATTCTTGTT  
 TTTTAGATTTTGTCTTTTAAATCTCTCTATAGCCCTTAGGTTATTTATTGATATTTTA  
 CTTTTATTGAAATAATTGTAGATTACAGGAAGTTACAAGAGAGAGTCCTGTGTAC  
 TCTTCAACCAGATTTCTCAATGCTTAGATTTATATACTGTAATAACAATATGAAACC  
 AGGAAACTGATATTGGTTCAATATATGTGTATACTTCTATGCCATTTATCATGTGTAGA  
 TGTAACCACCATCATGACCAAGCTGCAGAACTGTTCCATCACCACGAAGATCTGCCACCT  
 GTTGCTCCTTTAAAGTCATACCAGCCCTCTTCCCTGTCCCCACCCACTGTCACTATGCTT  
 AACCTTGGTAACCACTAATCTGTTTTCCCATCTCTATAG

The following amino acid sequence <SEQ ID NO. 223> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 89:

LCYCVIIIIIVPFPISIPQTHTYVEILRGDDVLFTSACLMLSPVLGTNAIVFLEHEIHQKHEWIWGHKRLTP  
 GSRNLGGETSGLEGAEDHCVRSTWFWLAGLARMQSRFWLLKFKTTIINIHLVLTMCQSLIAFYVFSHSS  
 KFGLDIFPVYTIHMRKRVEQGAETCPRIHSKNGNWDWSPRDSCLDFVFLISLPLRLFLIDIFTFYFEIIV

DSQEVTRERSCVLFTQISPMRLRFYITVIOYENQETDIGSIYVYTSMPFHHVMPSPSPSCRTVPSPRRSATCC  
SFKVIPALFPVPTHCHYAPLVTTNLFSLY

The following DNA sequence Seq-2425 <SEQ ID NO. 90> was identified in  
*H. sapiens*:

ATGCATACACCAGAGCCGACCCGACAGCTCTGCAACCCAGGCCAGCTGCACGGTCAGTT  
TGGAAGTCTACACAAGCATCTAGAGGACCTGGACACAAACAGGGCTAATTCAGGTGCCCA  
ATTCATGTCCCAACTCTGTCTGTGAGGCGACTAAGGCAGGGCTCTGGGAATCCAGGGAC  
AGGTGGAGTAACTCGTACACAGTCAGTGTGGGAGTCTTAGCAGGTGACTGGGTCTGCCC  
GGACTCGTGTGGGATGGAGGGCTGGGTAAACTCATTGCTGCAATAAAAGGGACAGAATCT  
CAGTGCAAAAGAGACTAGAAAAATGTTAGGTTCCAGAGAGAGGCTGGAATTCAGAGGG  
GAAGATGGAAGCCATTGATATAGTAGTGGTGAAGATGGAAGGTGGCCCTGCCGTGAG  
GAAGACACCTGAGCTATGAAGAGTGGAGTATAAGCTTGAACCATAGTGCACATACCCA  
GAGTTCATGTCCAACATATCTCAAATCTTTGCAAGTCTGTGTGGATCCTTAAAACTG  
GGGAGGGCAGAGCCAGCAGTGGGCGAGTGGCCCCACCTGGAGGAATGGGATTATAGAGT  
CCAGGAGTGAAGCAGCGCCCTACAGTTTGTCTCATCCTTCCATTTCCACACTTCCAGT  
TTCCTTTCAACCACTTCAGAAAAAAGTCCAGAAAGTCTAATGTTGCCAAGTTTA  
GAAACCAAGTCGTATTAGTGTGAGTGAATCAACGTTGATTACAGTCTGGTCTTTTCA  
AGTTTCTTTGATATCTTCAAAGCCCAATCATCTGTTCATCTAGGACATTAAGAAAA  
TACACCCAAAGATAGTCTTTCAAGTACATTGCCACCGTAGCTAGATGATTATTATCTG  
ACTATTAATTACTATTATGATTACTGTTGCCATGGTTTTATGTTTTTCTGTGTGCCAT  
CCAATCCACATCCAGCCACCACAGCCACTGCTGGGTTTT

The following amino acid sequence <SEQ ID NO. 224> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 90:

KPSSGCGGWMWDMGTQKNIKTMTVIIIVINSQDNNHLATVAMYLKDYSLGVFFLMSMEQDDWAFEDIKE  
TKGPDCNQRFHSHRPGFTWQHTFWTFFFFSGKETGSVENGRMRTNCRALPHSWTLSHSSRWGPPAHCWLCF  
PQFLRIHTDFAKILRYVGHELWVCAHLVPSLYSTLHSSGVFLTAGATFHLHHYIKWASIFPSEFQPLSGN  
LTFFLVSFALRFCPFYCSNEFTQPSIPHESQDPVTCDSHTDCVRVTPPVPGFPEPCLSRLTGQSWDMNWA  
PELALFVSRSSRCLRLPNPCSWAWVAESAGRLWCMH

The following DNA sequence Seq-2426 <SEQ ID NO. 91> was identified in  
*H. sapiens*:

TATTATGTTATTGTGTAATTATTTGAATTATTGTCCCTTTCCATCAATCCCCAAACAC  
ACACATATTAGGTGAAATCCTTAGGGGCTGAGATGATGTTTTATTACATCTGCATGCC  
TGATGTTAAGCCAGTGCTGGGCACGAATGCGATTTAGTGAGTGTCTTGAACATGAAT  
AATGAATTCACCAGTGAAAGCATGAGTGGATCTGGTGGGGGCACAAAAGGCTGACTCCAG  
GTTCCAGGAATCTGGGTGGAGAACTTCTGGGCTGGAGGGAGCAGAGGACCACTGTGTTA  
GGTCTACGTGGTTCTGGCTGGCAGGGTTAGCAAGGATGCAGAGGAGTTTCTGGGTCTTGC  
TCAAATGATAATTTAAACAACAATAATAATTAACATTCATTTAGTTCTTACTATGTGTC  
AGTCCCTTATTGCCTTCTATGTATTAGCCACTAATCCTCAAAATCTAGGGGTTAGATA  
TTTTTCCGGTCTATACTATACATATGAGAAAAAGGTTAGAACAGGGAGGTGCAGAACTT  
GCCCCAGGATACACAGCAAGTAAATGGGAAGTGGGATTGGTCACCTAGGGATTCTTGTT  
TTTTAGATTTTGTTTTTTAATCTCTATAGCCCCCTTAGGTTATTTATTGATATTTTA  
CTTTTATTTTGAATAATTGTAGATTCACAGGAAGTTACAAGAGAGAGGTCTGTGTAC  
TCTTCAACCCAGATTTCTCAATGCTTAGATTTATATACTGTAATACAATATGAAAACC  
AGGAAACTGATATTGGTCAATATATGTGTATACTTCTATGCCATTTTCATCATGTGTAGA  
TGTAACCAACATCATGACCAAGCTGCAGAACTGTTCCATCACCACGAAGATCTGCCACCT  
GTTGCTCCTTTAAAGTCATACCAGCCCTCTTCCCTGTCCCCACCACTGTCACTATGCTT  
AACCCTTGGTAACCACTAATCTGTTTCCCATCTCTATAG

The following amino acid sequence <SEQ ID NO. 225> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 91:

LCYCVIIIIIVPPPSIPQTHTYVEILRGDDVLFTSACLMLSPVLGTNAIVFLEHEIHQKHEWIWWGHKRLTP  
GSRNLGGETSGLEGAEDHCVRSTWFWLAGLARMQSRFWLLKFKTTIIINIHLVLTMCQSLIAFYVFSHSS  
KFGLDIFPVYTIHMRKRVEQGGAECPRIHSKNGNWDWSPRDSCLDFVFLISLPLRLFIDIFTFYFEIIV  
DSQEVTRERSCVLFTQISPMRLRFYITVIQYENQETDIGSIYVYTSMPFHHVMPSPSCRTVPSPRRSATCC  
SFKVIPALFPVPTHCHYAPLVTTNLFSLHY

The following DNA sequence Seq-2427 <SEQ ID NO. 92> was identified in *H. sapiens*:

TAGTTTCTCTGGTCTGCCTTGGGGAAGAAAGGAGAGCAGGAGAAAGAAAGGTGGGAGAAG  
GCCAGAAAGACTTTGTTTCTGAAGCTCTTTCAGTTTCCTTCAGTTCAAAGCACTCATCAC  
ACCAAGACACCATACTGTGGGGTATCACATTCTGAGCCCTAACACTTCCAATATTATGCT  
ATGAATTTACATCATGATTTTCAGGTAATTATTCCAACAATGCCACAAGGTGAGCATTGT  
GTTATCCAGTTTCACAGATGCAGAACTGAAGTGGAAAAAATTGACTAGCATTATATGGC  
TGGCAAGTGATCAAACAGGATTTTCTCATTATTTCACTCAATAGTTATTGAGCTCA  
TAATATATGCCAGGCATTATGTGCACTTCATGGATACAGACAGGTACACAGTAAACAAG  
GTGGCCACTGCCCAAATGGAGCTTGCACTCTGGTGGGGAAGACAGATAATAACAACAAG  
AAAGAAGCAATATAACAGATTGGGACAGTGCTATTAATATAAGTAAATGAAGGAGGGATA  
TCATCAGGAGAATCTGGGAAGGAGTGGATGCTACCTGAGACAGGATGGTCAAGGATCTGC  
CTAGTTGCAAAGCACTAGACTTTCCACAACCCCTTCTACCCTCCAGTGGGCCTCTGCAGT  
ATATATGGCAACCAATTCTGGTTTCATGTATTCTACCACTTACTCCAACCTTAGTAAATA  
TCTGCAAAGCTTACCATTGCCTAGCACTCTCAGATTATTTCCCAAGATGCTGCAGAATC  
CTTATAATGTTTCTCAGCCTCAATAGAATGAAAAGCAGGTCTGTGCTTATATCACTTAAT  
GACCAAAGGAAGGAAATTTACAATTAAAGTGACTTTGCCAACTGTGGATGAATTAGT  
TAGGTCACTGTGATCTACAGGTTAGATGTCTGTTACAGCAGTGTCTCTACTTGAGATTCC  
AAGGAGGTTGAAGCTCACTACTCGCCACCCCTCGCACCCC

The following amino acid sequence <SEQ ID NO. 226> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 92:

GARGGEASTSLESQVEDTAEQTSNLITVTLLIHPQLAKYTLIVNPLPLWSLSDISTDLLFILLRLRNIIRIL  
QHLGEIIESAMVSFADIYSWSKWNTNQNLWPYILQRPTGGKGLWKVCFATRQILDHPVSGSIHSFPDSPDD  
IPPSFTYINSTVPICYIASFLLFIICLPQNASSIWAATLFTVYLSVSMKSDIMPGIYYELNNYVNEIMR  
KSCLITCQPYNASQFFPLQFLHLNWITQMLTLWHCWNNYLKSKFIAYWKCGSECDTPQYGVLVVLTEGNK  
SFRNKVFLAFSHLSFSCSPFFPKADQRN

The following DNA sequence Seq-2428 <SEQ ID NO. 93> was identified in *H. sapiens*:

ATCAGCGACACCATCCTGGTTGTTCATCCCTGAGCTGTGTTAAGTAGGCCTTCCCCTAAG  
AGAGTTAAAGGGGCACTCGTGAGATACTAAGAAGACTCCTTCCCCAGCCCCAGGCCCTCC  
TTGTACCTTTGCTCTTTCATTCTGTCTGCTGCCTTCTGGGAAATGATGGGACTGGCAGG  
CTGTACTATGCAGCAGGGATAGCAGGGCTGTTTGCTCTGCCCTCAGGAAGGCAGATAACC  
CCTAGAAACAGGAAGAGCCAAATGAGGTTGTGTAAGTCTGAGGCAGAAACATTAGTCGTG  
AGAGCAAGACTTGCAATTGCAAGAGCCAGGCTGTGTGTGTGTTGTGTGTGTGTGTGTGTGT  
GT  
AAAAGTGGATGGCCAAGAGCCAACCCCTGGAGGGCACGGAGACAGGGAAGAAAACAGAGT  
GAAACAAAAATATTTGTGTAGAAGGCATAAAAGTTATCATCACAGACTCCACTGTGTAAA  
GGCATAACTTGCTTTATTTATCTCTAGTGTATATGAAGTCTAGCCTCCCTTCCATTACG





SACGGFNGLHFYSNISHQLYIYYLKVFLFVVFQFIFQIRSKQNYSWRLCCLHPQYQMFMASTEPGVSMESELRDCLSFSEESVMFSIPEEAEITLHYFFELCAGRHGSEICLSDSNSSSICVLFVVFVAFCIQLPDNFFLMFCCNLVKLLFYKLMFWYFGHQILARGKIRTRSTSCCKLIFLVDVFNGLFCFPICVYFLKSCRCIYEYLFH

The following DNA sequence Seq-2431 <SEQ ID NO. 96> was identified in *H. sapiens*:

CCTGCAAAGTCTCTCCTGCTGCACCTTCCTTCTGAAACCATTAATCACCACGACCCACT  
GAATGAAGCCCAATCTCAAATCACAGTGAAAAATCCTGCAACGTGCAGGGTGATGAGTGT  
TTACATTAGCTGAAATGAAATGATGTAATACCCAGAATCGAGGGAGGGCTGCGATCCAGA  
GTCAGGGCATTGCAAAAACCTCTGTGAAACATAACTTTTCTACATTACAAAAAATGTCC  
TTGCGTTTTAGTAATCTGGCTTCTGTAAATTTAGGATTACTTGGATTTTCTGATCTCAT  
CAATTTGTTTTCCAAATAGAAATTCAGAACTTCCCAATTACTCACTGTTTTAGTCAAGTT  
TAAAAAAGGGTAGCAAATAGAACCCAAAGTGATACATGTGCAAGAACCAGTATCA  
AGGGAATAATAATAGAAGGCAGCCATCCAGGTATGTGGGCACCTGCCATGCTGCAGAATA  
GCAGAGCCTCCCAAGGGTCTAAGTGCCTTCAAAGTAAAGCAACTCCTAAGAAAGACAGT  
ATTTGTTTAAGCCAGTGGCCAATTTTCTTCTATAACTGATGATGAACAAGAAAACCA  
GGAGTTCCTAGCCCTATTATTGATGGGCAACTGCTATTGATTAC

The following amino acid sequence <SEQ ID NO. 230> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 96:

VINSSCPSIIIGLGTGFGSCSSSVIGRKIGHWLKQILSFLGVVFTLKALRPLGGSAILQHGRCPHTWMAAFY  
YYSLDTGFFAHVYTLGSICYFFFTLKQVIGKFISIWKNDQKNPSNPKFTEARLLKRKIDIFLCRKVMFHRG  
FCNALTLDRSPPSILGITSFHFSCKHSSPCTLQDFSLFEIGLHVSVGRGDWFQKEGAAGRDA

The following DNA sequence Seq-2432 <SEQ ID NO. 97> was identified in *H. sapiens*:

ACAAGGTCGGTGTACACCCCTGTGATTCTGGGAGTAATATCTTCTCCTCCCTTGGATAT  
TAGGAACAATATCACGGCGGGGTGGGGTTGTGTACAGCCTCTGCAATATTGGGAGTAA  
TATCATCCTTTCTCCCACTGGATATTAGGAACAATATCACAGGAGGTCTGGACACCCCC  
TGCGATATTGGGAGTAACATCATTTTCTTTCCAGTGGATATTAGGAACAATATTGCAT  
TGGGGTGTACACCCCTCCGACATTAGGAGTAATATCATCCTCTCCACAGTGGATATTA  
GGAACAATATCTCAGAAGGAGTGTAGAACCCCTGCGGTATTAGGAGTAATATCATCCTCT  
CCCTCCCTGGATATTAGGAACAATAACACAGGGAGAGTATACAGCCCTGTGATATTGAG  
AGTAATATAATCCTCTCCCATCTGAATATTAGGAACAATATCAGGGGGGTGGGGTACAC  
CATTTGCGATAGTGGGAGGAATATCATCCTCTCCCACTGGATATTAGGAACAATATCA  
CAAGTGGAGTATACACCCCTGCGATATTGGGAGTAATATCT

The following amino acid sequence <SEQ ID NO. 231> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 97:

QGRCTPPVILGVISSPPLDIRNNITAGVGVVYSLCNIGSNIILSPHWILGTISQEVWTPPAILGVTSFSFP  
SGYEQYICIGVYTPSDIRSNIIILSHSGYEQYLRRSVEPLRYEYHPLPPWILGTITQGEYTAPVILRVISSPH  
LNIRNNIRGVGYTICDSGRNIILSPPGYEQYHKWSIHPLRYWEY

The following DNA sequence Seq-2433 SEQ ID NO. 98> was identified in *H. sapiens*:

TATTTAATCATATAATACTAAATATACTGTATTGAGAAGTTTTTGTGTTTTAGTCAGGT  
AAGATGCAGGGTGTAGAGGTGTTAACCTTCCTTAAATTTAATGGCTAGATATCTTGA  
GATCTGTCTGATGTAGAGTGGAAAGTGGGTGGTTCTTTCTTCCCATCATAAAGGCTC

ACAGCTGATACCCCTATAAAGAAAGACTGGTTAACAGAGAAAAGCACAAACAATTTATG  
 AATGTGAATAAGTATGAGAGCCATACAAAATATGAAAATTCAAAGAAATGGTTAGACGA  
 TTGATGCTTAACCTTCTTCATTAGGGAGAGGAAAGTTGGGGCGGGAGTGGGGGAGTG  
 GGAATGGGGCCCCCTCCATCTCCAGGAGTGGATAATGGTTTGTAATAATTCTGTTTGG  
 ACACTGAATGGAGCGGAATGGAAGGACAAACAATAGGAATGTGAGGGGTGGAAGTGCAT  
 GGTGAACAAAGGTTGTCTTATT

The following amino acid sequence <SEQ ID NO. 232> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 98:

DNLCSPCSSTPHIPIVCPFHSAFVSVQTELFTHYPLLEMEGAPFPTPLPPLQSSPRRLSINRLTISLNF  
 HIFVWLSYLFETFINLLCFSLVNQSFFIGVSAVSLYDGEKNHPLSTPTSDRSQDIPKFGKVNSTPCILP  
 DNTKNFIQYIYMIK

The following DNA sequence Seq-2434 <SEQ ID NO. 99> was identified in *H. sapiens*:

CCGAAGCCGAAAAGTCTGAAACTGGCCCAAAGTGGGAATTTATATCCCTGTTCTGCTGCT  
 GGAATGTTGCCTTTTCTAAACCAACCATGGTCCCGCCCTACACCATCCTGTACCTATAC  
 AAACCCCATACTCAGCCAGTAGACAGGACTATGGTTGGACATTGGAGAGAAGCAGCTTGA  
 TGGCTTAAACACCGAAGAAAATCCAGCCAGAGACGCCAGAACTTCCGGGGAGGGTTACG  
 CTACCGACCCTGTCTCCTTCTCAGCTCCCTTCTGCGGAGAGCCACGTTTCATTACAA  
 TAAAATCCCCCACATCCACCACCTTCAATTTATTCGTGCAACCTCATTTTTCTGGCTG  
 GTGGACAAGAGCGCGGAGCCACAGGTGGAGATACAAAAGCTGTACATTGGCCCTTTG  
 CCCTTGCTGGCGGAGGGCAGCCGCTCACACAGAGGCAGAGGGCCACTGAACTGTTAAC  
 ACTTAAGCCATCTGCAGATGGCAGAGCAAAACAGCACTGGAACATGCCCTCTGGGGCTT  
 C

The following amino acid sequence <SEQ ID NO. 233> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 99:

RSRKVNWPVKGIYIPVLLLECCLFLNHPWSRPTPSCITYTNPILSQTGLWLDIGEKQLDGLTPKKNPARDGO  
 NFRGGLRYRPCLLLSSPSCREPRFIHNKIPHIHPSIYSCNLIFFGWWTREPOVEIQKAVTLALCPCWR  
 RAAASHRGRGPTELLTLKPSADGRAKTALEHALWGF

The following DNA sequence Seq-2435 <SEQ ID NO. 100> was identified in *H. sapiens*:

ATAGAGACGAAGTTAAACACTTAATTTGCAAACTACTGAGAAGTAAATTTCTTGTTCCA  
 AGGTAAGTGGAGTAATTGCCAAATGCAGATAAATCCTCCCCCTGAGTAGGAAGCCCCACA  
 CTGTTTTGAAAACAATTCTAGACTTTGCCCTGTGAAGCTGATTGAATGCTCAACAC  
 AAGACTCCACTGTTGTTAGCTCTCGCTTACTGCTTTAGGGGCGGAGTTAAACACTTTTCA  
 AAAATCCGAGCTTCCCTAATAAATACAGGATTTAGTGAAGATTTGATTGTCTGGGGTT  
 GGCATTCTGAGGACAGAATAATTTATTTGCTCTAAGCAGGTGTGTTATGAGAACAGAG  
 GCTATGTTGATAAGAGATCCCTGGGAGCTGGTAATATATTATCTTCTGTAATTTCTCCA  
 AAAATAGACTTAATGGAAAGAGGATGCATAATATACCCCTCTCAAAGGAAGCGTTCCCC  
 AATACAACAGAAGCAGTCATTCTAAAACAGCTTTATGGCTCTGCAGTCAATAACTCTAT  
 TTCTCCCCCTTACAACTTCTCCTTCTGCTATGTAAGAACTTATGTGAGGGCACACA  
 CACATTCACG

The following amino acid sequence <SEQ ID NO. 234> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 100:

IETKLNTFAKLLRSKFLVPRLELPNADKSSSPVGSPTLFKQFLDFAPVEADMLNHKTPILLALAYCFGRSHF  
SKIRASLINTGIRFLSGVGIPEDRIIFYFALSRCVMRTEAMLIRDPWELVIYYLLFLPKIDLMERGCIIYPL  
SKEAFPNTTEAVILKTALWLCSQLYFLPFHNFPSAMELMGHTHIH

The following DNA sequence Seq-2436 <SEQ ID NO. 101> was identified in  
*H. sapiens*:

AAAAAAAAAACCCTCATGATATGGATATTGTTATCATTCTCTTCTCACAAATGGTAAT  
ATTGAAATTAATAGAGTTGTATATCGTGCCACAGTCACACAGTTAGAAAAGCGTCAGAG  
CCAGGGTTTGAACTCAAGTAGCCTAACTATAGAACCCTATTTTAACTACTATACAGTA  
TTTACTATCTGTTCCATCAAAAGAAATCATTTTTCAGAGTGGAGATGATAGAACATACA  
TGAGAACAAGAGTATTTAAATCCAAGATACCTGCAAAGCATCTAGACACTCTAGATTTAG  
ACTTTTAGCTCCTTGGCCAAGATTAATTACCTTTCAGGAAAATAAACTACATACCAATG  
AGATCACTAGACCTCTCGCAATGATCTATGAAGAATAATGGGAACAGCTATCTGGGTATC  
TAATGGGCTAGAGTCAGATAAATGGTTTCTCAATAGATTTCCAGAATAATGGGAAATTT  
GGTTTTCATTAAACAATAGGCTACGTATGTTATATTCATTCTAG

The following amino acid sequence <SEQ ID NO. 235> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 101:

KKKTPMIWILLSFLFSQMVLKLEVVYRVHSHTVRKRQSQGLNSSSLTIEPIFLITIQYFTICSIKRNHF  
SEWRNIHENKSI IQDTCKASRHSRFRLLAPWPRLLTFQENKTTYQDHTSRNDLRIMGTAIWVSNGLES DKW  
FLNRFPEWGNLVLHQATYVIFIL

The following DNA sequence Seq-2437 <SEQ ID NO. 102> was identified in  
*H. sapiens*:

TCCTTCTCTCTTTCAATCGTGTGGAGAAAATAATTATCAGTTGGGAACCATCATTTTTC  
TACTACCATGAATGCAATGTACTTCCATGACCCATCTTCTTTACGAATAAAGTTACAA  
TATAAGAAATACCACTACACATATCTGAGTTTATCTTTTAACTGTCTTTTAGAGCCCAT  
CTCTTCTGCCTTCCTAGAACCTCTACTATGGATTATCCCTTTACCATAGCATTGTCTATC  
TCTTCTTTTAAATGCATTTGTTTCCCACTGATTTTAAACATGATTGAGTCATTTTCATT  
AGAGACTAAATAAACATCCTCATTACATGGTTCCTAGGACCACTCCCTCTTCAGTTGTG  
TGGAGAACTAAGCTTTTAGAAAAGAGACGTCCAACTCAGTATCTCTATTTCTGCATGCCA  
CACAAATCCAGTTTGATTTTCATCCTCATCAGTCTACTAAAAGATGTCACTAAGGACACC  
AATGAATTCAAAAAAGCCCTGAAATCCAATGGAAATTTGACATTTTGGACCACTTTCT  
CTTTCTTCAAACATTCTTCCCTTAGTTTTCAGATAGTTTCTTCTTCTTCTTCTACTC  
ACTCTATTTTGATCTCTTTGAAAATTCATCCACCTCTACCCAGTCAATAAATGTTAACA

AGTTGGACTGCATTCTGGTTCTCTCTGAAGTTTGCTTTTAGGCAAGTACCAGATGGATTG  
 TATTTTAGAAAAGATTTGTCTGGAACATTTCTGATGTCATTATCCAGAGACAATGAGAC  
 AACTCATTGCTTATGAGGTTTTTACTACAGCAATCTAGAGATGGAATTTCCAATGGAAA  
 TAAAAAAGGGTTTTTATAATTTCTATATTGACACTGGCAGCTCCGCCTTTTAAAAAATTA  
 GTTCCTTTTAAATGAATGTATTTGGGAGTAGATTATAGTGATTTAGTAAATTGGCACTG  
 TGTTTAGA

The following amino acid sequence <SEQ ID NO. 237> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 103:

TQCQFTKYTIISQNTFIKRNFFKRRSCQCQYRNYKNPFLFPLEIPSLDCCSKNLISKVVSLSLDNDIRKC  
 SRQIFSKIQSIWYLPKSKLQREPCSPTAFSSSTQWISYMLNCHVCASLKCAFLFTEMRDVLFMIFSL

The following DNA sequence Seq-2439 <SEQ ID NO. 104> was identified in *H. sapiens*:

TCTCATTTTGAAAATGTAAGTGGATATCACTACTGCATTGCCTGGAAATCCACGAGGAA  
 GATAATGCCATAAATAACAGGGAGGTAGTGCATCTTGAGTGGGATGTTTTCATCAGTGCA  
 ATTTCCAAAAGCAGCTGCATAATCGGGGAAATCAGAAGCATTGCTAAATAGTCTAGTGG  
 CTCATTGATGTTGTCTCCTTTTCATCTTGCAAGAAAACAAGAGAGTTTCAGTTTGGCAATA  
 TGAATCAAATGAGCAGTAACTCGCTGATAAAGGAAAACAGAAAACATTAATGATAGGGTA  
 ATAAAAACAAGGATCTACTTTTAAATGAAAATTATTCTAACATCCTAAATTTGCCACTTC  
 TCTCTCTTTAATCTCAAAGAGACCCTGTGGAGAAGAAATTGAATTTCCAAGAAAATGAC  
 TATGAGGCAAGTTACTAAATGCATCTAATAAAAAATATAAAAGTTAAATTACCATGAGAGT  
 TAAATGAGGGATTGGGAGAAAAAAGCCACATGTCGCTTTGGAAAAACAATTGGCAAGGT  
 CACCATTGGGAGAAGCCATAGGGTATCGCCATTAGAGACTTAACAACAGGACCTACTATT  
 ACCCAAGTGTGATGCATGCCACCATCACTTACTTCTACATGTCACAAAATACTGAA

The following amino acid sequence <SEQ ID NO. 238> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 104:

FQYFVTCRSKWWHASHLVNSRSCCVSNGDTLWLLQMVTLNCFPKRHVAFFSQSLILTLMVILLYFYMHV  
 TCLIVIFLEIQFLHRVSFEIKEREVANLGCNNFHLKVDPCFYYPPIINVFCPLSASYCSFDSYQTELS  
 FLARKETTMNEPLDYLANASDFPDYAAAFGNCTDENIPLKMHYLPVIYGIIFLVGFPNGAVVISTYIFKMR

The following DNA sequence Seq-2440 <SEQ ID NO. 105> was identified in *H. sapiens*:

CCACCTGCTGTCTGCTAGACGTGGAAAGATTGCGAGCAACAGAGCAGGGAAAATGAGTC  
 AAATGGAGGCCAAAAATGAGAACTAAGAGATTGTGAGAATATTCAAGCAAGGCAAGGAG  
 AAAATAAGAGAAGGAAAGTAAATATAGCCACAAGCAAAAGTGGTAACAAAATGCTTGAT  
 ATGAAGTCCTATTTACCAGTGATAAGCCACATGGATAGTTAGTTATGAGCTTTTTTGTAA  
 TCAACAGGAAAAGGAAAATCACAATTTTCAAGATTCCAGTGTCTCTAAGGTATAAAGCC  
 CAAGTAATTGGAGAGAAGCACAATTTTGTGGAATAAGATAAAAATGAATTGCCTCTA  
 GTCAGTTTTTGAAGAGCCACTTGTCCAGGGTCTCACAGCTGCTCGGCCAGAATTGAACC  
 CCAACCATAGTTCCAGAGCCCACATTCTCAGACATAGCCCCCAATACTGCCTCTGGGC  
 TGGAGCTGGTATTCTCAATAACTGTTTGTGAGTGGATAGGTGAATCACCATT

The following amino acid sequence <SEQ ID NO. 239> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 105:

WFTYPLNKQLLRIPAPAQRQYWGLCLRMWALELCGWSNSGRAAVRPWTSGSSKTDQRQFIFILVPQIVVIL

SNYLGPIPRHWESKLPFSFCLQKSSLIHVAYHWIGLHIKHFVTTFACGYILLSFSYFLLALLEYSHKSL  
SHFWPPFDSFSLCCCESFHVQDSRW

The following DNA sequence Seq-2441 <SEQ ID NO. 106> was identified in *H. sapiens*:

TATCCACATAAATGTGCATTTTCTTTGGGCCAAAATGAGGCAGAGGTGTCATGTGAATT  
TTTCATTCCTTCACACAACGATAGTCTCTCACAAAACAAAGAACAAAAGGAAACATATGT  
TCACAGTGGGAAGGATTATTACTCGATCATCTGTATAAGCATGGCCCAAGGAGCCTTTGC  
CAACCTACTGGGGATGTACATGTAAAAAGGTTTCTCCAAAAGGTTGGCAATATGATTTA  
TTAAAGGAGTCAGATGACATGGGAGTTAAGGGCAGCAAACCTCATTGTGATGGAAAGGAT  
CTAAGCTGCTCCAGCAAATGAAAGGATTATGGTTCACCTGCCAACACTGTGCAATTTAT  
GGATGAAACCTCAACCACGAAAAGTGAACCTTCTTTGTGTGTGTATGGGGTTGCGAGG  
GGAGACATAGGAAAGGAAAGGCAGACAGACCGTGGAAAACAGATATTTCCCTGGATAAG  
AGTGGATGGCCAGTCTCATAACACTCATGTATTATAGAATTAAATATAAACCTGTTTCA  
GAAAGTACAATATTAAGACCCTTTTAAATCTTGATATTCTTTGATGATATCTCT

The following amino acid sequence <SEQ ID NO. 240> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 106:

STMCIFWAKMRQRCHVNFSFLHTTIVSHKTKNKRKHMFTVGRIITRSSVAVWPKEPLPTYWGCHMKGF  
SKRLAIFIKGVRHSGSQQTSLWKSGSKLLQONERIMVHLPTLCNLWMKPQPRKVKLLCVCVWGCEGRHRK  
GKADR  
PWKTDISPGEWNGQSHNTHVLNITCFRKYNIKTLFKSYSLSMIS

The following DNA sequence Seq-2442 <SEQ ID NO. 107> was identified in *H. sapiens*:

TTTCCTACTGATCAGAGTTACTGTAGAATTTGATTTAGGTGTGTAAATTAGTCTGAGGCA  
CACATTCAGTCTTAGGCAACCCTCTCTGTGATGGCATGCCTCAAAGCAGTGGTTTGAATT  
AGGGGCAACCTTCAACCCTGAGGGACACTTGGCAACATCTTGAAATATTCAATGGTCTT  
AAGTGAGAAAGTGCTATTGGCATCTGGTAGATTCAAGCCAGGGATGATGCCAAAGATTG  
ACAAAACACAGAACAGGCCATACACAGAGAATTATCTGGTCCAAAATGTCAATGGTGCC  
ATGGTTGACAAAACCTGAGATAAGCTTAGGGAAGGATCCAGCACAGAGCAGAATGTATTC  
TCTCTGTAAAGAAGCCAATCCCAAAGAGAAAGAAGTTGAGTAATGCTGCGTATATTACT  
CACTTTCTCTTTCCAAATTTCTTAGTTTGATAATTCACTCGACTTGCCCTGGTAAGGAAT  
GAGGGAGGAAGCAAAAAGACCAAGCTTGTTTACCTAATTACTGTCCCTCAACAGAAA  
AACGTGAGGTGAGGGGTAAGAAAGTCCCCCATTCTCACATCTATATCCAATACAT

The following amino acid sequence <SEQ ID NO. 241> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 107:

VLDIDVRMGGLSYSPSPHVFLLRDSNCNTSLVFFASSLIPYQGKSSELSNEIWKEKVSQYTHYSTSFSLGL  
ASLQREYILLCAGSFPKLISGFVNHGTDILDQIILCCMACSVFCQIFGIIPGLNLPDANSTFSLKTIEIF  
QDVAKCPSGLKVAPNSNHCPEACHHREGCLRLNVCLRLIYTPKSNSTVTLISRK

The following DNA sequence Seq-2443 SEQ ID NO. 108> was identified in *H. sapiens*:

TTTGCTCTTTTCTATGTTTCATCATCTCATTGAATGGCACCCCATCTGCATGGTAGCC  
TGGGAAATATATTAAGGTATTATCCTTGAACCTTCTTTCTTTATCATCCCTATGTCCAGG  
TAATCTGAAATCTGTCAGAAATATGCATCTTTAATCTATCTTAAACTGGCCCATTTTAA  
AAATTTCTATCTTACCTTACCTTTACCTAAATGATTATCACTCTCCTAATTGTTTC  
CTAATGGGCCTCATAGGCAAGACAAATCTGTTCTTATACTGCCTCTAGAATTATCTTTT

CAAACACGGATGTGGCCATCCTTCTTTCTTACAAATGACCTCATAGTCCCAAAGACAAAG  
TCTATACTCTCCCTAAATAACATTCAAGGCCCTCACTCACGCAGCTCCCTGATTCCCACG  
TCAGTATTTTGTCTCCTCCCCTTCCCAAAGCACACTCTCACATACGCGTTATTCTACC  
TGGAGTCATATTAAGCTACTTTCAATTCTGGGCTTTCTTTAGCCTTCAACCCCTCTCTTA  
GGCTGGTGCAATCCTGGGGAGTGGTCCAATCCATGCACGTGCTACCATGCACCCACCTTT  
CTT

The following amino acid sequence <SEQ ID NO. 242> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 108:

FALFFPMFIISLNGTPICMVAVEIYGIILEPSFFIIPMSRSEILSEYASLIYKLAHFKFLSILTLLYLNDY  
HSPNCFLMGLIGKTNLFLILPLELSFQTRMWPSFFLTNDLIVPKTSILSLNNIQGPHSRSSLIPTSVFLS  
SSPSQSTLSHTRYSTWSHIKLLSILGFLLA FNPLLGCIPGEWSNPCTCYHAPTFL

The following DNA sequence Seq-2444 <SEQ ID NO. 109> was identified in *H. sapiens*:

CTGCATGTTGTCTATTGGTCTGATCCATGGGTTGCTTTTGCTCCAAGGTCCAGGCTAAAG  
GAGATGCCCTCTCTTGGGGAATGTCATGCCCTGCTAGAGGTAGTCTCTGCTTGGACTGG  
GCACACTGCTACTTCGCTGCTCATTTTCATCAACCCAGCCAGCCACTGTGGGGCAAGCCA  
GTGTTCTCTTCTTGTGAGAGATGCTGTACTTTGCATACAATGGTGAAGAGAGTGAACAGC  
AGGGTGTAATTAAACAGTCAACCACAACCTGAAGCCACTTTCCCTGCTAAGTGGACCTCA  
ACTCAATGGTCTCATTTCTGAAAGATGTGGCCTAAATTCTTGCTTGAATGGTAATTCCTC  
TCTAATAGACTCTGCTGTTCTCTTGCCAGTCAAGAGGACTGAAGGGGATTGAAGGTCTGA  
ACCTAGGCTCAGTGGCTACTGCCCTCCTCCACAGCCGCTGGCTTCCAGCAGACATTCCT  
GATGCTGATGTGCTCCTTGGAGTCTGAGCTTTGGGGGAAATCCTGTTGCATGGTGCCAG  
ACCTCTCCTTCCCCATCTCATAACTCCATCACAGAG

The following amino acid sequence <SEQ ID NO. 243> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 109:

LCDGVMRWGRRVWHHATGFPPKLSTPRSTSASGMSAGSQRLWRRGSSHAVQTFNPLQSSLAREQQSLLERN  
YHSKQEFRPHLSEDHVEVHLAGKVASGCGLFNYTLTFLFTIVCKVQHLQARNTGLPHSGWLGLMKAQKQC  
AQSKQRLPLAGAHSPREGISFSLDLGAKATHGSDQTT

The following DNA sequence Seq-2445 <SEQ ID NO. 110> was identified in *H. sapiens*:

TTGTGGAGCAGTTAGAGACACATGGCAGTGTCTTGAGTGGCTCTGAGTGTGGGACCATT  
TTCTAGGTGATCACTCAGCATAGCTTACCGATCAGACTCAAGTGAATGGAACCTGCCCTC  
TTCCCTTTCTCCTGGCTTTGGAACAGTTGCTACCAGGTGAGTGGTTTTTCCCTCCAGAC  
AGTTACTGAGAGTAATCCCTGAGCACTCACTGGGTGCCTGTTCTGTGCTGACAGTCATCT  
CATTTCATCCTAACAGCAATTCCATTCTGCATCTTCTCTGGACACCCCAAGGACCATCCAG  
GACAACCTGCCTGACACCGGCTAGTGTGGCTCCATGATAACAAAGACGCAGGTCCAG  
AGACAATCCCCCTACATGGTGCCTGCATCTGATTCCCCTTGG

The following amino acid sequence <SEQ ID NO. 244> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 110:

VEQLETHGVSLEWLVDHFLGDHSALTDQTVNGTCPLFPFPGFGTVATRVVFPQRQLLRVIPHERSLGACS  
VLTVISFILTAIPFCIFSGHPQDHPGQPCLTPLGLVWLHDNDKAGPETIPLHGACIFPL

The following DNA sequence Seq-2446 <SEQ ID NO. 111> was identified in

*H. sapiens:*

TCTTGCACTCTGGGCCCCCAAACAAGAGGCCACTCAGAAATCACAGTTTGAGAACAAGGC  
 ACCATTGCCCTGAGCCTGGGCTTTCCTGAGGCTTGGGTAAGAGAAAGAGAGATGAGAA  
 GGCTCCCTGGGCTACAGAGGTCTGGAGAGAAGCTGGCACCTGGGAAGAACAATTTCCCCA  
 GCAGCTAGCCAAGCTGGGGTCTTCCAAGTGGATGCAGAGACCTGCCCTGCTGCCCTCCCC  
 ATCCTCTGAGAGTGCCTTCTCTGGGCTTTTGCTTCAAAGAGCCATCTTTTCCACATGGC  
 ACTCATCTTCTTGTCTTTGCTTCATGACACCTTGAGCGTGTTAGAAGCTAATCCTGAA  
 CAAGCATAGAAGGGGCACTTGGGGTAGGAGCTGCAGTGGCACCACCCGAGAGGCCAGCTT  
 TACCTCCCCCAAAGATCCACTGCCAGAAGGAAGACCAGGGGCTCCCTGGTGCCAAGG  
 GCTTGAGAGTATGCATCCAATGCAGCTAGGTCCTCCACACACTGTGGTGGGGCCCCTCAC  
 CCTCAGATCAGCATCTTACTCTCA

The following amino acid sequence <SEQ ID NO. 245> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 111:

ESKMLIGGAPPQCVEDLAALDAYSQLGTREAPGLPEFVAVDLWGRSWPLGWCHCSSYPKPCFYACSLASN  
 TLKVSSKQGGRVPCGKRWLFEAKAQRHSQRMGRAAGQVSASTWKTPAWLAAGEIVLPRCQLLSRPLPREP  
 SHLSFSYPSLRKAQAQGMVPCSQTIVISEWPLVWGPRVQ

The following DNA sequence Seq-2447 <SEQ ID NO. 112> was identified in *H. sapiens*:

TAACAAACACTTTTTATCATATATGAACTCCTGTACAATGATTGGCTAGAAGAAAA  
 AATAGTTGGAAGGTCAAATTTGTTTAAACATCTGTTCAAAGCCTGCATTAACTTTT  
 ATCTGTCTGACAAAACATGTCTCAATTTCTTCTAAAGCAGCTCTATTGCTCTAGCATA  
 TGCTCACCAGTTCTTTAAAGGGCATTTCCAACCTTAGTCTGACAATGAAGACACAAA  
 GTAGGTTAGGTTCCAAACCACCTTCCTAGCCCTCCCTGTAGAAAATACCATGTTGCAC  
 AGTTACATGTGTCCCCTGACACAAACGACACTCATTTTACGTAGGTCAGTGGACCTCAA  
 CTGTTGTTGCTGTCTCCAGCCAATTCAAGAGTGAAGGAAGATGTAACCAGACATACA  
 TATCTCCCTTTCT

The following amino acid sequence <SEQ ID NO. 246> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 112:

QNTFYHINSCTMIWLEEKNSWKVFKVLKHLFKSLHTFICPDKTCLNFFLKQLYCPSICLTKFFKGHFQPFQ  
 RHKVGVPKPPFLALPVENTMLHSYMCPLTQTLLILRRSLDLKLLLVANPANSRVKEDVTRHTYLPF

The following DNA sequence Seq-2448 <SEQ ID NO. 113> was identified in *H. sapiens*:

CAGTCCAATGCTCCAGTTTATAGATTGGGAAAACCTAGAGCCTAAGGGGTCACTTGTTA  
 TAGCTCCTATCCCCAACTTACAAAACAAAGAGTTTACAGAATGAGTCAAATATAATTT  
 GTTTGGGCTACTATTTCAATTTTACCATTTTATCCCTATTAGTATTTATCACCATACATTC  
 AAAGGAATTCATACATGTAGACACATCTGAGGTGTTCTGATTCTCCTGTTGACCTGT  
 GGTAAACCTCCTGTGGCACTATAGCACCTTTAGCTTATCAGTCTTCTTTCCCTCACCTCA  
 TAGATCAGAACTTATCAGCCCCATCCTGGTCCTTCTGAATCTTTGTCAAGTCATTGCT  
 TTCCAATCTCTGATAAAGTGTGAAAGGTACCATATGCCTCTCAGAGATACACACAGT  
 CATGTGCCACCTAACTATGTTTCAGTCAGTGAGGGACCATA

The following amino acid sequence <SEQ ID NO. 247> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 113:

SPMLQFYRLGKLRAGVTCYSSYPQTYKTKSETEVKYNLFGLLFHFTILSLLVFITIHSKEFIHVDTSEVFL  
ISPVRPVVKLLWHYSTFSLSVFFPSPHRSELISPHPGPSEFVKSLLSNLSVERVPLCLSEIHTVMCHLTM  
FQSVRDH

The following DNA sequence Seq-2449 <SEQ ID NO. 114> was identified in  
*H. sapiens*:

CCAATACCACCATCTGAGGGTCTAGAGAAGGCTTGATTTACTTTCATGAGTCCCGGAATA  
AGATCTCCTCAAACAAGGAATTTTTTTTAAATCATGGAAGTATGGCAATGGGCAACTAAA  
CCAAAAGTCTCAGTGCTCCTCTCAGATATAGCTTCGCTCAGAAACAGGCAGCCTGGGTAG  
AGAGATGGAATGTAAAGTCTTATTAAATGCTCAGCTGAAGTGTCAAGTAGGGGGCTTTGG  
TGCTGTCTTCAGGATGTAATATATGTACTAAACAGTGACCGAATACTATACAGAAATCA  
GTAGTACCTAAAATACATGGATTTTATACCAAGGCTTAGACATAGAATCAGCACTTGTA  
ACTATCAAATGGTTGAGGAATTTCTACTTCATTTGTCCACAATTACGCTGGATTAGAAGT  
GTTTGCATCCTTGCATCTGTGTGT

The following amino acid sequence <SEQ ID NO. 248> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 114:

PIPPSEGLEKAFTFMSPGIRSPQTRNFFLIMEVWQWATKPKVSVLLSDIASLRNRQPRDGMSLIKCSAEV  
SSRGLWCCPSCGNICTKPVTEYYTESVVPKIHGFLYQGLDIESALVTIKWLRNFYFICPQLRWIRSVCIL  
SVC

The following DNA sequence Seq-2450 <SEQ ID NO. 115> was identified in  
*H. sapiens*:

TTTGTACAAATATTAAAAGTGTGTCCAAGGTCCAGAGATAGCATGTAACACTAACAAAT  
CTGTGGGATGGTGGTGTGATGTCAATACCAAGAAAAGCTTTCAGAGAGCTTGGGGTTTCAG  
CCAAGACTCCACAAAGGCATAGGGGCTTTGTGGGAGAATGGCAGTCTCTCTGGAGAAGTG  
GCAGATAAAAAGGTAAAGATCTGTGAGCAACGTCATCTTGAGTTCAGGAATTGACAATAG  
TTTGGTATTAGAAGAAGAGTAAGAGTGTCAAAGGAGCATTGTGTGAATCTTTCACTCCA  
GAGATTTTAAATCTCTTAATAGAAAGTTGTTTGTATTGATTGAATGATTAACTTTATTA  
AGAATTTTGTGTCTCAGGCACTGGATTAGTAGCTTTACACATTTCAATTAAATCTCACA  
TTTTGATAGCTTCTACTATGGTTATTATTTTACAGAAGAACTGAAGTTAAGA

The following amino acid sequence <SEQ ID NO. 249> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 115:

LTSVSSVKPKLSKCEIMKCVKLLIQCLRQONSRLIIQSIQTTFYGDNLWSERLHKCSFHSYSSSNTKLLSI  
PELKMTLLLDLYLFICHFSRRTAILPQSPYAFVESWLKPQALCKAFLGIDITTIPQNLLVLHAISGPWTHE  
YCNK

The following DNA sequence Seq-2451 <SEQ ID NO. 116> was identified in  
*H. sapiens*:

CCTGAAACCATGGGCTCTTCGTACCTCCAGTGCCGCTCACATCTTATGACACATAGTAGG  
GGCGTTAATAAATGCTTATTAAGTTGACGACTATGCCAGAAAAGGGTGAGGGATTACAC  
AAAGTTTTAAACAAATCTCACGGTAACTCTCAGAAGCAAAAATAAAATAAACATTTA  
ATAAAGTGCTCTCAAGGCCTGCAGCCCAATTCAGGTTTGCTCCAAATGTTGATGGC  
CTTGAGCTTTCTTGTGTGAAA

The following amino acid sequence <SEQ ID NO. 250> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 116:



FTQESSRPSTFGANLELGCPRPAGTFIKCYFFIFASEELPDFVKTLCPSPFFFWHSRQLNKHLLTPLLCVIR  
CERHWRYEPMVS

The following DNA sequence Seq-2452 <SEQ ID NO. 117> was identified in  
*H. sapiens*:

CTGCTCCATGGGGATGGGCCTCAGTGAGTGTATGTGCCAGGCTTGAAATGGCTTCACGGT  
ATGGGTTGCAGGAGCACCATGAGGTTTCATCTAATCTTTGCCTTCCTCTGCCAGCATGTGT  
GCCATCTGCAATGTCTCACTGAGCACTGAGTGGGCCTGCTATGTGGGCAGTATCCCTGC  
CATCTTCATATCA

The following amino acid sequence <SEQ ID NO. 251> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 117:

APWGWASVSVCARLEMASRYGLQEHHEVHLIFAFILCQHVCHLQCLTEHVGPMWAVSLPSSY

The following DNA sequence Seq-2453 SEQ ID NO. 118> was identified in  
*H. sapiens*:

ATCTCATTTGGTATGTAGTTTTATTTCTGAAAGGTAATTAATCTTGGCCAAGGAGCTAA  
AAGTCTAAATCTAGAGTGTCTAGATGCTTTGCAGGTATCTTGGATTTAAATACTCTTGTT  
CTCATGTATGTTCTATCATCTCCACTCTGAAAAATGATTTCTTTGATGGAACAGATAGG  
AAAATACTGTATAGTGATTAATAATATGGGTTCTATAGTTAGGCTACTGAGTTCAAACC  
CTGGCTCTGACGCTTTCTAACTGTGTGACTGTGGACACGATATACAACCTCTATTAATTT  
CAATATTACCATTGTGAGAAAAGGAATGATAACAATATCCATATCATGGTGGGTTCTTT  
TTTT

The following amino acid sequence <SEQ ID NO. 252> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 118:

KKEPTMIWILLSFLFSQMVLKILIEVVYRVHSHTVRKRQSQGLNSSSLTIEPIFLITIOYFPCSIK  
RNHFSEWRNIHENKSIIQDTCASRHSRFRLLAPWPRLLITFQENKTTYQD

The following DNA sequence Seq-2454 <SEQ ID NO. 119> was identified in  
*H. sapiens*:

AGAGATCTTTAAATACTCAAAGAAAATGTACCTAGAATTTGATAACTCTTGAAAATA  
TCTTGCAAAAATGAAGGCTAAATAAATGATTTTTTGACAAAGAAAGCTGAAAAAATTTA  
TTGTGAGCAGACCTGTACTACAAGAAAGGTTAAAGAAGTTATTTAGGTAGAAAGAAAAT  
GATATCAAATAAGCAGATCTACACAAAGGAATGAAGATCTTCAGAAATCGTAAATTTGTG  
GGTAAATCTAAAAGCCATTTTAAAAATTTGAGTCATCTTAAGATTATGTCTATAGCAA  
AGAAAAATGCTAGCAATTTGTTATGAGGTTTAAATATGCAGAAGCAGAAGTAAATCATA  
TAATGATAGCAACATGACAACCTGGGGGAAAATGAAAGTCCACTGAAGAAATGCTTAATAA  
ATGTT

The following amino acid sequence <SEQ ID NO. 253> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 119:

TFIKHFFSGLSFSPSCHVAIIIFTASAYFKPHNKLIAFFFAIDNNLKMNTQNFNGFIYPQFYDFRSSFLCV  
DLLIYHFLSTITSFNLSCSTGLLTINFFSFLSKNHLFSLHFCKIFSRVIKVFVITFFEYFKDL

The following DNA sequence Seq-2455 <SEQ ID NO. 120> was identified in

*H. sapiens:*

ACTTTCCTTTCCAGGCATTTCTTGATGTGGAAGAGATTACTGAGTCTGATACCTTTAAA  
 GGTCTGACAAGAGACATTTGCTGCCTATGCCTTCTGTTCTCTTGAGGAGTGCTACCAAT  
 AAGGCTTCGTCAACATAACAAGGCCACCTTAGCTAGACAGGCCTCTTCCTTTCTTCCTCT  
 CATAACCTGTCTTGCCACTAAACCTGAATTACCAGCACAACCTCTTGGGGCCATGCTCT  
 GAGCCACATTTCTTTCTATAACCTCAAGTAGGTATATAAGCTTCTGCGCCTTATTGTCTT  
 CATTCTGAAGGCTCTTATGTACATGCATTAAACAAATTTGTATCTCCTATTAATGTGCCT  
 TTTGCGAGTTGATTTTTCAGTGAACTTCAGAGGTCCAACGGCAGTAGCCCTACCAAGT  
 TCAAGATGCTCCACTTAC

The following amino acid sequence <SEQ ID NO. 254> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 120:

TFLSRHFLMWKRFTESDTFKGLTRDICCLLFSWRSATNKASSTQGHLSLGLFLSSSHNLSCHTITSTTS  
 LGPCSEPTFFLPQVGIASAPYCLHSEGSYVHALNKFVSPINVPFASFFSETSEVQRQLPSSRCSTY

The following DNA sequence Seq-2456 <SEQ ID NO. 121> was identified in *H. sapiens*:

GTGATGTAAGACTGGTGGACTTAAATTAATTTTTTAAAGGCATCATGGGATTTTGTATCG  
 GCTATCTCTGTATCTAGAAGATGTCAGACTCATGGAAGTTTTGTCCATTTTATCCCTTT  
 GCTTATCCATTCTTTCTGTTTACAGAAAGACTTAATTTCTGTCTCATATCTCTGTCTT  
 TCTTGCCCTACTATTTTCCCCCTTCTCCAAAATCCAGCCCCAAAACAGTCTACATA  
 TTGTGAAAAGATTTCTCAAACCAAGGGTGATGTAACCTTAGGCCTGTGTTTCTCTC  
 TCACACACACAAAATATTGGATATGAGTGAGATTTTAAAAAATGGTTTTTAAATGTGAT  
 GAAAAGAGTGTCTTTTCAACAGAACAAAACAACCTTAATGCTGAAGCCTCTTCCCGA  
 TATGGGTGGCTTCCAAATATGAAGAAATCTGTGCATTGGGCCACAGGCTCCAGACAAAGT  
 CT

The following amino acid sequence <SEQ ID NO. 255> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 121:

CKTGGLKLIFRHHGILYRLSLYLEDVRLMEVLSILFPLLIHSFLFETERLNFLSHISVLLAPLFFPLLOKSO  
 PQKQSTYCEKDFSNHKGDVTLGLCFLSHTHKILDMSEILKNWFLNVMKRVSFSPQNNPCSLLPDMGGFQI  
 RNLICIGPQAPDKV

The following DNA sequence Seq-2457 <SEQ ID NO. 122> was identified in *H. sapiens*:

CCTTGGCAGCTCCAACCTGAACATGTAAAGGGTGATTCAACAGACAAGTGAGAGAAGGA  
 ACCTCACACAGCCTGAGTGGGCCTGAGATAGGCTGAGGGGCCTAAGCTTCAATTGCATAA  
 GCAGGGCTAGGTCACTCCAGTTACCAAAGACAGAAACAGATAGTCCAGAGCCGTCCAGGG  
 GATGCTAGCCACTGCCAGGAGATGATCAGAGAACACACAACAGAAATCAGAAAATGTAG  
 TACAAGAAGAATTTGCTGATAGGTGCAATCGCCTCAGCAAGGCACAGGAACTCAACTCA  
 GAAGGCAGTCTGTCTGTCATCCACCAATTCTCTGGGTCAAGTCTGATGTGCACTCATAAA  
 GTAAAAATGCACTGTTATTGTGACTGAGAAAAAAATAAAGCTAAAAGGTAAGTGCCTAT  
 AAAATAAGATTTTACTAATGCAACAAAAGCCCTAAAGAAGTGTGGTTTGAGCCCAGTGT  
 CCTCCTCTATTAGACCAACAATGGATAGGTGGTTGAGTCTGTCAAAATGCCTCTGGGTT  
 TACAGAAATGAAAGCTTGGTCTGTGCCC

The following amino acid sequence <SEQ ID NO. 256> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 122:

GHRPSFHFCKPRGILTDSTTYPLLVLIEEDTGLKPHFFRAFCISKILFYRHLPPSFIFFLSHNNNSAFLLY  
ECTSDLTQRIGGQTDCLLSVSCALLRRLHLSANSSCTTFSDFCVFSDDLGLSGHPLDGLSVSVFNGWS  
DLALLMQKLRLPLSLSQAHSGCVRFLLSLVCIHPLHVQVGAAG

The following DNA sequence Seq-2458 <SEQ ID NO. 123> was identified in  
*H. sapiens*:

CATTTTTACCACATATACTATAAGAATTAGTATTATTTTATTAAATAAATGTTATT  
TTCAGAGGTGCAATTTTGTCTTTCAGTAAGATTCTAATTTAAGGAAGTCATTTTAAAG  
GCTAAATTTAATGAGAAAAAGAGCTTGTGCACTTGTGATCCAGTTGGATCCAGTTT  
CTCTGCTGGTCCATTTTGTATCCCTTTGAGTTTGCACTTCTTTTAAACATTTTGG  
TATAGCAGATTTTATTTTGTGACATTTGTGCACATAAACTTCTTGGTGTGGAGGAGA  
GGTTAAATTTAATAGCTAATGGGACAAAGGTATATAGGGATATATAGGTACAACCCCTAG  
CTCTTATTCTTTCTTCTCCATAGTATTCTGGTGATGTAGGGATAAAATT

The following amino acid sequence <SEQ ID NO. 256> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 123:

HFLPHILELVLELIKINVIFRGAIFCFQDFKEVILKAKFKEKELVALVDPVGSSEFLCWSIFCIPFEFAFL  
FNIFWYSRFLFFGTFVHINFLVWRRGILIANGTKVYRDIRVQPLLEFLFLHSILVMGN

The following DNA sequence Seq-2459 <SEQ ID NO. 124> was identified in  
*H. sapiens*:

CCAAGCAAAGTTATTTGTATTTTATTTTACATTTATTTTGTATATTCCTTTTATCTA  
CTTAGGTTTCTCTCTACTTCCCTTTTAAATGAAGAGTTTAAATGCATGTATCTGTGTGT  
TTGCTTGAAAAAACACCAAGTATAACATGTTCTATCTATGAATACTTCTGGCCATTAA  
CTCAAAGGTACTATATTACAGACAGAAAAAGCACCAGAAAGCAATCAGGGACTTCATCTA  
AGAGGTAGGACAGCATAGTTGGTAAAAATACAGACCCTGGAGGCAAACTGCCTGGGCTTG  
AATCCCAGCTTTATTACTTTGGGAAAACTACTTATCTTCTTTACTTGTTTTGGTATCCAT  
GTCTGTGAAATGGAAGTAATAATAATCTCTCATAGCATTGTTGTGAGGTTTCAATAGAT  
GAAGTGAAGACTTTAGAAGGGCACATGATAAGAATTATATAAGGGTTACCTATTATTGCT  
ATCCAATTTGTCATAGCAAGCTAAGGGACCTTGGGCAAGTTACTC

The following amino acid sequence <SEQ ID NO. 258> is the predicted  
amino acid sequence derived from the DNA sequence of SEQ ID NO. 124:

KQSYICILFYIYFVIFLLSTVSSLLPFLIEEFNACICVFAKKTSPITCSIYEFWPLTQKVLYYRQKSTRK  
QSGTSSKRDSIVGKNTDPGKLPGLSOLYYFGKTTYLLYLFWYPCLNGSNNNPLIALLGFNRSDFRRAH  
DKNYIRVTYYCYPICHSKLRDLGQVT

The following DNA sequence Seq-2460 <SEQ ID NO. 125> was identified in  
*H. sapiens*:

ACTGGTAGAATGGGCTCATTCAAGCATGTAACGCCCTTAAATTTTTCATTTAAATTTTCT  
GTGCCTTAGAAATGAACTTTACAGTAATCTTTGCTTTCTAAAAATAAATGTGTTTCTTGT  
TAAGCATTTAGTCTCATCAAAATCTGTTTGAAGAAAAACAACAGAAATAGTGAATG  
AGAAGGGTAGGAGACTTAGGACTCAGCGAATCTATCTCAGTGCCAAGACTTTAAACTG  
GGAATAAATGCTACTTCTCCATGACCTGGGTCTGATAATTTGTCTGAGGAACACTGTTT  
CTAGAGGGTGGTGTGGTACAGTGGGAGGAATGGACTTTGGAGTGAGATCCATGTTCAAT  
CCCAAGTCACTACCTTCTCTGATCCTCAGTTTCTCTCATCTGTAATAATGACCATAATCAA  
CACCATCTCGAAGATTTGTGGTGACAACACAGCATTTACTTCTGCTGTATACTTCCCAT

TTCTCTTGTAGAGACAGAATTTTCCACTTTATTTTAATCTATAATTATGTAATCCCAT  
TAAAAATCACCCCTTCGACTTTCAGTTCACCAAGGC

The following amino acid sequence <SEQ ID NO. 259> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 125:

LVEWAHSSMRPIFHLNFLCLRNELYSNLCFLKINVFLVKHLVSSQILFKKTTENSEEGETDSANSISVPRL  
NWEMLLLHDLGLIICLQEHCFRVVWYSGRNGLWSEIHVQIPSHLPSLILSFLICKMTIINTISKICGDNTA  
FTSCCILPISSCRDRIFHFILIYNYVIPFKNHPSTFSSTR

The following DNA sequence Seq-2461 <SEQ ID NO. 126> was identified in *H. sapiens*:

ATTGCTCTCTCTAGATTTTCTAATGTTGGTCCGTCCTTCGTAAGTTGTGTACAAAGC  
TGGATCCAGTACTCCAAGGGTGATCTGACCTCACAGAGCACAGTGCCTGGGGAGTGCCCT  
TAATCTGGACTTGAATTCATCATAAGAGGCCAAGTCTCTGACCATGATGTTCTCTCT  
GTGTAAGTGGGGGTGCTGAAACCAAGTATTTGTGAGCAGTGCCTGCTCCAGCCATGCT  
TGTGTCTTTTAAGAAGTGACAGTAACTGCTATTTGTGGAGATGGCTATTCATAGGGACTC  
CTTTTCTTTTGCTGACAGAGGCCAGTGTCTAAGCTCTAAGAGGGGCTCTGATGCCAGC  
ATGTGAGTCACACTCACTTGCTACTGTTCTTTCCAGAGTTTGGGCCACTTGTTGCTGC  
ACATCACTACCTCCTCTCCCCCTGCCAGCTTGCAATGTCGCCCTTCCCCTATCTACCATG  
CTGTCCTTGAACATAAGGCGCTTCTCTGCATTCCATGTGTCTACTTTGTAGTTATGTGCT  
GCATTTTGAAGAGCTGAATCTATGTCCAGGTTCAAGAAAGAATGCTGATCAACTGTTGG  
CAATAGATGGGTTTAATATATCTTATGATTGGTTCTTG

The following amino acid sequence <SEQ ID NO. 260> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 126:

CSLLDFMLVLGALRKLCTKLDPLVQGSDDLTEHSAWGPVLIWTWNSIIQRPSLPCSLCVTGAAETQVLSASA  
GLQPCLCLLRSDSNICYLWRWLFITPFLCLTEAQCSKLEGLCQHVSHTHLLFFSRVLGHLLHITTSPP  
AQLALSPFPYHAVLEHKALLCIPCVYFVVMCCILKELNLCPSGRKNADQLLAIDGFNISYDWFL

The following DNA sequence Seq-2462 <SEQ ID NO. 127> was identified in *H. sapiens*:

TAGTCTAGACTCTTTTTCCCTTTTAAGGTCAGCTGATTAACCTTAATCCATCTAATAC  
CTTGATTTCCCTTTGCCATGTATGTCCTGGGGATGAGGATGTGGATGGATCTAGGGGGG  
CGGTATTCTGGCTACCATAGCTATCTTGCTCTTTTGTATTATAATTATGATATGTTCCAA  
AAAGGAGTAAACGTAATACAAGAAGATAAAATACATTTACCATTAAAGTAAGAAAAAAG  
ACAAGGGAGAAGAGAATAAGAAAATGAGTCAGGAGTGGGATTATACAAAAAATTAGTGA  
GTCCACTTTACTTCTGGAAGTGGATGGTGAGCTTTTCTTGCCAGCCTTCTTGAAGAGGG  
AAGCACTGTCTAGTTATGTTGTAGTGTGTCGATCTAGTAAATCCAAGTGTGTTGAT  
ACCTAGATGAATATTCTTGATAGGAAGATGAAAAAAATTTCTTCCAAAGTCTTCATGG  
ATACATAAAGTGTATAATGAGCAAAACCTTTGACATGTTTACAGTAAACCAATGGTGTG  
TTTCACCTGGCCTTTCTCTTTCTGTTTACTG

The following amino acid sequence <SEQ ID NO. 261> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 127:

QTKEEKGOVKHTIGFTVNMSKVLLIIHFMYPRWLKKFFHLPKNIHLGITTSWILLDRHTTTLTVLPSSR  
RLARKAHNPLPGSKVDSLIFCINPTPDSFSYLLPCLFSYLMVNVFLSSCITFYFLEHIIINKKSKIAM  
VARIPAPLDPSTSSSPGHTWQREIKVLDGIKVNQLTKGEKESRL

The following DNA sequence Seq-2463 SEQ ID NO. 128> was identified in *H. sapiens*:

CATCTATTTCGACGACCTTGAGTTACCGCTGAGACATTTCTGAGGCACAACACTAAGAAAA  
CGCATGTAATTGTCAAGCGTGGCAGGGCAGTATTGCTCTCAAAGTCCCGTCTGACTGACA  
GGGCAGAGGTTCTTCCTCACTGCCCCGAATCTGCTTCCCGACAGCTCCAGGGTTCCCTCAG  
GAAGCCGCCCTCCACCTTCACCTCAGGCATGTCTGTCAGAGCCCTCTGGAGAACCAGCTT  
CAGGTTCTGCCTATTTTACGCTGCCTAAAGGAGCCACGAAGAAGTAAATGACGGGGTT  
GGCACTACCGTTTAGAGGAGACAGGAAAATGGAAACTAGATGGACATGACAGAAAATGAC  
TTCCAAATCCAGGTGTATCCAGTAGACAGAGCCACCGAATGCCGAAGGGCAGGCTGCG  
GAGTAGGAAGACTAGCACTGTGAGCAGGATCGTCACGTACA

The following amino acid sequence <SEQ ID NO. 262> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 128:

YVTILLTVLVFLRLSPFGIRWALSTGIHLDLEVIFCHVHLVSIPLSPLNGSANPVIYFFVGSFRQRQNRQ  
RLKLVLQRLQDMPEVKVEGGFLREPWSCREADSGSEEEPLPCQSDGTLRLAILPCHAQLHAFSCCASEMSQ  
RLKVVEM

The following DNA sequence Seq-2464 <SEQ ID NO. 129> was identified in *H. sapiens*:

TCACTGGAGAAGCCTAGTCACCTGGGCAGAATATCTTGAACCTAGGATAAGTTCATCCAT  
GGTAGACCAACTCTGTGATGGAGTTATGAGATGGGGAAGGAGGTCTGGCACCATGCAAC  
AGGATTTCCCCCAAAGCTCAGCACTCCAAGGAGCACATCAGCATCAGGAATGTCTGCTGG  
AAGCCAGCGGCTGTGGAGGAGGGCAGTAGCCACTGAGCCTAGGTTAGAGCTTCAATCC  
CCTTCAGTCCTCTGACTGGCAAGAGAACAGCAGAGTCTATTAGAGAGGAATTACCATTC  
CAAGCAAGAATTAGGCCCATCTTTTCAAGATGAGACCATTGAGTTGAGGTCCACTTAGC  
AGGGAAAGTGGCTTCAGGTTGTGGTTGACTGTTAATTACACCCTGTCTGTTCACTCTCTT  
CACCATTGTATGCAAGTACAGCATCTCTGACAAGCAAGGAACACTGGCTTGCCCCACAG  
TGGCTGGCTGGGCTTGATGAAATGAGCAACGAAGTAGCAGTGTGCCAGTCCAAGCAGAG  
ACTACCTCTAGCAGGGGCATGACATTCCCCAAGAGAGGGCATCTCCTTTAGCCTGGACCT  
TGGAGCAAAGCAACCCATGGATCAGACCAATAGACAACATGCAGCCCTCATCTA

The following amino acid sequence <SEQ ID NO. 263> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 129:

HWRSLVTWAEYLEPRISSEMVDQLCDGVMRWGRRVWHATGFPPKLSTPRSTSASGMSAGSQRLWRRGSSH  
AVQSFNPLQSSSLAREQQSLERNYHSKQEFRPHLSEHVEVHLAGKVASGCGLFNYTLFTLFTIVCKVQH  
LQARNTGLPHSGWLGLMKATKQCAQSKQRLPLAGAHSPREGISFSLDLGAKATHGSDQTTCSPHL

The following DNA sequence Seq-2465 <SEQ ID NO. 130> was identified in *H. sapiens*:

AAGAGTTAGAGCAGGATTTTACCTTGTTTTACAAAAAGAAAAGTTTATTTTGAAAAAAA  
TTCCAACCTTGCTCCTCCGAACATAGTGAAGATAATTTCCACATCCCTTTGTTC  
GGAAATGAGGACACAGTGGTGTCAATGGGTTTTGATTGTCCACTTGGAAAAGGTTAAAC  
CTGTCTACAGTCATGATGACTTCAGTTCCATTAAAGTGGGGTCTGTCTCTCACTCT  
CCACCGACTGTACCTTTACTATAACATGGCCTTATATAGATAGCTTTGAGTAAGTGTGTG  
TTAAATGACTGCCCAAGTGAATGGAAAATTGAGAAGGGCTCCAGCACTGGAGTATGGAA  
AGGAGCACTGGGTTCAATGACTCTTTGGATTCTCCCTTGCTACGTAAGTCCGTTCCCTA  
AAGGACATGGATCTTGACAGTGTGGAATCTTCAGAAATAATTGCAATACCAGAAGTTAT  
TTAAGATTTTACCATTTTCAAAGTATTTGTACGTAACACTTTCATATGTTTTGTTTCT

AGCTACCTCAGTTTCCCTGTTGGCTTGAGCAGATTAGTGTAAAGAGGTGGTGACATCAGG  
GGAAACAGGTTTACTCAGCCATCTTCATTACCATATTATCACTGACTTGAGGCTCCT

The following amino acid sequence <SEQ ID NO. 264> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 130:

GASSQYGNEDGVNLFPLMSPLYTNLLKPTGKRLGNKNIKCYVQILKWNLLVQLFLKIPTLSRSMSE  
RERTYVAREKSKESMNPVLLSILQWRPFSSIFHSLGQSFNTHLLKAIYIRPCYSKGTVGGEERQDPTMELK  
SSLDREFFPPSGQSKPNDTTVSSFPEQRDVENYLFITVRRRQGWNFQNKLFFVVKQGKILL

The following DNA sequence Seq-2466 <SEQ ID NO. 131> was identified in *H. sapiens*:

TAGTCGCTGCTTTCTGTTTCCGCTTAAAGATGGAGATATTTTTCTTTTCATGCTTGAGG  
AGTCTCGAAAGTTTTCACACTCTCCACCTCCTGGAACCTCACTGTGCCATTAGGGTG  
ACTACTGCTGTCTGGCTCCACTCGAGGGAAGCCAGGTAACCTGTGTTAGGCGCGCTTTT  
CCTGGCGGCCTTGTAATCTGTTAGTACATGAAAAGCATGACGCACATGGGGATTAGGAT  
GCCAATGCGGTGGAGTAAATCGTGTAGCCAAAGTCTTGACTGACCAAGCACACCTTATCA  
TCGTTTACATTCTGAGCCCGACCAAAATGGTAGGTAAAGTGACAAAGGCGGAAAGAAGG  
CAGACAGAAAGAATCATCTCGTCATGCATTCCCCCTCTGCCTCATAGGTTACGTGAGA  
GGCTTCATGATCCCAAGGTACCTGTGCTGATCAGTACAGGTCAAGATCCAGGCC  
GTGCAGCACATGACATTACGGAGAAGACGTTACAGAAAAAGTGCCAAAGATCCACTTG  
CCCCGATGAGGTGGTGACACTGAT

The following amino acid sequence <SEQ ID NO. 265> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 131:

ISVTDLIGGKWIFGHFFCNVFSVNVMCCTAWILTLYVISIDRYLGIMKPLTYPMRQKGKCMTKMILSVCLL  
SAFVTLPTIFGRAQNVNDKVLVSQDFGYTIYSTALASSPCASCFSCCTNRFTTRPPGKARPNTGYLASLEW  
SQTAVVTLNGTVKQFEVEECALSRLLKHERKKYLHLAETESSD

The following DNA sequence Seq-2467 <SEQ ID NO. 132> was identified in *H. sapiens*:

AGTGTACAGCTGGGCAGCCAGAGAGACAGCATGTAGTCCTCATGAAGCAGAAAGACAG  
AGGGTTCTGAGACAGAGGTCTCCAGGAAAAAAAAAAGAACCTGACTTACTGGATAAACA  
AGTCTTTAGTTTAAAAACAACAAAAAAGTGTATACATATATATATAAATCAGGTAG  
TATAAAGAAAAACAGAACTCCAGAGATTCTGGGTACAGAGGGGAAAGGGCTGTTCAA  
GAAAGTGAAATGAACTAACTGAAAATACAGCTATCTTTATATGGAAGGACAGTCAGGA  
AGTCAACAGATAAGGCCTAAACTGCATAAAGCAGGAAACAGCAGACTAAAGACATTATTA  
AGAAATATGGAACACAACCAAAAGAAATAGCAAAACAATGAAAGTGACTGTTTTTCAT  
AAGTGAGGCAGGGGAAGAGAAGGGGTATTTTTTTCCCATTATATGTCTTTAAGAACTA  
CTTGCTAAAAATATTGGGCACATATGAATTTGATAAAGCGAAAAAAGTTTACTTCACA  
AGTGCAGCTTTAACATACGTTGATTACAGTGAAGT

The following amino acid sequence <SEQ ID NO. 266> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 132:

FTVINVCSCTEVKSFSLSNSYVPNIFSKFLKTYNGEKNPFSSPASLMKNSHFSLELLLVVVFHISCL  
SAVSCFMQFRPYLLTSLSFQYKDSCTFSFNFTFLNSPFFCDPGISGVLEFFILPDFIYICVYSELLFFKL  
KTCCLSSKSGSFFFSWRPLSQNPLSFCFNEDYMLSLWLPSCNT

The following DNA sequence Seq-2468 <SEQ ID NO. 133> was identified in

*H. sapiens:*

AAAGGTGACAGAGAAGTAGGTGAGGAATTCAGTTTAAATTTATTCAATTTTAAGTTGTG  
 TCAGGTCTCCCCAAGATTATCCCTCGGTTCTGTGATTCATAGGACTTAGCATATAGTTGT  
 ATTCACAGCTATGACTTATTAACAGAGGGATACCGAAGCATAAATCAGCAAAAGGAAAAGA  
 TGCATGAGGAAAAGTCTGAAGAAACCAGGGACAGCTTCCAAGATTCTTTTCCAGTGAAA  
 TTACACAGGATATGCTTAATTCTTTCAGCAAGGAATGTGACAAGACATGTGAAACACTA  
 CCTGCCAGGGAAGTTCTTAGTGACTCAGTGCCCATGGTTATTATTGGGGACTGGTCACG  
 TATGCCCTCTTTCCTCATACTTAGAGAATTCAGTTCAGAAAGGAAAGCAGGTATTAG  
 TATAAGCCATATTATTTGCATAGACCAGTTTAGGATCAAGGAATTGTAGGAAGCTTTTCA  
 AAATCTAAGACCCCAAATACCAGCCAAGAGCCAGCTTGCAAGCAGGACATTTTAAGAGT  
 AGCAGTCTTGGGTCTGCTGTATTAACCTCTTTCTGCACAGAAATGATAGTATGACATCTA  
 AGTTATTATTATCAAGGGACCGAGAAATGCATGTTTTTAGGCTAGGGAAG

The following amino acid sequence <SEQ ID NO. 267> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 133:

FPSLKNMHFSVPLRCHTII SVQKRVNTADPRLLLLKCPACKAGSWLVFGVLD FEKLPTIPSTGLCKYGLYI  
 PAFLLELEFSKYEAKRAYVTSPQPWALSHGTS LAGSVSHVLSQFLAERIKHILCNFTGKRILEAVPGFFRL  
FLMHLFLLLI MLRYPSVNKS LIQLYAKSYESQNRGIILGRPD TTKINLKLNSSPTSLSP

The following DNA sequence Seq-74 <SEQ ID NO. 134> was identified in *H. sapiens*:

ATGAACCAGACTTTGAATAGCAGTGGGACCGTGGA  
 GTCAGCCCTAAACTATTCCAGAGGGAGCACAGTGCACACGGCTACCTGG  
 TGCTGAGCTCCCTGGCCATGTTACCTGCCTGTGCGGGATGGCAGGCAAC  
 AGCATGGTGATCTGGCTGCTGGGCTTTTGAATGCACAGGAACCCCTTCTG  
 CATCTATATCCTCAACCTGGCGGCAGCCGACCTCCTCTTCTCTTCAGCA  
 TGGCTTCCACGCTCAGCCTGGAAACCCAGCCCTGGTCAATACCACTGAC  
 AAGTCCACGAGCTGATGAAGAGACTGATGTACTTGCCTACACAGTGGG  
 CCTGAGCCTGCTGACGGCCATCAGCACCCAGCGCTGTCTCTGTCTCTCT  
 TCCCTATCTGGTTCAAGTGTACCCGGCCAGGCACCTGTGACGCTGGGTG  
 TGTGGCCTGCTGTGGACACTCTGTCTCCTGATGAACGGGTTGACCTCTTC  
 CTCTGACGCAAGTTCTTGAATTCATGAAGATCGGTGCTTCAGGGTGG  
 ACATGGTCCAGGCCGCCCTCATCATGGGGGTCTTAACCCAGTGATGACT  
 CTGTCCAGCCTGACCCTCTTTGTCTGGGTGCGGAGGAGCTCCAGCAGTG  
 GCGGCGGCAGCCACACGGCTGTTGCTGGTGGTCTGCGCTCTGTCTCTGG  
 TGTTCCTCATCTGTTCCCTGCCTCTGAGCATCTACTGGTTTGTGCTCTAC  
 TGGTTGAGCCTGCGGCCGAGATGCAGGTCTGTGCTTCAGCTTGTACAG  
 CCTCTCCTCGTCCGTAAGCAGCAGCGCCAACCCCGTCATCTACTTCTCTGG  
 TGGGACGCGGAGGAGCCACAGGCTGCCACAGGTCCCTGGGGACTGTG  
 CTCCAACAGGCGCTTCGCGAGGAGCCGAGCTGGAAGGTGGGGAGACGCC  
 CACCGTGGGCACCAATGAGATGGGGGCTTGA

The following amino acid sequence <SEQ ID NO. 268> is the predicted amino acid sequence derived from the DNA sequence of SEQ ID NO. 134:

MNQTLNSSGTVESALNYSRGSTVHTAYLVLSLAMFTCLCGMAGNSMVIWLLGFRMHRNPFCIYILNLAAA  
DLLFLFSMASTLSLETQPLVNTTDKVHLMKRLMYFAYTVGLSLLTAISTORCLSVLFPIWFKCHRPRHLS  
AWVCGLLWTLCLLMNGLTSSFCSKFLKFNEIDRCFRVDMVQAALIMGVLPVMTLSLLTLFVWVRRSSQQR  
RQPTRLFFVVVLASVLF LICSLPLSIYWFVLYWLSLPEMQVLCFSLSRLSSSVSSSANFVIYFLVGSRRS  
 HRLPTRSLGTVLQALREBPELEGGETPTVGTNEMGA

**EXAMPLE 2: CLONING OF nGPCR-x**

cDNAs may be sequenced directly using an ABI377 or ABI373A fluorescence-based sequencer (Perkin Elmer/Applied Biosystems Division, PE/ABD, Foster City, CA) and the ABI PRISM Ready Dye-Deoxy Terminator kit with Taq FS polymerase. Each ABI cycle sequencing reaction contains about 0.5µg of plasmid DNA. Cycle-sequencing is performed using an initial denaturation at 98°C for 1 min, followed by 50 cycles: 98°C for 30 sec, annealing at 50°C for 30 sec, and extension at 60°C for 4 min. Temperature cycles and times are controlled by a Perkin-Elmer 9600 thermocycler. Extension products are purified using Centriflex gel filtration (Advanced Genetic Technologies Corp., Gaithersburg, MD). Each reaction product is loaded by pipette onto the column, which is then centrifuged in a swinging bucket centrifuge (Sorvall model RT6000B table-top centrifuge) at 1500 x g for 4 min at room temperature. Column-purified samples are dried under vacuum for about 40 min and then dissolved in 5µl of a DNA loading solution (83% deionized formamide, 8.3 mM EDTA, and 1.6 mg/ml Blue Dextran). The samples are then heated to 90°C for three min and loaded into the gel sample wells for sequence analysis by the ABI377 sequencer. Sequence analysis is performed by importing ABI373A files into the Sequencer program (Gene Codes, Ann Arbor, MI). Generally, sequence reads of 700 bp are obtained. Potential sequencing errors are minimized by obtaining sequence information from both DNA strands and by re-sequencing difficult areas using primers at different locations until all sequencing ambiguities are removed.

To isolate a cDNA clone encoding full length nGPCR, a DNA fragment corresponding to a nucleotide sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134, or a portion thereof, can be used as a probe for hybridization screening of a phage cDNA library. The DNA fragment is amplified by the polymerase chain reaction (PCR) method. The PCR reaction mixture of 50µl contains polymerase mixture (0.2mM dNTPs, 1x PCR Buffer and 0.75µl Expand High Fidelity Polymerase (Roche Biochemicals)), 1µg of 3206491 plasmid, and 50pmoles of forward primer and 50pmoles of reverse primer. The primers are preferably 10 to 25 nucleotides in length and are determined by procedures well known to those skilled in the art. Amplification is performed in an Applied Biosystems PE2400 thermocycler, using the following program:



95°C for 15 seconds, 52°C for 30 seconds and 72°C for 90 seconds; repeated for 25 cycles. The amplified product is separated from the plasmid by agarose gel electrophoresis, and purified by Qiaquick gel extraction kit (Qiagen).

A lambda phage library containing cDNAs cloned into lambda ZAPII phage-vector is plated with *E. coli* XL-1 blue host, on 15 cm LB-agar plates at a density of 50,000 pfu per plate, and grown overnight at 37°C; (plated as described by Sambrook *et al.*, *supra*). Phage plaques are transferred to nylon membranes (Amersham Hybond NJ), denatured for 2 minutes in denaturation solution (0.5 M NaOH, 1.5 M NaCl), renatured for 5 minutes in renaturation solution (1 M Tris pH 7.5, 1.5 M NaCl), and washed briefly in 2xSSC (20x SSC: 3 M NaCl, 0.3 M Na-citrate). Filter membranes are dried and incubated at 80°C for 120 minutes to cross link the phage DNA to the membranes.

The membranes are hybridized with a DNA probe prepared as described above. A DNA fragment (25ng) is labeled with  $\alpha$ -<sup>32</sup>P-dCTP (NEN) using Rediprime random priming (Amersham Pharmacia Biotech), according to the manufacturer's instructions. Labeled DNA is separated from unincorporated nucleotides by S200 spin columns (Amersham Pharmacia Biotech), denatured at 95°C for 5 minutes and kept on ice. The DNA-containing membranes (above) are pre-hybridized in 50ml ExpressHyb (Clontech) solution at 68°C for 90 minutes. Subsequently, the labeled DNA probe is added to the hybridization solution, and the probe is left to hybridize to the membranes at 68°C for 70 minutes. The membranes are washed five times in 2x SSC, 0.1% SDS at 42°C for 5 minutes each, and finally washed 30 minutes in 0.1x SSC, 0.2% SDS. Filters are exposed to Kodak XAR film (Eastman Kodak Company, Rochester, N.Y., USA) with an intensifying screen at -80°C for 16 hours. One positive colony is isolated from the plates, and re-plated with about 1000 pfu on a 15 cm LB plate. Plating, plaque lift to filters and hybridization are performed as described above. About four positive phage plaques are isolated from this secondary screening.

cDNA containing plasmids (pBluescript SK-) are rescued from the isolated phages by in vivo excision by culturing XL-1 blue cells co-infected with the isolated phages and with the Excision helper phage, as described by the manufacturer (Stratagene). XL-blue cells containing the plasmids are plated on LB plates and grown at 37°C for 16 hours. Colonies (18) from each plate are replated on LB plates and grown. One colony from each

plate is stricken onto a nylon filter in an ordered array, and the filter is placed on a LB plate to raise the colonies. The filter is then hybridized with a labeled probe as described above. About three positive colonies are selected and grown up in LB medium. Plasmid DNA is isolated from the three clones by Qiagen Midi Kit (Qiagen) according to the  
5 manufacturer's instructions. The size of the insert is determined by digesting the plasmid with the restriction enzymes NotI and SalI, which establishes an insert size. The sequence of the entire insert is determined by automated sequencing on both strands of the plasmids.

### EXAMPLE 3: SUBCLONING OF THE CODING REGION OF nGPCR-X VIA 10 PCR

Additional experiments may be conducted to subclone the coding region of nGPCR and place the isolated coding region into a useful vector. Two additional PCR primers are designed based on the coding region of nGPCR, corresponding to either end. To protect against exonucleolytic attack during subsequent exposure to enzymes, *e.g.*, Taq  
15 polymerase, primers are routinely synthesized with a protective run of nucleotides at the 5' end that were not necessarily complementary to the desired target.

PCR is performed in a 50µl reaction containing 34µl H<sub>2</sub>O, 5 µl 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5µl 15mM MgSO<sub>4</sub>, 2µl dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10 mM), 3µl genomic phage DNA  
20 (0.25µg/µl), 0.3µl Primer 1 (1µg/µl), 0.3µl Primer 2 (1µg/µl), 0.4µl High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction was started with 1 cycle of 94°C for 2 minutes; followed by 25 cycles at 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1.3 minutes.

The contents from the PCR reaction are loaded onto a 2% agarose gel and  
25 fractionated. The DNA band of expected size is excised from the gel, placed in a GenElute Agarose spin column (Supelco) and spun for 10 minutes at maximum speed in a microfuge. The eluted DNA is precipitated with ethanol and resuspended in 6µl H<sub>2</sub>O for ligation.

The PCR-amplified DNA fragment containing the coding region is cloned into  
30 pCR2.1 using a protocol standard in the art. In particular, the ligation reaction consists of 6µl of GPCR DNA, 1µl 10X ligation buffer, 2µl pCR2.1 (25ng/µl, Invitrogen), and 1µl T4

DNA ligase (Invitrogen). The reaction mixture is incubated overnight at 14°C and the reaction is then stopped by heating at 65°C for 10 minutes. Two microliters of the ligation reaction are transformed into One Shot cells (Invitrogen) and plated onto ampicillin plates. A single colony containing a recombinant pCR2.1 bearing an insert is used to inoculate a 5 ml culture of LB medium. Plasmid DNA is purified using the Concert Rapid Plasmid Miniprep System (GibcoBRL) and sequenced. Following confirmation of the sequence, a 50 ml culture of LB medium is inoculated with the transformed One Shot cells, cultured, and processed using a Qiagen Plasmid Midi Kit to yield purified pCR-GPCR.

#### nGPCR-74

10 PCR was performed in a 50 µl reaction using components that come with PLATINUM<sup>®</sup> Pfx DNA Polymerase (GibcoBRL) containing 30.5 µl H<sub>2</sub>O, 5 µl 10X Pfx Amplification buffer, 5 µl 10X Enhancer solution, 1.5 µl 50mM MgSO<sub>4</sub>, 2 µl 10 mM dNTP, 5 µl human genomic DNA (0.3 µg/µl) (Clontech), 0.3 µl of LW1591 (SEQ ID NO: 3) (1 µg/µl), 0.3 µl of LW1592 (SEQ ID NO: 4) (1 µg/µl), 0.4 µl PLATINUM<sup>®</sup> Pfx DNA  
15 Polymerase (2.5 U/µl). The PCR reaction was performed in a Robocycler Gradient 96 (Stratagene) starting with 1 cycle of 94°C for 5 min followed by 30 cycles at 94°C for 30 sec, 55°C for 2 min, 68°C for 3 min. Following the final cycle, 0.5 µl of AmpliTaq DNA Polymerase (5 U/µl) was added and the tube was incubated at 72°C for 5 min. The PCR reaction was loaded onto a 1.2% agarose gel. The DNA band was excised from the gel,  
20 placed in GenElute Agarose spin column (Supelco) and spun for 10 min at maximum speed in a microcentrifuge. The eluted DNA was EtOH precipitated and resuspended in 121 H<sub>2</sub>O for ligation. The forward PCR primer sequence was:

LW1591: GATCAAGCTTGGATGAACCAGACTTTGAATAGC (SEQ ID NO: 272) and the reverse PCR primer was:

25 LW1592: GATCCTCGAGCTCAAGCCCCATCTCATTGG (SEQ ID NO: 273)

The ligation reaction used solutions from the TOPO TA Cloning Kit (Invitrogen) which consisted of 4 µl PCR product DNA and 1 µl pCRII-TOPO vector that was incubated for 5 minutes at room temperature. To the ligation reaction one microliter of 6X TOPO Cloning Stop Solution was added then the reaction was placed on ice. Two microliters of the  
30 ligation reaction was transformed in One-Shot TOP10 cells (Invitrogen), and placed on ice

for 30 minutes. The cells were heat-shocked for 30 seconds at 42°C, placed on ice for two minutes, 250 µl of SOC was added, then incubated at 37°C with shaking for one hour and then plated onto ampicillin plates. A single colony containing an insert was used to inoculate a 5 ml culture of LB medium. Plasmid DNA was purified using a Concert Rapid  
5 Plasmid Miniprep System (GibcoBRL) and then sequenced.

The DNA subcloned into pCRII-TOPO was sequenced using the ABI PRISM™ 310 Genetic Analyzer (PE Applied Biosystems) which uses advanced capillary electrophoresis technology and the ABI PRISM™ BigDye™ Terminator Cycle Sequencing Ready Reaction Kit. Each cycle-sequencing reaction contained 6 µl of H<sub>2</sub>O, 8  
10 µl of BigDye Terminator mix, 5 µl mini-prep DNA (0.1 µg/µl), and 1 µl primer (25 ng/µl) and was performed in a Perkin-Elmer 9600 thermocycler with 25 cycles of 96°C for 10 sec, 50°C for 10 sec, and 60°C for 4 min. The product was purified using a Centriflex™ gel filtration cartridge, dried under vacuum, then dissolved in 16 µl of Template Suppression Reagent (PE Applied Biosystems). The samples were heated at 95°C for 5  
15 min then placed in the 310 Genetic Analyzer.

#### **EXAMPLE 4: HYBRIDIZATION ANALYSIS TO DEMONSTRATE nGPCR-X EXPRESSION IN BRAIN**

The expression of nGPCR-x in mammals, such as the rat, may be investigated by  
20 *in situ* hybridization histochemistry. To investigate expression in the brain, for example, coronal and sagittal rat brain cryosections (20µm thick) are prepared using a Reichert-Jung cryostat. Individual sections are thaw-mounted onto silanized, nuclease-free slides (CEL Associates, Inc., Houston, TX), and stored at -80°C. Sections are processed starting with post-fixation in cold 4% paraformaldehyde, rinsed in cold phosphate-buffered saline  
25 (PBS), acetylated using acetic anhydride in triethanolamine buffer, and dehydrated through a series of alcohol washes in 70%, 95%, and 100% alcohol at room temperature. Subsequently, sections are delipidated in chloroform, followed by rehydration through successive exposure to 100% and 95% alcohol at room temperature. Microscope slides containing processed cryosections are allowed to air dry prior to hybridization. Other  
30 tissues may be assayed in a similar fashion.

A nGPCR-x-specific probe is generated using PCR. Following PCR amplification, the fragment is digested with restriction enzymes and cloned into pBluescript II cleaved with the same enzymes. For production of a probe specific for the sense strand of nGPCR-x, the nGPCR-x clone in pBluescript II is linearized with a suitable restriction enzyme, which provides a substrate for labeled run-off transcripts (*i.e.*, cRNA riboprobes) using the vector-borne T7 promoter and commercially available T7 RNA polymerase. A probe specific for the antisense strand of nGPCR-x is also readily prepared using the nGPCR-x clone in pBluescript II by cleaving the recombinant plasmid with a suitable restriction enzyme to generate a linearized substrate for the production of labeled run-off cRNA transcripts using the T3 promoter and cognate polymerase. The riboprobes are labeled with [<sup>35</sup>S]-UTP to yield a specific activity of about  $0.40 \times 10^6$  cpm/pmol for antisense riboprobes and about  $0.65 \times 10^6$  cpm/pmol for sense-strand riboprobes. Each riboprobe is subsequently denatured and added (2 pmol/ml) to hybridization buffer which contained 50% formamide, 10% dextran, 0.3 M NaCl, 10 mM Tris (pH 8.0), 1 mM EDTA, 1X Denhardt's Solution, and 10 mM dithiothreitol. Microscope slides containing sequential brain cryosections are independently exposed to 45 $\mu$ l of hybridization solution per slide and silanized cover slips are placed over the sections being exposed to hybridization solution. Sections are incubated overnight (15-18 hours) at 52°C to allow hybridization to occur. Equivalent series of cryosections are exposed to sense or antisense nGPCR-x-specific cRNA riboprobes.

Following the hybridization period, coverslips are washed off the slides in 1X SSC, followed by RNase A treatment involving the exposure of slides to 20  $\mu$ g/ml RNase A in a buffer containing 10mM Tris-HCl (pH 7.4), 0.5M EDTA, and 0.5M NaCl for 45 minutes at 37°C. The cryosections are then subjected to three high-stringency washes in 0.1 X SSC at 52°C for 20 minutes each. Following the series of washes, cryosections are dehydrated by consecutive exposure to 70%, 95%, and 100% ammonium acetate in alcohol, followed by air drying and exposure to Kodak BioMax™ MR-1 film. After 13 days of exposure, the film is developed. Based on these results, slides containing tissue that hybridized, as shown by film autoradiograms, are coated with Kodak NTB-2 nuclear track emulsion and the slides are stored in the dark for 32 days. The slides are then developed and counterstained with hematoxylin. Emulsion-coated sections are analyzed

microscopically to determine the specificity of labeling. The signal is determined to be specific if autoradiographic grains (generated by antisense probe hybridization) are clearly associated with cresyl violet-stained cell bodies. Autoradiographic grains found between cell bodies indicates non-specific binding of the probe.

5           As discussed above, it is well known that GPCRs are expressed in many different tissues and regions, including in the brain. Expression of nGPCR-x in the brain provides an indication that modulators of nGPCR-x activity have utility for treating neurological disorders, including but not limited to, mental disorder, affective disorders, ADHD/ADD (i.e., Attention Deficit-Hyperactivity Disorder/Attention Deficit Disorder), and neural  
10 disorders such as Alzheimer's disease, Parkinson's disease, migraine, and senile dementia. Some other diseases for which modulators of nGPCR-x may have utility include depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, and the like. Use of nGPCR-x modulators, including nGPCR-x ligands and anti-nGPCR-x antibodies, to treat individuals having such disease states is intended as an aspect of the  
15 invention.

#### EXAMPLE 5: TISSUE EXPRESSION PROFILING

Tissue specific expression of nGPCR-74 was detected using a PCR-based method. Tissue specific expression of cDNAs encoding nGPCR-x may be accomplished using  
20 similar methods.

A PCR-based system (RapidScan™ Gene Expression Panel, OriGene Technologies, Rockville, MD) may be used to generate a comprehensive expression profile of the putative nGPCR-x in human tissue, and in human brain regions. The RapidScan Expression Panel is comprised of first-strand cDNAs from various human  
25 tissues and brain regions that are serially diluted over a 4-log range and arrayed into a multi-well PCR plate. Human tissues in the array may include: brain, heart, kidney, spleen, liver, colon, lung, small intestine, muscle, stomach, testis, placenta, salivary gland, thyroid, adrenal gland, pancreas, ovary, uterus, prostate, skin, PBL, bone marrow, fetal brain, and fetal liver.

Expression of nGPCR-x in various tissues is detected using PCR primers designed based on the available sequence of the receptor that will prime the synthesis of a predetermined size fragment in the presence of the appropriate cDNA.

PCR is performed in a 50 $\mu$ l reaction containing 34 $\mu$ l H<sub>2</sub>O, 5 $\mu$ l 10X TT buffer (140 mM ammonium sulfate, 0.1% gelatin, 0.6 M Tris-tricine, pH 8.4), 5 $\mu$ l 15mM MgSO<sub>4</sub>, 2 $\mu$ l dNTP mixture (dGTP, dATP, dTTP, and dCTP, each at 10mM), 0.3 $\mu$ l forward primer (1 $\mu$ g/ $\mu$ l), 0.3 $\mu$ l reverse primer (1 $\mu$ g/ $\mu$ l), 0.4 $\mu$ l High Fidelity Taq polymerase (Boehringer Mannheim). The PCR reaction mixture is added to each well of the PCR plate. The plate is placed in a MJ Research PTC100 thermocycler, and is then exposed to the following cycling parameters: Pre-soak 94°C for 3 min; denaturation at 94°C for 30 seconds; annealing at primer 57°C for 45 seconds; extension 72°C for 2 minutes; for 35 cycles. PCR productions are then separated and analyzed by electrophoresis on a 1.2%-agarose gel stained with ethidium bromide.

The 4-log dilution range of cDNA deposited on the plate ensures that the amplification reaction is within the linear range and, hence, facilitates semi-quantitative determination of relative mRNA accumulation in the various tissues or brain regions examined.

Primers were synthesized by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System™ (Roche Molecular Biochemicals, Indianapolis, IN).

For nGPCR-74, the above procedure was followed. Multiple Choice™ first strand cDNAs (OriGene Technologies, Rockville, MD) from 12 human tissues were serially diluted over a 3-log range and arrayed into a multi-well PCR plate. This array was used to generate a comprehensive expression profile of the putative GPCR in human tissues. Human tissues arrayed include: brain, heart, kidney, peripheral blood leukocytes, liver, lung, muscle, ovary, prostate, small intestine, spleen and testis. The forward primer used was:

5'CTGTCTCTCTGTCTCTTCC (SEQ ID NO: 270),

and the reverse primer used was:

5'GCACCGATCTTCATTGAATTTC (SEQ ID NO: 271). This primer set primed the synthesis of a 157 base pair fragment in the presence of the appropriate cDNA. For

detection of expression within brain regions, the same primer set was used with the Human Brain Rapid Scan™ Panel (OriGene Technologies, Rockville, MD). This panel represents serial dilutions over a 2 log range of first strand cDNA from the following brain regions arrayed in a 96 well format: frontal lobe, temporal lobe, cerebellum, hippocampus, substantia nigra, caudate nucleus, amygdala, thalamus, hypothalamus, pons, medulla and spinal cord. Primers were synthesized by Genosys Corp., The Woodlands, TX. PCR reactions were assembled using the components of the Expand Hi-Fi PCR System™ (Roche Molecular Biochemicals, Indianapolis, IN). Twenty-five microliters of the PCR reaction mixture was added to each well of the RapidScan PCR plate. The plate was placed in a GeneAmp 9700 PCR thermocycler (Perkin Elmer Applied Biosystems). The following cycling program was executed: Pre-soak at (94° for 3min.) followed by 35 cycles of [(94° for 45 sec.) (53°C for 2 min.) (72° for 45 sec)]. PCR reaction products were then separated and analyzed by electrophoresis on a 2.0% agarose gel stained with ethidium bromide.

nGPCR-74 was expressed in the brain, heart, kidney, peripheral blood leukocytes, liver, lung, muscle, ovary, prostate, small intestine, spleen, and testis. Within the brain, nGPCR-74 was expressed in the frontal and temporal lobes, cerebellum, hippocampus, substantia nigra, amygdala, thalamus, pons, and spinal cord.

Expression of the nGPCR-74 in the brain provides an indication that modulators of nGPCR-74 activity have utility for treating neurological disorders, including but not limited to, schizophrenia, affective disorders, ADHD/ADD (*i.e.*, Attention Deficit-Hyperactivity Disorder/Attention Deficit Disorder), neural disorders such as Alzheimer's disease, Parkinson's disease, migraine, senile dementia, depression, anxiety, bipolar disease, epilepsy, neuritis, neurasthenia, neuropathy, neuroses, metabolic disorders, inflammatory disorders, cancers and the like. Use of nGPCR-74 modulators, including nGPCR-74 ligands and anti-nGPCR-74 antibodies, to treat individuals having such disease states is intended as an aspect of the invention.

#### EXAMPLE 6: NORTHERN BLOT ANALYSIS

Northern blots are performed to examine the expression of nGPCR-x mRNA. The sense orientation oligonucleotide and the antisense-orientation oligonucleotide, described



above, are used as primers to amplify a portion of the GPCR-x cDNA sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134.

Multiple human tissue northern blots from Clontech (Human II # 7767-1) are hybridized with the probe. Pre-hybridization is carried out at 42 C for 4 hours in 5xSSC, 1X Denhardt's reagent, 0.1% SDS, 50% formamide, 250 mg/ml salmon sperm DNA. Hybridization is performed overnight at 42°C in the same mixture with the addition of about  $1.5 \times 10^6$  cpm/ml of labeled probe.

The probe is labeled with  $\alpha$ - $^{32}$ P-dCTP by Rediprime™ DNA labeling system (Amersham Pharmacia), purified on Nick Column™ (Amersham Pharmacia) and added to the hybridization solution. The filters are washed several times at 42°C in 0.2x SSC, 0.1% SDS. Filters are exposed to Kodak XAR film (Eastman Kodak Company, Rochester, N.Y., USA) with intensifying screen at -80°C.

#### EXAMPLE 7: RECOMBINANT EXPRESSION OF nGPCR-X IN EUKARYOTIC HOST CELLS

##### A. Expression of nGPCR-x in Mammalian Cells

To produce nGPCR-x protein, a nGPCR-x-encoding polynucleotide is expressed in a suitable host cell using a suitable expression vector and standard genetic engineering techniques. For example, the nGPCR-x-encoding sequence described in Example 1 is subcloned into the commercial expression vector pzeoSV2 (Invitrogen, San Diego, CA) and transfected into Chinese Hamster Ovary (CHO) cells using the transfection reagent FuGENE6™ (Boehringer-Mannheim) and the transfection protocol provided in the product insert. Other eukaryotic cell lines, including human embryonic kidney (HEK 293) and COS cells, are suitable as well. Cells stably expressing nGPCR-x are selected by growth in the presence of 100µg/ml zeocin (Stratagene, LaJolla, CA). Optionally, nGPCR-x may be purified from the cells using standard chromatographic techniques. To facilitate purification, antisera is raised against one or more synthetic peptide sequences that correspond to portions of the nGPCR-x amino acid sequence, and the antisera is used to affinity purify nGPCR-x. The nGPCR-x also may be expressed in-frame with a tag sequence (e.g., polyhistidine, hemagglutinin, FLAG) to facilitate purification. Moreover, it

will be appreciated that many of the uses for nGPCR-x polypeptides, such as assays described below, do not require purification of nGPCR-x from the host cell.

#### B. Expression of nGPCR-x in HEK-293 cells

For expression of nGPCR-x in mammalian cells HEK293 (transformed human,  
5 primary embryonic kidney cells), a plasmid bearing the relevant nGPCR-x coding  
sequence is prepared, using vector pSecTag2A (Invitrogen). Vector pSecTag2A contains  
the murine IgK chain leader sequence for secretion, the  $\alpha$ -myc epitope for detection of the  
recombinant protein with the anti-myc antibody, a C-terminal polyhistidine for  
purification with nickel chelate chromatography, and a Zeocin resistant gene for selection  
10 of stable transfectants. The forward primer for amplification of this GPCR cDNA is  
determined by routine procedures and preferably contains a 5' extension of nucleotides to  
introduce the *HindIII* cloning site and nucleotides matching the GPCR sequence. The  
reverse primer is also determined by routine procedures and preferably contains a 5'  
extension of nucleotides to introduce an *XhoI* restriction site for cloning and nucleotides  
15 corresponding to the reverse complement of the nGPCR-x sequence. The PCR conditions  
are 55°C as the annealing temperature. The PCR product is gel purified and cloned into  
the *HindIII-XhoI* sites of the vector.

The DNA is purified using Qiagen chromatography columns and transfected into  
HEK-293 cells using DOTAP™ transfection media (Boehringer Mannheim, Indianapolis,  
20 IN). Transiently transfected cells are tested for expression after 24 hours of transfection,  
using western blots probed with anti-His and anti-nGPCR-x peptide antibodies.  
Permanently transfected cells are selected with Zeocin and propagated. Production of the  
recombinant protein is detected from both cells and media by western blots probed with  
anti-His, anti-Myc or anti-GPCR peptide antibodies.

#### 25 C. Expression of nGPCR-x in COS cells

For expression of the nGPCR-x in COS7 cells, a polynucleotide molecule having a  
sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 can be  
cloned into vector p3-CI. This vector is a pUC18-derived plasmid that contains the  
HCMV (human cytomegalovirus) promoter-intron located upstream from the bGH (bovine  
30 growth hormone) polyadenylation sequence and a multiple cloning site. In addition, the

plasmid contains the *dhfr* (dihydrofolate reductase) gene which provides selection in the presence of the drug methotrexane (MTX) for selection of stable transformants.

The forward primer is determined by routine procedures and preferably contains a 5' extension which introduces an *XbaI* restriction site for cloning, followed by nucleotides which correspond to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134. The reverse primer is also determined by routine procedures and preferably contains 5'- extension of nucleotides which introduces a *SalI* cloning site followed by nucleotides which correspond to the reverse complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134. The PCR consists of an initial denaturation step of 5 min at 95°C 30 cycles of 30 sec denaturation at 95°C, 30 sec annealing at 58°C and 30 sec extension at 72°C, followed by 5 min extension at 72°C. The PCR product is gel purified and ligated into the *XbaI* and *SalI* sites of vector p3-CI. This construct is transformed into *E. coli* cells for amplification and DNA purification. The DNA is purified with Qiagen chromatography columns and transfected into COS 7 cells using Lipofectamine™ reagent from BRL, following the manufacturer's protocols. Forty-eight and 72 hours after transfection, the media and the cells are tested for recombinant protein expression.

nGPCR-x expressed from a COS cell culture can be purified by concentrating the cell-growth media to about 10 mg of protein/ml, and purifying the protein by, for example, chromatography. Purified nGPCR-x is concentrated to 0.5 mg/ml in an Amicon concentrator fitted with a YM-10 membrane and stored at -80°C.

#### D. Expression of nGPCR-x in Insect Cells

For expression of nGPCR-x in a baculovirus system, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 can be amplified by PCR. The forward primer is determined by routine procedures and preferably contains a 5' extension which adds the *NdeI* cloning site, followed by nucleotides which correspond to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134. The reverse primer is also determined by routine procedures and preferably contains a 5' extension which introduces the *KpnI* cloning site, followed by

nucleotides which correspond to the reverse complement of a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134.

The PCR product is gel purified, digested with *NdeI* and *KpnI*, and cloned into the corresponding sites of vector pACHTL-A (Pharmingen, San Diego, CA). The pACHTL expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV), and a 6XHis tag upstream from the multiple cloning site. A protein kinase site for phosphorylation and a thrombin site for excision of the recombinant protein precede the multiple cloning site is also present. Of course, many other baculovirus vectors could be used in place of pACHTL-A, such as pAc373, pVL941 and pAcIM1. Other suitable vectors for the expression of GPCR polypeptides can be used, provided that the vector construct includes appropriately located signals for transcription, translation, and trafficking, such as an in-frame AUG and a signal peptide, as required. Such vectors are described in Luckow *et al.*, Virology 170:31-39, among others.

The virus is grown and isolated using standard baculovirus expression methods, such as those described in Summers *et al.* (A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures, Texas Agricultural Experimental Station Bulletin No. 1555 (1987)).

In a preferred embodiment, pACHTL-A containing nGPCR-x gene is introduced into baculovirus using the "BaculoGold™" transfection kit (Pharmingen, San Diego, CA) using methods established by the manufacturer. Individual virus isolates are analyzed for protein production by radiolabeling infected cells with <sup>35</sup>S-methionine at 24 hours post infection. Infected cells are harvested at 48 hours post infection, and the labeled proteins are visualized by SDS-PAGE. Viruses exhibiting high expression levels can be isolated and used for scaled up expression.

For expression of a nGPCR-x polypeptide in a Sf9 cells, a polynucleotide molecule having a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 can be amplified by PCR using the primers and methods described above for baculovirus expression. The nGPCR-x cDNA is cloned into vector pACHTL-A (Pharmingen) for expression in Sf9 insect. The insert is cloned into the *NdeI* and *KpnI* sites, after elimination of an internal *NdeI* site (using the same primers described above for

expression in baculovirus). DNA is purified with Qiagen chromatography columns and expressed in Sf9 cells. Preliminary Western blot experiments from non-purified plaques are tested for the presence of the recombinant protein of the expected size which reacted with the GPCR-specific antibody. These results are confirmed after further purification and expression optimization in HiG5 cells.

#### EXAMPLE 8: INTERACTION TRAP/TWO-HYBRID SYSTEM

In order to assay for nGPCR-x-interacting proteins, the interaction trap/two-hybrid library screening method can be used. This assay was first described in Fields *et al.*, *Nature*, 1989, 340, 245, which is incorporated herein by reference in its entirety. A protocol is published in Current Protocols in Molecular Biology 1999, John Wiley & Sons, NY, and Ausubel, F. M. *et al.* 1992, Short protocols in molecular biology, Fourth edition, Greene and Wiley-interscience, NY, each of which is incorporated herein by reference in its entirety. Kits are available from Clontech, Palo Alto, CA (Matchmaker Two-Hybrid System 3).

A fusion of the nucleotide sequences encoding all or partial nGPCR-x and the yeast transcription factor GAL4 DNA-binding domain (DNA-BD) is constructed in an appropriate plasmid (*i.e.*, pGBKT7) using standard subcloning techniques. Similarly, a GAL4 active domain (AD) fusion library is constructed in a second plasmid (*i.e.*, pGADT7) from cDNA of potential GPCR-binding proteins (for protocols on forming cDNA libraries, see Sambrook *et al.* 1989, Molecular cloning: a laboratory manual, second edition, Cold Spring Harbor Press, Cold Spring Harbor, NY), which is incorporated herein by reference in its entirety. The DNA-BD/nGPCR-x fusion construct is verified by sequencing, and tested for autonomous reporter gene activation and cell toxicity, both of which would prevent a successful two-hybrid analysis. Similar controls are performed with the AD/library fusion construct to ensure expression in host cells and lack of transcriptional activity. Yeast cells are transformed (*ca.* 10<sup>5</sup> transformants/mg DNA) with both the nGPCR-x and library fusion plasmids according to standard procedures (Ausubel *et al.*, 1992, Short protocols in molecular biology, fourth edition, Greene and Wiley-interscience, NY, which is incorporated herein by reference in its entirety). *In vivo* binding of DNA-BD/nGPCR-x with AD/library proteins results in

transcription of specific yeast plasmid reporter genes (*i.e.*, lacZ, HIS3, ADE2, LEU2). Yeast cells are plated on nutrient-deficient media to screen for expression of reporter genes. Colonies are dually assayed for  $\beta$ -galactosidase activity upon growth in Xgal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside) supplemented media (filter assay for  $\beta$ -galactosidase activity is described in Breeden *et al.*, Cold Spring Harb. Symp. Quant. Biol., 1985, 50, 643, which is incorporated herein by reference in its entirety). Positive AD-library plasmids are rescued from transformants and reintroduced into the original yeast strain as well as other strains containing unrelated DNA-BD fusion proteins to confirm specific nGPCR-x/library protein interactions. Insert DNA is sequenced to verify the presence of an open reading frame fused to GAL4 AD and to determine the identity of the nGPCR-x-binding protein.

#### EXAMPLE 9: MOBILITY SHIFT DNA-BINDING ASSAY USING GEL ELECTROPHORESIS

A gel electrophoresis mobility shift assay can rapidly detect specific protein-DNA interactions. Protocols are widely available in such manuals as Sambrook *et al.* 1989, *Molecular cloning: a laboratory manual*, second edition, Cold Spring Harbor Press, Cold Spring Harbor, NY and Ausubel, F. M. *et al.*, 1992, *Short Protocols in Molecular Biology*, fourth edition, Greene and Wiley-interscience, NY, each of which is incorporated herein by reference in its entirety.

Probe DNA(<300 bp) is obtained from synthetic oligonucleotides, restriction endonuclease fragments, or PCR fragments and end-labeled with  $^{32}\text{P}$ . An aliquot of purified nGPCR-x (*ca.* 15  $\mu\text{g}$ ) or crude nGPCR-x extract (*ca.* 15 ng) is incubated at constant temperature (in the range 22-37 C) for at least 30 minutes in 10-15  $\mu\text{l}$  of buffer (*i.e.* TAE or TBE, pH 8.0-8.5) containing radiolabeled probe DNA, nonspecific carrier DNA (*ca.* 1  $\mu\text{g}$ ), BSA (300  $\mu\text{g}/\text{ml}$ ), and 10% (v/v) glycerol. The reaction mixture is then loaded onto a polyacrylamide gel and run at 30-35 mA until good separation of free probe DNA from protein-DNA complexes occurs. The gel is then dried and bands corresponding to free DNA and protein-DNA complexes are detected by autoradiography.

**EXAMPLE 10: ANTIBODIES TO nGPCR-X**

Standard techniques are employed to generate polyclonal or monoclonal antibodies to the nGPCR-x receptor, and to generate useful antigen-binding fragments thereof or variants thereof, including "humanized" variants. Such protocols can be found, for example, in Sambrook *et al.* (1989) and Harlow *et al.* (Eds.), Antibodies A Laboratory Manual; Cold Spring Harbor Laboratory; Cold Spring Harbor, NY (1988). In one embodiment, recombinant nGPCR-x polypeptides (or cells or cell membranes containing such polypeptides) are used as antigen to generate the antibodies. In another embodiment, one or more peptides having amino acid sequences corresponding to an immunogenic portion of nGPCR-x (e.g., 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more amino acids) are used as antigen. Peptides corresponding to extracellular portions of nGPCR-x, especially hydrophilic extracellular portions, are preferred. The antigen may be mixed with an adjuvant or linked to a hapten to increase antibody production.

**A. Polyclonal or Monoclonal antibodies**

As one exemplary protocol, recombinant nGPCR-x or a synthetic fragment thereof is used to immunize a mouse for generation of monoclonal antibodies (or larger mammal, such as a rabbit, for polyclonal antibodies). To increase antigenicity, peptides are conjugated to Keyhole Limpet Hemocyanin (Pierce), according to the manufacturer's recommendations. For an initial injection, the antigen is emulsified with Freund's Complete Adjuvant and injected subcutaneously. At intervals of two to three weeks, additional aliquots of nGPCR-x antigen are emulsified with Freund's Incomplete Adjuvant and injected subcutaneously. Prior to the final booster injection, a serum sample is taken from the immunized mice and assayed by western blot to confirm the presence of antibodies that immunoreact with nGPCR-x. Serum from the immunized animals may be used as polyclonal antisera or used to isolate polyclonal antibodies that recognize nGPCR-x. Alternatively, the mice are sacrificed and their spleen removed for generation of monoclonal antibodies.

To generate monoclonal antibodies, the spleens are placed in 10 ml serum-free RPMI 1640, and single cell suspensions are formed by grinding the spleens in serum-free RPMI 1640, supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate, 100 units/ml penicillin, and 100 µg/ml streptomycin (RPMI) (Gibco, Canada). The cell suspensions are

filtered and washed by centrifugation and resuspended in serum-free RPMI. Thymocytes taken from three naive Balb/c mice are prepared in a similar manner and used as a Feeder Layer. NS-1 myeloma cells, kept in log phase in RPMI with 10% fetal bovine serum (FBS) (Hyclone Laboratories, Inc., Logan, Utah) for three days prior to fusion, are  
5 centrifuged and washed as well.

To produce hybridoma fusions, spleen cells from the immunized mice are combined with NS-1 cells and centrifuged, and the supernatant is aspirated. The cell pellet is dislodged by tapping the tube, and 2 ml of 37°C PEG 1500 (50% in 75 mM HEPES, pH 8.0) (Boehringer-Mannheim) is stirred into the pellet, followed by the  
10 addition of serum-free RPMI. Thereafter, the cells are centrifuged, resuspended in RPMI containing 15% FBS, 100  $\mu$ M sodium hypoxanthine, 0.4  $\mu$ M aminopterin, 16  $\mu$ M thymidine (HAT) (Gibco), 25 units/ml IL-6 (Boehringer-Mannheim) and  $1.5 \times 10^6$  thymocytes/ml, and plated into 10 Corning flat-bottom 96-well tissue culture plates (Corning, Corning New York).

15 On days 2, 4, and 6 after the fusion, 100 $\mu$ l of medium is removed from the wells of the fusion plates and replaced with fresh medium. On day 8, the fusions are screened by ELISA, testing for the presence of mouse IgG that binds to nGPCR-x. Selected fusion wells are further cloned by dilution until monoclonal cultures producing anti-nGPCR-x antibodies are obtained.

20 B. Humanization of anti-nGPCR-x monoclonal antibodies

The expression pattern of nGPCR-x as reported herein and the proven track record of GPCRs as targets for therapeutic intervention suggest therapeutic indications for nGPCR-x inhibitors (antagonists). nGPCR-x-neutralizing antibodies comprise one class of therapeutics useful as nGPCR-x antagonists. Following are protocols to improve the  
25 utility of anti-nGPCR-x monoclonal antibodies as therapeutics in humans by "humanizing" the monoclonal antibodies to improve their serum half-life and render them less immunogenic in human hosts (*i.e.*, to prevent human antibody response to non-human anti-nGPCR-x antibodies).

The principles of humanization have been described in the literature and are  
30 facilitated by the modular arrangement of antibody proteins. To minimize the possibility of binding complement, a humanized antibody of the IgG4 isotype is preferred.



For example, a level of humanization is achieved by generating chimeric antibodies comprising the variable domains of non-human antibody proteins of interest with the constant domains of human antibody molecules. (See, *e.g.*, Morrison *et al.*, Adv. Immunol., 44:65-92 (1989)). The variable domains of nGPCR-x-neutralizing anti-nGPCR-x antibodies are cloned from the genomic DNA of a B-cell hybridoma or from cDNA generated from mRNA isolated from the hybridoma of interest. The V region gene fragments are linked to exons encoding human antibody constant domains, and the resultant construct is expressed in suitable mammalian host cells (*e.g.*, myeloma or CHO cells).

To achieve an even greater level of humanization, only those portions of the variable region gene fragments that encode antigen-binding complementarity determining regions ("CDR") of the non-human monoclonal antibody genes are cloned into human antibody sequences. (See, *e.g.*, Jones *et al.*, Nature 321:522-525 (1986); Riechmann *et al.*, Nature 332:323-327 (1988); Verhoeven *et al.*, Science 239:1534-36 (1988); and Tempest *et al.*, Bio/Technology 9: 266-71 (1991)). If necessary, the  $\beta$ -sheet framework of the human antibody surrounding the CDR3 regions also is modified to more closely mirror the three dimensional structure of the antigen-binding domain of the original monoclonal antibody. (See Kettleborough *et al.*, Protein Engin., 4:773-783 (1991); and Foote *et al.*, J. Mol. Biol., 224:487-499 (1992)).

In an alternative approach, the surface of a non-human monoclonal antibody of interest is humanized by altering selected surface residues of the non-human antibody, *e.g.*, by site-directed mutagenesis, while retaining all of the interior and contacting residues of the non-human antibody. See Padlan, Molecular Immunol., 28(4/5):489-98 (1991).

The foregoing approaches are employed using nGPCR-x-neutralizing anti-nGPCR-x monoclonal antibodies and the hybridomas that produce them to generate humanized nGPCR-x-neutralizing antibodies useful as therapeutics to treat or palliate conditions wherein nGPCR-x expression or ligand-mediated nGPCR-x signaling is detrimental.

#### C. Human nGPCR-x-Neutralizing Antibodies from Phage Display

Human nGPCR-x-neutralizing antibodies are generated by phage display techniques such as those described in Aujame *et al.*, Human Antibodies 8(4):155-168

(1997); Hoogenboom, TIBTECH 15:62-70 (1997); and Rader *et al.*, Curr. Opin. Biotechnol. 8:503-508 (1997), all of which are incorporated by reference. For example, antibody variable regions in the form of Fab fragments or linked single chain Fv fragments are fused to the amino terminus of filamentous phage minor coat protein pIII. Expression of the fusion protein and incorporation thereof into the mature phage coat results in phage particles that present an antibody on their surface and contain the genetic material encoding the antibody. A phage library comprising such constructs is expressed in bacteria, and the library is screened for nGPCR-x-specific phage-antibodies using labeled or immobilized nGPCR-x as antigen-probe.

10       D.     Human nGPCR-x-neutralizing antibodies from transgenic mice

Human nGPCR-x-neutralizing antibodies are generated in transgenic mice essentially as described in Bruggemann *et al.*, Immunol. Today 17(8):391-97 (1996) and Bruggemann *et al.*, Curr. Opin. Biotechnol. 8:455-58 (1997). Transgenic mice carrying human V-gene segments in germline configuration and that express these transgenes in their lymphoid tissue are immunized with a nGPCR-x composition using conventional immunization protocols. Hybridomas are generated using B cells from the immunized mice using conventional protocols and screened to identify hybridomas secreting anti-nGPCR-x human antibodies (*e.g.*, as described above).

20     **EXAMPLE 11: ASSAYS TO IDENTIFY MODULATORS OF nGPCR-X ACTIVITY**

Set forth below are several nonlimiting assays for identifying modulators (agonists and antagonists) of nGPCR-x activity. Among the modulators that can be identified by these assays are natural ligand compounds of the receptor; synthetic analogs and derivatives of natural ligands; antibodies, antibody fragments, and/or antibody-like compounds derived from natural antibodies or from antibody-like combinatorial libraries; and/or synthetic compounds identified by high-throughput screening of libraries; and the like. All modulators that bind nGPCR-x are useful for identifying nGPCR-x in tissue samples (*e.g.*, for diagnostic purposes, pathological purposes, and the like). Agonist and antagonist modulators are useful for up-regulating and down-regulating nGPCR-x activity, respectively, to treat disease states characterized by abnormal levels of nGPCR-x activity.

The assays may be performed using single putative modulators, and/or may be performed using a known agonist in combination with candidate antagonists (or visa versa).

A. cAMP Assays

In one type of assay, levels of cyclic adenosine monophosphate (cAMP) are measured in nGPCR-x-transfected cells that have been exposed to candidate modulator compounds. Protocols for cAMP assays have been described in the literature. (See, *e.g.*, Sutherland *et al.*, *Circulation* 37: 279 (1968); Frandsen *et al.*, *Life Sciences* 18: 529-541 (1976); Dooley *et al.*, *Journal of Pharmacology and Experimental Therapeutics* 283 (2): 735-41 (1997); and George *et al.*, *Journal of Biomolecular Screening* 2 (4): 235-40 (1997)). An exemplary protocol for such an assay, using an Adenylyl Cyclase Activation FlashPlate® Assay from NEN™ Life Science Products, is set forth below.

Briefly, the nGPCR-x coding sequence (*e.g.*, a cDNA or intronless genomic DNA) is subcloned into a commercial expression vector, such as pzeoSV2 (Invitrogen), and transiently transfected into Chinese Hamster Ovary (CHO) cells using known methods, such as the transfection protocol provided by Boehringer-Mannheim when supplying the FuGENE 6 transfection reagent. Transfected CHO cells are seeded into 96-well microplates from the FlashPlate® assay kit, which are coated with solid scintillant to which antisera to cAMP has been bound. For a control, some wells are seeded with wild type (untransfected) CHO cells. Other wells in the plate receive various amounts of a cAMP standard solution for use in creating a standard curve.

One or more test compounds (*i.e.*, candidate modulators) are added to the cells in each well, with water and/or compound-free medium/diluent serving as a control or controls. After treatment, cAMP is allowed to accumulate in the cells for exactly 15 minutes at room temperature. The assay is terminated by the addition of lysis buffer containing [<sup>125</sup>I]-labeled cAMP, and the plate is counted using a Packard Topcount™ 96-well microplate scintillation counter. Unlabeled cAMP from the lysed cells (or from standards) and fixed amounts of [<sup>125</sup>I]-cAMP compete for antibody bound to the plate. A standard curve is constructed, and cAMP values for the unknowns are obtained by interpolation. Changes in intracellular cAMP levels of cells in response to exposure to a test compound are indicative of nGPCR-x modulating activity. Modulators that act as agonists of receptors which couple to the G<sub>s</sub> subtype of G proteins will stimulate

production of cAMP, leading to a measurable 3-10 fold increase in cAMP levels. Agonists of receptors which couple to the  $G_{i/o}$  subtype of G proteins will inhibit forskolin-stimulated cAMP production, leading to a measurable decrease in cAMP levels of 50-100%. Modulators that act as inverse agonists will reverse these effects at receptors  
5 that are either constitutively active or activated by known agonists.

#### B. Aequorin Assays

In another assay, cells (e.g., CHO cells) are transiently co-transfected with both a nGPCR-x expression construct and a construct that encodes the photoprotein apoaequorin. In the presence of the cofactor coelenterazine, apoaequorin will emit a measurable  
10 luminescence that is proportional to the amount of intracellular (cytoplasmic) free calcium. (See generally, Cobbold, *et al.* "Aequorin measurements of cytoplasmic free calcium," *In*: McCormack J.G. and Cobbold P.H., eds., *Cellular Calcium: A Practical Approach*. Oxford:IRL Press (1991); Stables *et al.*, *Analytical Biochemistry* 252: 115-26 (1997); and Haugland, *Handbook of Fluorescent Probes and Research Chemicals*. Sixth  
15 edition. Eugene OR: Molecular Probes (1996).)

In one exemplary assay, nGPCR-x is subcloned into the commercial expression vector pzeoSV2 (Invitrogen) and transiently co-transfected along with a construct that encodes the photoprotein apoaequorin (Molecular Probes, Eugene, OR) into CHO cells using the transfection reagent FuGENE 6 (Boehringer-Mannheim) and the transfection  
20 protocol provided in the product insert.

The cells are cultured for 24 hours at 37°C in MEM (Gibco/BRL, Gaithersburg, MD) supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10 µg/ml streptomycin, at which time the medium is changed to serum-free MEM containing 5 µM coelenterazine (Molecular Probes, Eugene, OR). Culturing is then  
25 continued for two additional hours at 37°C. Subsequently, cells are detached from the plate using VERSEN (Gibco/BRL), washed, and resuspended at 200,000 cells/ml in serum-free MEM.

Dilutions of candidate nGPCR-x modulator compounds are prepared in serum-free MEM and dispensed into wells of an opaque 96-well assay plate at 50 µl/well. Plates are  
30 then loaded onto an MLX microtiter plate luminometer (Dynex Technologies, Inc., Chantilly, VA). The instrument is programmed to dispense 50µl cell suspensions into

each well, one well at a time, and immediately read luminescence for 15 seconds. Dose-response curves for the candidate modulators are constructed using the area under the curve for each light signal peak. Data are analyzed with SlideWrite, using the equation for a one-site ligand, and  $EC_{50}$  values are obtained. Changes in luminescence caused by the compounds are considered indicative of modulatory activity. Modulators that act as agonists at receptors which couple to the  $G_q$  subtype of G proteins give an increase in luminescence of up to 100 fold. Modulators that act as inverse agonists will reverse this effect at receptors that are either constitutively active or activated by known agonists.

### C. Luciferase Reporter Gene Assay

The photoprotein luciferase provides another useful tool for assaying for modulators of nGPCR-x activity. Cells (*e.g.*, CHO cells or COS 7 cells) are transiently co-transfected with both a nGPCR-x expression construct (*e.g.*, nGPCR-x in pzeoSV2) and a reporter construct which includes a gene for the luciferase protein downstream from a transcription factor binding site, such as the cAMP-response element (CRE), AP-1, or NF-kappa B. Agonist binding to receptors coupled to the  $G_s$  subtype of G proteins leads to increases in cAMP, thereby activating the CRE transcription factor and resulting in expression of the luciferase gene. Agonist binding to receptors coupled to the  $G_q$  subtype of G protein leads to production of diacylglycerol that activates protein kinase C, which activates the AP-1 or NF-kappa B transcription factors, in turn resulting in expression of the luciferase gene. Expression levels of luciferase reflect the activation status of the signaling events. (See generally, George *et al.*, Journal of Biomolecular Screening 2(4): 235-240 (1997); and Stratowa *et al.*, Current Opinion in Biotechnology 6: 574-581 (1995)). Luciferase activity may be quantitatively measured using, *e.g.*, luciferase assay reagents that are commercially available from Promega (Madison, WI).

In one exemplary assay, CHO cells are plated in 24-well culture dishes at a density of 100,000 cells/well one day prior to transfection and cultured at 37°C in MEM (Gibco/BRL) supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10 µg/ml streptomycin. Cells are transiently co-transfected with both a nGPCR-x expression construct and a reporter construct containing the luciferase gene. The reporter plasmids CRE-luciferase, AP-1-luciferase and NF-kappaB-luciferase may be purchased from Stratagene (LaJolla, CA). Transfections are performed using the FuGENE

6 transfection reagent (Boehringer-Mannheim) according to the supplier's instructions. Cells transfected with the reporter construct alone are used as a control. Twenty-four hours after transfection, cells are washed once with PBS pre-warmed to 37°C. Serum-free MEM is then added to the cells either alone (control) or with one or more candidate  
5 modulators and the cells are incubated at 37°C for five hours. Thereafter, cells are washed once with ice-cold PBS and lysed by the addition of 100 µl of lysis buffer per well from the luciferase assay kit supplied by Promega. After incubation for 15 minutes at room temperature, 15 µl of the lysate is mixed with 50 µl of substrate solution (Promega) in an opaque-white, 96-well plate, and the luminescence is read immediately on a Wallace  
10 model 1450 MicroBeta scintillation and luminescence counter (Wallace Instruments, Gaithersburg, MD).

Differences in luminescence in the presence versus the absence of a candidate modulator compound are indicative of modulatory activity. Receptors that are either constitutively active or activated by agonists typically give a 3 to 20-fold stimulation of  
15 luminescence compared to cells transfected with the reporter gene alone. Modulators that act as inverse agonists will reverse this effect.

D. Intracellular calcium measurement using FLIPR

Changes in intracellular calcium levels are another recognized indicator of G protein-coupled receptor activity, and such assays can be employed to screen for  
20 modulators of nGPCR-x activity. For example, CHO cells stably transfected with a nGPCR-x expression vector are plated at a density of  $4 \times 10^4$  cells/well in Packard black-walled, 96-well plates specially designed to discriminate fluorescence signals emanating from the various wells on the plate. The cells are incubated for 60 minutes at 37°C in modified Dulbecco's PBS (D-PBS) containing 36 mg/L pyruvate and 1 g/L glucose with  
25 the addition of 1% fetal bovine serum and one of four calcium indicator dyes (Fluo-3™ AM, Fluo-4™ AM, Calcium Green™-1 AM, or Oregon Green™ 488 BAPTA-1 AM), each at a concentration of 4 µM. Plates are washed once with modified D-PBS without 1% fetal bovine serum and incubated for 10 minutes at 37°C to remove residual dye from the cellular membrane. In addition, a series of washes with modified D-PBS without 1%  
30 fetal bovine serum is performed immediately prior to activation of the calcium response.

A calcium response is initiated by the addition of one or more candidate receptor agonist compounds, calcium ionophore A23187 (10  $\mu$ M; positive control), or ATP (4  $\mu$ M; positive control). Fluorescence is measured by Molecular Device's FLIPR with an argon laser (excitation at 488 nm). (See, *e.g.*, Kuntzweiler *et al.*, Drug Development Research, 44(1):14-20 (1998)). The F-stop for the detector camera was set at 2.5 and the length of exposure was 0.4 milliseconds. Basal fluorescence of cells was measured for 20 seconds prior to addition of candidate agonist, ATP, or A23187, and the basal fluorescence level was subtracted from the response signal. The calcium signal is measured for approximately 200 seconds, taking readings every two seconds. Calcium ionophore A23187 and ATP increase the calcium signal 200% above baseline levels. In general, activated GPCRs increase the calcium signal approximately 10-15% above baseline signal.

#### E. Mitogenesis Assay

In a mitogenesis assay, the ability of candidate modulators to induce or inhibit nGPCR-x-mediated cell division is determined. (See, *e.g.*, Lajiness *et al.*, Journal of Pharmacology and Experimental Therapeutics 267(3): 1573-1581 (1993)). For example, CHO cells stably expressing nGPCR-x are seeded into 96-well plates at a density of 5000 cells/well and grown at 37°C in MEM with 10% fetal calf serum for 48 hours, at which time the cells are rinsed twice with serum-free MEM. After rinsing, 80 $\mu$ l of fresh MEM, or MEM containing a known mitogen, is added along with 20 $\mu$ l MEM containing varying concentrations of one or more candidate modulators or test compounds diluted in serum-free medium. As controls, some wells on each plate receive serum-free medium alone, and some receive medium containing 10% fetal bovine serum. Untransfected cells or cells transfected with vector alone also may serve as controls.

After culture for 16-18 hours, 1 $\mu$  Ci of [ $^3$ H]-thymidine (2 Ci/mmol) is added to the wells and cells are incubated for an additional 2 hours at 37°C. The cells are trypsinized and collected on filter mats with a cell harvester (Tomtec); the filters are then counted in a Betaplate counter. The incorporation of [ $^3$ H]-thymidine in serum-free test wells is compared to the results achieved in cells stimulated with serum (positive control). Use of multiple concentrations of test compounds permits creation and analysis of dose-response curves using the non-linear, least squares fit equation:  $A = B \times [C / (D + C)] + G$  where A is the percent of serum stimulation; B is the maximal effect minus baseline; C is the EC<sub>50</sub>;

D is the concentration of the compound; and G is the maximal effect. Parameters B,C and G are determined by Simplex optimization.

Agonists that bind to the receptor are expected to increase [ $^3\text{H}$ ]-thymidine incorporation into cells, showing up to 80% of the response to serum. Antagonists that  
5 bind to the receptor will inhibit the stimulation seen with a known agonist by up to 100%.

F. [ $^{35}\text{S}$ ]GTP $\gamma$ S Binding Assay

Because G protein-coupled receptors signal through intracellular G proteins whose activity involves GTP binding and hydrolysis to yield bound GDP, measurement of binding of the non-hydrolyzable GTP analog [ $^{35}\text{S}$ ]GTP $\gamma$ S in the presence and absence of  
10 candidate modulators provides another assay for modulator activity. (See, e.g., Kowal *et al.*, *Neuropharmacology* 37:179-187 (1998).)

In one exemplary assay, cells stably transfected with a nGPCR-x expression vector are grown in 10 cm tissue culture dishes to subconfluence, rinsed once with 5 ml of ice-cold  $\text{Ca}^{2+}/\text{Mg}^{2+}$ -free phosphate-buffered saline, and scraped into 5 ml of the same buffer.  
15 Cells are pelleted by centrifugation (500 x g, 5 minutes), resuspended in TEE buffer (25 mM Tris, pH 7.5, 5 mM EDTA, 5 mM EGTA), and frozen in liquid nitrogen. After thawing, the cells are homogenized using a Dounce homogenizer (one ml TEE per plate of cells), and centrifuged at 1,000 x g for 5 minutes to remove nuclei and unbroken cells.

The homogenate supernatant is centrifuged at 20,000 x g for 20 minutes to isolate  
20 the membrane fraction, and the membrane pellet is washed once with TEE and resuspended in binding buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 10 mM  $\text{MgCl}_2$ , 1 mM EDTA). The resuspended membranes can be frozen in liquid nitrogen and stored at -70°C until use.

Aliquots of cell membranes prepared as described above and stored at -70°C are  
25 thawed, homogenized, and diluted into buffer containing 20 mM HEPES, 10 mM  $\text{MgCl}_2$ , 1 mM EDTA, 120 mM NaCl, 10  $\mu\text{M}$  GDP, and 0.2 mM ascorbate, at a concentration of 10-50  $\mu\text{g}/\text{ml}$ . In a final volume of 90  $\mu\text{l}$ , homogenates are incubated with varying concentrations of candidate modulator compounds or 100  $\mu\text{M}$  GTP for 30 minutes at 30°C and then placed on ice. To each sample, 10  $\mu\text{l}$  guanosine 5'-O-(3[ $^{35}\text{S}$ ]thio) triphosphate  
30 (NEN, 1200 Ci/mmol; [ $^{35}\text{S}$ ]-GTP $\gamma$ S), was added to a final concentration of 100-200 pM.



Samples are incubated at 30°C for an additional 30 minutes, 1 ml of 10mM HEPES, pH 7.4, 10 mM MgCl<sub>2</sub>, at 4°C is added and the reaction is stopped by filtration.

Samples are filtered over Whatman GF/B filters and the filters are washed with 20 ml ice-cold 10 mM HEPES, pH 7.4, 10 mM MgCl<sub>2</sub>. Filters are counted by liquid  
5 scintillation spectroscopy. Nonspecific binding of [<sup>35</sup>S]-GTPγS is measured in the presence of 100 μM GTP and subtracted from the total. Compounds are selected that modulate the amount of [<sup>35</sup>S]-GTPγS binding in the cells, compared to untransfected control cells. Activation of receptors by agonists gives up to a five-fold increase in [<sup>35</sup>S]GTPγS binding. This response is blocked by antagonists.

10 G. MAP Kinase Activity Assay

Evaluation of MAP kinase activity in cells expressing a GPCR provides another assay to identify modulators of GPCR activity. (See, e.g., Lajiness *et al.*, *Journal of Pharmacology and Experimental Therapeutics* 267(3):1573-1581 (1993) and Boulton *et al.*, *Cell* 65:663-675 (1991).)

15 In one embodiment, CHO cells stably transfected with nGPCR-x are seeded into 6-well plates at a density of 70,000 cells/well 48 hours prior to the assay. During this 48-hour period, the cells are cultured at 37°C in MEM medium supplemented with 10% fetal bovine serum, 2mM glutamine, 10 U/ml penicillin and 10μg/ml streptomycin. The cells are serum-starved for 1-2 hours prior to the addition of stimulants.

20 For the assay, the cells are treated with medium alone or medium containing either a candidate agonist or 200 nM Phorbol ester- myristoyl acetate (*i.e.*, PMA, a positive control), and the cells are incubated at 37°C for varying times. To stop the reaction, the plates are placed on ice, the medium is aspirated, and the cells are rinsed with 1 ml of ice-cold PBS containing 1mM EDTA. Thereafter, 200μl of cell lysis buffer (12.5 mM MOPS,  
25 pH 7.3, 12.5 mM glycerophosphate, 7.5mM MgCl<sub>2</sub>, 0.5mM EGTA, 0.5 mM sodium vanadate, 1mM benzamidine, 1mM dithiothreitol, 10 μg/ml leupeptin, 10 μg/ml aprotinin, 2μg/ml pepstatin A, and 1μM okadaic acid) is added to the cells. The cells are scraped from the plates and homogenized by 10 passages through a 23 3/4 G needle, and the cytosol fraction is prepared by centrifugation at 20,000 x g for 15 minutes.

Aliquots (5-10  $\mu$ l containing 1-5  $\mu$ g protein) of cytosol are mixed with 1 mM MAPK Substrate Peptide (APRTPGGRR (SEQ ID NO: 269), Upstate Biotechnology, Inc., N.Y.) and 50 $\mu$ M [ $\gamma$ - $^{32}$ P]ATP (NEN, 3000 Ci/mmol), diluted to a final specific activity of ~2000 cpm/pmol, in a total volume of 25  $\mu$ l. The samples are incubated for 5 minutes at 30°C, and reactions are stopped by spotting 20  $\mu$ l on 2 cm<sup>2</sup> squares of Whatman P81 phosphocellulose paper. The filter squares are washed in 4 changes of 1% H<sub>3</sub>PO<sub>4</sub>, and the squares are subjected to liquid scintillation spectroscopy to quantitate bound label. Equivalent cytosolic extracts are incubated without MAPK substrate peptide, and the bound label from these samples are subtracted from the matched samples with the substrate peptide. The cytosolic extract from each well is used as a separate point. Protein concentrations are determined by a dye binding protein assay (Bio-Rad Laboratories). Agonist activation of the receptor is expected to result in up to a five-fold increase in MAPK enzyme activity. This increase is blocked by antagonists.

#### H. [ $^3$ H]Arachidonic Acid Release

The activation of GPCRs also has been observed to potentiate arachidonic acid release in cells, providing yet another useful assay for modulators of GPCR activity. (See, e.g., Kanterman *et al.*, Molecular Pharmacology 39:364-369 (1991).) For example, CHO cells that are stably transfected with a nGPCR-x expression vector are plated in 24-well plates at a density of 15,000 cells/well and grown in MEM medium supplemented with 10% fetal bovine serum, 2 mM glutamine, 10 U/ml penicillin and 10  $\mu$ g/ml streptomycin for 48 hours at 37°C before use. Cells of each well are labeled by incubation with [ $^3$ H]-arachidonic acid (Amersham Corp., 210 Ci/mmol) at 0.5  $\mu$ Ci/ml in 1 ml MEM supplemented with 10mM HEPES, pH 7.5, and 0.5% fatty-acid-free bovine serum albumin for 2 hours at 37°C. The cells are then washed twice with 1 ml of the same buffer.

Candidate modulator compounds are added in 1 ml of the same buffer, either alone or with 10 $\mu$ M ATP and the cells are incubated at 37°C for 30 minutes. Buffer alone and mock-transfected cells are used as controls. Samples (0.5 ml) from each well are counted by liquid scintillation spectroscopy. Agonists which activate the receptor will lead to

potentiation of the ATP-stimulated release of [<sup>3</sup>H]-arachidonic acid. This potentiation is blocked by antagonists.

#### I. Extracellular Acidification Rate

In yet another assay, the effects of candidate modulators of nGPCR-x activity are assayed by monitoring extracellular changes in pH induced by the test compounds. (See, e.g., Dunlop *et al.*, Journal of Pharmacological and Toxicological Methods 40(1):47-55 (1998).) In one embodiment, CHO cells transfected with a nGPCR-x expression vector are seeded into 12 mm capsule cups (Molecular Devices Corp.) at 4 x 10<sup>5</sup> cells/cup in MEM supplemented with 10% fetal bovine serum, 2 mM L-glutamine, 10 U/ml penicillin, and 10 µg/ml streptomycin. The cells are incubated in this medium at 37°C in 5% CO<sub>2</sub> for 24 hours.

Extracellular acidification rates are measured using a Cytosensor microphysiometer (Molecular Devices Corp.). The capsule cups are loaded into the sensor chambers of the microphysiometer and the chambers are perfused with running buffer (bicarbonate-free MEM supplemented with 4 mM L-glutamine, 10 units/ml penicillin, 10 µg/ml streptomycin, 26 mM NaCl) at a flow rate of 100 µl/minute. Candidate agonists or other agents are diluted into the running buffer and perfused through a second fluid path. During each 60-second pump cycle, the pump is run for 38 seconds and is off for the remaining 22 seconds. The pH of the running buffer in the sensor chamber is recorded during the cycle from 43-58 seconds, and the pump is re-started at 60 seconds to start the next cycle. The rate of acidification of the running buffer during the recording time is calculated by the Cytosoft program. Changes in the rate of acidification are calculated by subtracting the baseline value (the average of 4 rate measurements immediately before addition of a modulator candidate) from the highest rate measurement obtained after addition of a modulator candidate. The selected instrument detects 61mV/pH unit. Modulators that act as agonists of the receptor result in an increase in the rate of extracellular acidification compared to the rate in the absence of agonist. This response is blocked by modulators which act as antagonists of the receptor.

**Example 12 - Using nGPCR-x proteins to isolate neurotransmitters**

Isolated nGPCR-x proteins of the present invention can be used to isolate novel or known neurotransmitters (Saito *et al.*, Nature 400: 265-269, 1999). The cDNAs that  
5 encode the isolated nGPCR-x can be cloned into mammalian expression vectors and used to stably or transiently transfect mammalian cells including CHO, Cos or HEK293 cells. Receptor expression can be determined by Northern blot analysis of transfected cells and identification of an appropriately sized mRNA band (predicted size from the cDNA). Brain regions shown by mRNA analysis to express each of the nGPCR-x proteins could be  
10 processed for peptide extraction using any of several protocols ((Reinscheid R.K. *et al.*, Science 270: 243-247, 1996; Sakurai, T., *et al.*, Cell 92: 573-585, 1998; Hinuma, S., *et al.*, Nature 393: 272-276, 1998). Chromatographic fractions of brain extracts could be tested for ability to activate nGPCR-x proteins by measuring second messenger production such as changes in cAMP production in the presence or absence of forskolin, changes in  
15 inositol 3-phosphate levels, changes in intracellular calcium levels or by indirect measures of receptor activation including receptor stimulated mitogenesis, receptor mediated changes in extracellular acidification or receptor mediated changes in reporter gene activation in response to cAMP or calcium (these methods should all be referenced in other sections of the patent). Receptor activation could also be monitored by co-  
20 transfecting cells with a chimeric  $GL_{q/13}$  to force receptor coupling to a calcium stimulating pathway (Conklin *et al.*, Nature 363: 274-276, 1993). Neurotransmitter mediated activation of receptors could also be monitored by measuring changes in [ $^{35}$  S]-GTPKS binding in membrane fractions prepared from transfected mammalian cells. This assay could also be performed using baculoviruses containing nGPCR-x proteins infected into  
25 SF9 insect cells.

The neurotransmitter which activates nGPCR-x proteins can be purified to homogeneity through successive rounds of purification using nGPCR-x proteins activation as a measurement of neurotransmitter activity. The composition of the neurotransmitter can be determined by mass spectrometry and Edman degradation if peptidergic.  
30 Neurotransmitters isolated in this manner will be bioactive materials which will alter

neurotransmission in the central nervous system and will produce behavioral and biochemical changes.

### Example 13 - Using nGPCR-x proteins to isolate and purify G proteins

5 cDNAs encoding nGPCR-x proteins are epitope-tagged at the amino terminus end of the cDNA with the cleavable influenza-hemagglutinin signal sequence followed by the FLAG epitope (IBI, New Haven, CT). Additionally, these sequences are tagged at the carboxyl terminus with DNA encoding six histidine residues. (Amino and Carboxyl  
10 Terminal Modifications to Facilitate the Production and Purification of a G Protein-Coupled Receptor, B.K. Kobilka, *Analytical Biochemistry*, Vol. 231, No. 1, Oct 1995, pp. 269-271). The resulting sequences are cloned into a baculovirus expression vector such as pVL1392 (Invitrogen). The baculovirus expression vectors are used to infect SF-9 insect cells as described (Guan, X. M., Kobilka, T. S., and Kobilka, B. K. (1992) *J. Biol. Chem.*  
15 **267**, 21995-21998). Infected SF-9 cells could be grown in 1000-ml cultures in SF900 II medium (Life Technologies, Inc.) containing 5% fetal calf serum (Gemini, Calabasas, CA) and 0.1 mg/ml gentamicin (Life Technologies, Inc.) for 48 hours at which time the cells could be harvested. Cell membrane preparations could be separated from soluble proteins following cell lysis. nGPCR-x protein purification is carried out as described for  
20 purification of the  $\beta_2$  receptor (Kobilka, *Anal. Biochem.*, 231 (1): 269-271, 1995) including solubilization of the membranes in 0.8-1.0 % *n*-dodecyl -D-maltoside (DM) (CalBiochem, La Jolla, CA) in buffer containing protease inhibitors followed by Ni-column chromatography using chelating Sepharose™ (Pharmacia, Uppsala, Sweden). The eluate from the Ni-column is further purified on an M1 anti-FLAG antibody column (IBI).  
25 Receptor containing fractions are monitored by using receptor specific antibodies following western blot analysis or by SDS-PAGE analysis to look for an appropriate sized protein band (appropriate size would be the predicted molecular weight of the protein).

This method of purifying G protein is particularly useful to isolate G proteins that bind to the nGPCR-x proteins in the absence of an activating ligand.

30

**EXAMPLE 14: CLONE DEPOSIT INFORMATION**

In accordance with the Budapest Treaty, clones of the present invention have been deposited at the Agricultural Research Culture Collection (NRRL) International Depository Authority, 1815 N. University Street, Peoria, Illinois 61604, U.S.A. Accession  
5 numbers and deposit dates are provided below in Table 6.

**Table 6: DEPOSIT INFORMATION**

Clone	Accession Number NRRL	Budapest Treaty Deposit Date
nGPCR-74 (SEQ ID NO:134)	UC20088	2000 Feb 22

Some of the preferred embodiments of the invention described above are outlined  
10 below and include, but are not limited to, the following embodiments. As those skilled in the art will appreciate, numerous changes and modifications may be made to the preferred embodiments of the invention without departing from the spirit of the invention. It is intended that all such variations fall within the scope of the invention.

The entire disclosure of each publication cited herein is hereby incorporated by  
15 reference.

**What is claimed is:**

1. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous to sequences selected from the group consisting of: SEQ ID NO:135 to SEQ ID NO:268; said nucleic acid molecule encoding at least a portion of nGPCR-x.
2. The isolated nucleic acid molecule of claim 1 comprising a sequence that encodes a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268.
3. The isolated nucleic acid molecule of claim 1 comprising a sequence homologous to a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134.
4. The isolated nucleic acid molecule of claim 1 comprising a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:134.
5. The isolated nucleic acid molecule of claim 1 wherein said nucleic acid molecule is DNA.
6. The isolated nucleic acid molecule of claim 1 wherein said nucleic acid molecule is RNA.
7. An expression vector comprising a nucleic acid molecule of any one of claims 1 to 4.
8. The expression vector of claim 7 wherein said nucleic acid molecule comprises a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:134.
9. The expression vector of claim 7 wherein said vector is a plasmid.

10. The expression vector of claim 7 wherein said vector is a viral particle.
11. The expression vector of claim 10 wherein said vector is selected from the group  
5 consisting of adenoviruses, baculoviruses, parvoviruses, herpesviruses, poxviruses, adeno-  
associated viruses, Semliki Forest viruses, vaccinia viruses, and retroviruses.
12. The expression vector of claim 7 wherein said nucleic acid molecule is operably  
connected to a promoter selected from the group consisting of simian virus 40, mouse  
10 mammary tumor virus, long terminal repeat of human immunodeficiency virus, maloney  
virus, cytomegalovirus immediate early promoter, Epstein Barr virus, rous sarcoma virus,  
human actin, human myosin, human hemoglobin, human muscle creatine, and human  
metallothionein.
13. A host cell transformed with an expression vector of claim 7.
14. The transformed host cell of claim 13 wherein said cell is a bacterial cell.
15. The transformed host cell of claim 14 wherein said bacterial cell is *E. coli*.
- 20 16. The transformed host cell of claim 13 wherein said cell is yeast.
17. The transformed host cell of claim 16 wherein said yeast is *S. cerevisiae*.
18. The transformed host cell of claim 13 wherein said cell is an insect cell.
- 25 19. The transformed host cell of claim 18 wherein said insect cell is *S. frugiperda*.
20. The transformed host cell of claim 13 wherein said cell is a mammalian cell.

30



21. The transformed host cell of claim 20 wherein mammalian cell is selected from the group consisting of chinese hamster ovary cells, HeLa cells, African green monkey kidney cells, human HEK-293 cells, and murine 3T3 fibroblasts.
- 5 22. An isolated nucleic acid molecule comprising at least 10 nucleotides, said isolated nucleic acid comprising a nucleotide sequence complementary to a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:134.
23. The nucleic acid molecule of claim 22 wherein said molecule is an antisense  
10 oligonucleotide directed to a region of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:134.
24. The nucleic acid molecule of claim 23 wherein said oligonucleotide is directed to a  
15 regulatory region of a sequence selected from the group of sequences consisting of SEQ ID NO:1 to SEQ ID NO:134.
25. A composition comprising a nucleic acid molecule of any one of claims 1 to 4 or 22 and an acceptable carrier or diluent.
- 20 26. A composition comprising a recombinant expression vector of claim 7 and an acceptable carrier or diluent.
27. A method of producing a polypeptide that comprises a sequence selected from the group of sequences consisting SEQ ID NO:135 to SEQ ID NO:268, and homologs thereof,  
25 said method comprising the steps of:
- a) introducing a recombinant expression vector of claim 8 into a compatible host cell;
  - b) growing said host cell under conditions for expression of said polypeptide; and
  - 30 c) recovering said polypeptide.

28. The method of claim 27 wherein said host cell is lysed and said polypeptide is recovered from the lysate of said host cell.
29. The method of claim 27 wherein said polypeptide is recovered by purifying the culture medium without lysing said host cell.
30. An isolated polypeptide encoded by a nucleic acid molecule of claim 1.
31. The polypeptide of claim 30 wherein said polypeptide comprises a sequence selected from the group of sequences consisting of SEQ ID NO:135 to SEQ ID NO:268.
32. The polypeptide of claim 30 wherein said polypeptide comprises an amino acid sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:135 to SEQ ID NO:268.
33. The polypeptide of claim 30 wherein said sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:135 to SEQ ID NO:268 comprises at least one conservative amino acid substitution compared to the sequences in the group of sequences consisting of SEQ ID NO:135 to SEQ ID NO:268.
34. The polypeptide of claim 30 wherein said polypeptide comprises an allelic variant of a polypeptide with a sequence selected from the group of sequences consisting of SEQ ID NO:135 to SEQ ID NO:268.
35. A composition comprising a polypeptide of claim 34 and an acceptable carrier or diluent.
36. An isolated antibody which binds to an epitope on a polypeptide of claim 30.
37. The antibody of claim 36 wherein said antibody is a monoclonal antibody.

38. A composition comprising an antibody of claim 36 and an acceptable carrier or diluent.

39. A method of inducing an immune response in a mammal against a polypeptide of claim 30 comprising administering to said mammal an amount of said polypeptide  
5 sufficient to induce said immune response.

40. A method for identifying a compound which binds nGPCR-x comprising the steps of:

- 10                   a)     contacting nGPCR-x with a compound; and  
                    b)     determining whether said compound binds nGPCR-x.

41. The method of claim 40 wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268.

15

42. The method of claim 40 wherein binding of said compound to nGPCR-x is determined by a protein binding assay.

43. The method of claim 40 wherein said protein binding assay is selected from the  
20 group consisting of a gel-shift assay, Western blot, radiolabeled competition assay, phage-based expression cloning, co-fractionation by chromatography, co-precipitation, cross linking, interaction trap/two-hybrid analysis, southwestern analysis, and ELISA.

44. A compound identified by the method of claim 40.

25

45. A method for identifying a compound which binds a nucleic acid molecule encoding nGPCR-x comprising the steps of:

- a)     contacting said nucleic acid molecule encoding nGPCR-x with a  
                    compound; and  
30                   b)     determining whether said compound binds said nucleic acid molecule.

46. The method of claim 45 wherein binding is determined by a gel-shift assay.
47. A compound identified by the method of claim 45.
- 5 48. A method for identifying a compound which modulates the activity of nGPCR-x comprising the steps of:
- a) contacting nGPCR-x with a compound; and
  - b) determining whether nGPCR-x activity has been modulated.
- 10 49. The method of claim 48 wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268.
50. The method of claim 48 wherein said activity is neuropeptide binding.
- 15 51. The method of claim 48 wherein said activity is neuropeptide signaling.
52. A compound identified by the method of claim 48.
- 20 53. A method of identifying an animal homolog of nGPCR-x comprising the steps:
- a) comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequence consisting of SEQ ID NO:1 to SEQ ID NO:134, and portions thereof, said portions being at least 10 nucleotides; and
  - b) identifying nucleic acid sequences of the animal that are
- 25 homologous to said sequence selected from the group sequence consisting of SEQ ID NO:1 to SEQ ID NO:134, and portions thereof, said portions comprising at least 10 nucleotides.
54. The method of claim 53 wherein comparing the nucleic acid sequences of the
- 30 animal with a sequence selected from the group of sequences consisting of SEQ ID NO:1

to SEQ ID NO:134, and portions thereof, said portions being at least 10 nucleotides, is performed by DNA hybridization.

55. The method of claim 53 wherein comparing the nucleic acid sequences of the animal with a sequence selected from the group of sequences consisting of SEQ ID NO:1  
5 to SEQ ID NO:134, and portions thereof, said portions being at least 10 nucleotides, is performed by computer homology search.

56. A method of screening a human subject to diagnose a disorder affecting the brain  
10 or genetic predisposition therefor, comprising the steps of:

(a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR-x that is expressed in the brain, wherein the nGPCR-x comprises an amino acid sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID  
15 NO:268, and allelic variants thereof, and wherein the nucleic acid corresponds to a gene encoding the nGPCR-x; and

(b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR-x in the nucleic acid correlates with an  
20 increased risk of developing the disorder.

57. A method according to claim 56, wherein the disease is a mental disorder.

58. A method according to claim 56, wherein the assaying step comprises at least one  
25 procedure selected from the group consisting of:

a) comparing nucleotide sequences from the human subject and reference sequences and determining a difference of at least a nucleotide or at least one codon between the nucleotide sequences from the human subject that encodes a nGPCR-x reference sequence;

(b) performing a hybridization assay to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences;

(c) performing a polynucleotide migration assay to determine whether  
5 nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences; and

(d) performing a restriction endonuclease digestion to determine whether nucleic acid from the human subject has a nucleotide sequence identical to or different from one or more reference sequences.

10

59. A method according to claim 58 wherein the assaying step comprises: performing a polymerase chain reaction assay to amplify nucleic acid comprising nGPCR-x coding sequence, and determining nucleotide sequence of the amplified nucleic acid.

15 60. A method of screening for an nGPCR-x hereditary mental disorder genotype in a human patient, comprising the steps of:

(a) providing a biological sample comprising nucleic acid from said patient, said nucleic acid including sequences corresponding to alleles of nGPCR-x; and

20 (b) detecting the presence of one or more mutations in the nGPCR-x allele;

wherein the presence of a mutation in a nGPCR-x allele is indicative of a hereditary mental disorder genotype.

25 61. The method according to claim 60 wherein said biological sample is a cell sample.

62. The method according to claim 60 wherein said detecting the presence of a mutation comprises sequencing at least a portion of said nucleic acid, said portion comprising at least one codon of said nGPCR-x allele, said portion comprising at least 10  
30 nucleotides.

63. The method according to claim 60 wherein said nucleic acid is DNA.

64. The method according to claim 60 wherein said nucleic acid is RNA.

5 65. A kit for screening a human subject to diagnose a mental disorder or a genetic predisposition therefor, comprising, in association:

(a) an oligonucleotide useful as a probe for identifying polymorphisms in a human nGPCR-x gene, the oligonucleotide comprising 6-50 nucleotides in a sequence that is identical or complementary to a sequence of a wild type human nGPCR-x gene sequence or nGPCR-x coding sequence, except for one sequence difference selected from  
10 the group consisting of a nucleotide addition, a nucleotide deletion, or nucleotide substitution; and

(b) a media packaged with the oligonucleotide, said media containing information for identifying polymorphisms that correlate with mental disorder or a genetic  
15 predisposition therefor, the polymorphisms being identifiable using the oligonucleotide as a probe.

66. A method of identifying a nGPCR-x allelic variant that correlates with a mental disorder, comprising the steps of:

20 (a) providing a biological sample comprising nucleic acid from a human patient diagnosed with a mental disorder, or from the patient's genetic progenitors or progeny;

(b) detecting in the nucleic acid the presence of one or more mutations in an nGPCR-x that is expressed in the brain, wherein the nGPCR-x comprises  
25 an amino acid sequence selected from the group consisting of SEQ ID NO:135 to SEQ ID NO:268, and allelic variants thereof, and wherein the nucleic acid includes sequence corresponding to the gene or genes encoding nGPCR-x;

wherein the one or more mutations detected indicates an allelic variant that correlates with a mental disorder.

30

67. A purified and isolated polynucleotide comprising a nucleotide sequence encoding a nGPCR-x allelic variant identified according to claim 66.
68. A host cell transformed or transfected with a polynucleotide according to claim 67  
5 or with a vector comprising the polynucleotide.
69. A purified polynucleotide comprising a nucleotide sequence encoding nGPCR-x of a human with a mental disorder;  
wherein said polynucleotide hybridizes to the complement of a sequence  
10 selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 under the following hybridization conditions:  
(a) hybridization for 16 hours at 42°C in a hybridization solution comprising 50% formamide, 1% SDS, 1 M NaCl, 10% dextran sulfate and  
(b) washing 2 times for 30 minutes at 60°C in a wash solution comprising  
15 0.1x SSC and 1% SDS; and  
wherein the polynucleotide that encodes nGPCR-x amino acid sequence of the human differs from the sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 by at least one residue.
- 20 70. A vector comprising a polynucleotide according to claim 69.
71. A host cell that has been transformed or transfected with a polynucleotide according to claim 69 and that expresses the nGPCR-x protein encoded by the polynucleotide.  
25
72. A host cell according to claim 71 that has been co-transfected with a polynucleotide encoding the nGPCR-x amino acid sequence set forth in a sequence selected from the group consisting of SEQ ID NO:1 to SEQ ID NO:134 and that expresses the nGPCR-x protein having the amino acid sequence set forth in SEQ ID NO:135 to SEQ  
30 ID NO:268.



73. A method for identifying a modulator of biological activity of nGPCR-x comprising the steps of:

a) contacting a cell according to claim 72 in the presence and in the absence of a putative modulator compound;

5 b) measuring nGPCR-x biological activity in the cell;  
wherein decreased or increased nGPCR-x biological activity in the presence versus absence of the putative modulator is indicative of a modulator of biological activity.

74. A method to identify compounds useful for the treatment of a mental disorder, said  
10 method comprising the steps of:

(a) contacting a composition comprising nGPCR-x with a compound suspected of binding nGPCR-x;

(b) detecting binding between nGPCR-x and the compound suspected of binding nGPCR-x;

15 wherein compounds identified as binding nGPCR-x are candidate compounds useful for the treatment of a mental disorder.

75. A method for identifying a compound useful as a modulator of binding between nGPCR-x and a binding partner of nGPCR-x comprising the steps of:

20 (a) contacting the binding partner and a composition comprising nGPCR-x in the presence and in the absence of a putative modulator compound;

(b) detecting binding between the binding partner and nGPCR-x;

25 wherein decreased or increased binding between the binding partner and nGPCR-x in the presence of the putative modulator, as compared to binding in the absence of the putative modulator is indicative a modulator compound useful for the treatment of a mental disorder.

76. A method according to claim 74 or 75 wherein the composition comprises a cell  
30 expressing nGPCR-x on its surface.

77. A method according to claim 76 wherein the composition comprises a cell transformed or transfected with a polynucleotide that encodes nGPCR-x.

78. A method of purifying a G protein from a sample containing said G protein comprising the steps of:

- a) contacting said sample with a polypeptide of claim 1 for a time sufficient to allow said G protein to form a complex with said polypeptide;
- b) isolating said complex from remaining components of said sample;
- c) maintaining said complex under conditions which result in dissociation of said G protein from said polypeptide; and
- d) isolating said G protein from said polypeptide.

79. The method of claim 78 wherein said sample comprises an amino acid sequence selected from the group of sequences consisting of SEQ ID NO:135 to SEQ ID NO:268.

80. The method of claim 78 wherein said polypeptide comprises an amino acid sequence homologous to a sequence selected from the group of sequences consisting of SEQ ID NO:135 to SEQ ID NO:268.

81. The method of claim 78 wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: SEQ ID NO:135 to SEQ ID NO:268.

82. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence homologous a sequence of SEQ ID NO:268; said nucleic acid molecule encoding at least a portion of nGPCR-x.

83. The isolated nucleic acid molecule of claim 82 comprising a sequence that encodes a polypeptide comprising a sequence of SEQ ID NO:268.

84. The isolated nucleic acid molecule of claim 82 comprising a sequence homologous to a sequence of SEQ ID NO:134.

85. The isolated nucleic acid molecule of claim 82 comprising a sequence of SEQ ID NO:134.
- 5 86. An expression vector comprising a nucleic acid molecule of any one of claims 82 to 85.
87. A host cell transformed with an expression vector of claim 86.
- 10 88. An isolated polypeptide encoded by a nucleic acid molecule of claim 82.
89. The polypeptide of claim 88 wherein said polypeptide comprises a sequence of SEQ ID NO:268.
- 15 90. The polypeptide of claim 88 wherein said polypeptide comprises an amino acid sequence homologous to a sequence of SEQ ID NO:268.
91. An isolated antibody which binds to an epitope on a polypeptide of claim 88.
- 20 92. A method for identifying a compound which binds nGPCR-x comprising the steps of:
- a) contacting nGPCR-x with a compound; and
  - b) determining whether said compound binds nGPCR-x.
- 25 93. A method for identifying a compound which modulates the activity of nGPCR-x comprising the steps of:
- a) contacting nGPCR-x with a compound; and
  - b) determining whether nGPCR-x activity has been modulated.
- 30 94. The method of claim 93 wherein the nGPCR-x comprises an amino acid sequence of SEQ ID NO:268.

95. A method of screening a human subject to diagnose a disorder affecting the brain or genetic predisposition therefor, comprising the steps of:

5 (a) assaying nucleic acid of a human subject to determine a presence or an absence of a mutation altering an amino acid sequence, expression, or biological activity of at least one nGPCR-x that is expressed in the brain, wherein the nGPCR-x comprises an amino acid sequence of SEQ ID NO:268, and allelic variants thereof, and wherein the nucleic acid corresponds to a gene encoding the nGPCR-x; and

10 (b) diagnosing the disorder or predisposition from the presence or absence of said mutation, wherein the presence of a mutation altering the amino acid sequence, expression, or biological activity of the nGPCR-x in the nucleic acid correlates with an increased risk of developing the disorder.

15

20

## SEQUENCE LISTING

<110> Pharmacia & Upjohn Company  
Vogeli, Gabriel  
Wood, Linda S.

<120> Novel G Protein-Coupled Receptors

<130> 00100PCT1

<150> 60/187,828  
<151> 2000-03-08

<150> 60/187,715  
<151> 2000-03-08

<150> 60/187,929  
<151> 2000-03-08

<150> 60/187,930  
<151> 2000-03-08

<150> 60/187,825  
<151> 2000-03-08

<150> 60/187,833  
<151> 2000-03-08

<150> 60/187,830  
<151> 2000-03-08

<150> 60/187,829  
<151> 2000-03-08

<150> 60/187,582  
<151> 2000-03-08

<150> 60/187,581  
<151> 2000-03-08

<150> 60/187,714  
<151> 2000-03-08

<150> 60/189,294  
<151> 2000-03-08

<150> 60/187,874  
<151> 2000-03-08

<150> 60/187,928  
<151> 2000-03-08

<150> 60/188,049  
<151> 2000-03-08

<160> 273

<170> PatentIn version 3.0

<210> 1  
<211> 642  
<212> DNA  
<213> Homo sapiens

<400> 1

ggaatttagt tgggcagaag ggaataaag tgaggatggt taatgggtac aaaaaatagt 60  
 taggaaaaaa atgaataaga tctagtatta gatagcacia cagggtgatt gtagtcaata 120  
 taatttagtt gtacaattta aaataactaa aagaatataa ctggattggt tgtaacacaa 180  
 atgataaacg cttgaggtaa tggatacgat atttaccctg atgtaattat tacacattgc 240  
 acgtctgtat tcaaaatacc ccatctaact cataaatatt tatatctact atctacacaa 300  
 aaaattaaaa attaaaaaaa tttttgcatg atgatcttaa ctgaattttt caataataaa 360  
 acattgtctg ttttcattaa gttcaattta gcaatttcaa ttatgtttta ttatttttgc 420  
 atcctgaata aaaaatcttc ttatactgca agattttgaa ggcaatctag acttacttct 480  
 agaattgtta tgttctacct gttataatca ggcttacaat tcatgtccaa ttaattttca 540  
 tatgtaaagt gagttatatt tttcatgaag ttgttcagtt tttcagcccc acttaaaaaa 600  
 atgtagaatt gtttcttgct cagttaaact gacctgcttt tt 642

<210> 2  
 <211> 660  
 <212> DNA  
 <213> Homo sapiens

<400> 2  
 cagggtgcagc atcgtgtcct cagtgtcctg cccctgctt ccacccggtg tcgacagctg 60  
 cacgggtccac cccacgcctg cctttccatc gttcctcatc agccctgtga tctttcctgt 120  
 ggccctgctg tgtgtgtgcc ctgtgaggtc ctgtggacac aagagactgc acgggccaca 180  
 ccccagctg ggtgagtcct ctccctcctg ggtactctgg acagtaaaga aagatggaca 240  
 cgtgggctcc gtggagcatg aggtagtcca ggacctcggc ggccacaggt cctgcctccc 300  
 tgcttctcgt gcctccctc cctttgggtc tctgtccac ctgggtaaac gcttcgttcc 360  
 caccctcga agggtaaatc gagctccttg gtggtaaac accactgcc ctagtcaga 420  
 gggctccctc tctctgatgt catgggtgcc tggctgcct ggtagaattt tagctgcttt 480  
 ataacctggt cctgaaatga accactggga agaaataggg taaaatgaac acacagctgc 540  
 cacactgcat cccaaccctg tgtgacctta tcaccgcaga cttttgtggc aagatgacag 600  
 catctcagtt tgcttgagaa gcttattttt gccaggctg ttaccaccag gcaggcacca 660

<210> 3  
 <211> 436  
 <212> DNA  
 <213> Homo sapiens

<400> 3  
 ctattatttc ttaacatact gcatttttcc gattctctct aagtatctgt ttctgtaact 60  
 cctattggac atttacttct ctttcacatt gtctgcttta tctcttaact ttgtgtttt 120  
 ctgtctctca ctgctgtatt gtgagttatt acttagctct tccagttaat tctttaagct 180  
 tttttgttaa tctctttatc agttcattgt gtttattatt tcagtgacta aatttaattg 240

ctcaaagttt tatttggtcg tttaaaattt gcttttgtct tcatagtat tttgtcctgt 300  
 tctctttatc tctttattta tttttgatgc tttcatctgc ttattatttt taagatat 360  
 atttcttagc ctctttgaga tattctatta tctctgtgcc taggatcatt aaatctccca 420  
 ctacgtctgt agactc 436

<210> 4  
 <211> 707  
 <212> DNA  
 <213> Homo sapiens

<400> 4  
 acttctgggc cacggaagc cctactgtct aaatgctttt caggccaatt tgaagaagta 60  
 attagactta ctggaagctt ctgtgaataa ttctgcaagt acaattatgg acttcccagg 120  
 aaatattgcc ttcaatatag aaaagcttgt cagttgattc tgatgagata tatgtaaaat 180  
 ttgagatttt gatattagaa tgagtaaaat gatgacatca cgatgtatta aagttggggg 240  
 ttattttttg gaattaattg tcatcaggta aaaagccagc tataagtcaa ataaaatata 300  
 atcatgttct tccgtcttta gcactcatct tttcttgttc taaatgttga caaatgactg 360  
 taaatttaac aagcttatag ataataattg aaaagtcttc taagaactga aaattgataa 420  
 acacatggca atggcaggct attgcagtgc aattataaga tgttgtgtgg atgccctga 480  
 agtgcctata aatgaatgtg acttcagtac tactgccaaa tgagtccaat atcccacaaa 540  
 tgaactgaaa ataaagtgcc tggaatactg tgtctacagt gtcactgtaa agttactgtc 600  
 atgctgtatt actgaaatga tttgctggaa agtaacatgg cacatatatg caccaagaga 660  
 gttaaatctc atcttattct atgaaaaatc atgttaacca ttcattga 707

<210> 5  
 <211> 529  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 aacattatta ctttctttta tgaatattct tggctcttcc aaaacaaaac aagctattgg 60  
 ttttaataat tatggtataa tcaaataatg aaactctatg catttgtaa agtaactttt 120  
 caaaagaata tcttgtaaca tagaataaca gatcctagtg cattaccac tctttgggct 180  
 ttatcgcttt tccaccatca ttatctgcat cactgcctgc aggttttcta cacggccagg 240  
 gttggtctct gcctgctcaa tagtcaagtc aaaagaggca ggaaattaac accctctgga 300  
 ggcagccttt gaggaatgat ccatgggagg tggagtataa atacctcagc tctgtttcct 360  
 ctagagatat aactaaggaa tgggttttac attgtttctc agagtttcct caaggtttta 420  
 aacttcaatc acccacaggg ggtagtgggc tttatcatag tatacatccc tttgtggctt 480  
 cccttccttc ttgtctcact tctccattcc aaactaggat ttatttctt 529

<210> 6  
 <211> 688  
 <212> DNA  
 <213> Homo sapiens

<400> 6  
 aagtattctt gtcacggaaa gaagaaaagg gttgggtagt tacaggggga caacaatgcc 60  
 agaactgggg agtgtggact gggatacaag agaatgaggg agctcaggat gagcagaagg 120  
 gcggggaagc aatattcatt aagcaccttc tatgtgccag tcaataggcc aggcttcaaa 180  
 ttattacctt gctgaaatct tcacagcagc cctctaatag gtatttatcc ctgattccat 240  
 atccatgctc tgcttcccct cctattacaa tggctgaaga attcaaacc ctttcaaagg 300  
 ctagcactgt catttgtcct ctagatccca tcccctccat tttctttttt attgaaacat 360  
 tctcaatggg attcaaacat actctgctct ctcttctatt aaataggcaa atgcaactca 420  
 tcaagctctt tttctccctt ggctactgcc ccatttctct acttctttc atggcagaac 480  
 ttctcgaag agtttttcac aatcacttca tttccacacc tctaactgac ttttgaacac 540  
 aactagagga ggagtaggag gggacactca ttccaaagt tccaattaag cccaatcctt 600  
 taaaagtatt atgttgctcat gatggctgtt aagagcatgg tgaaaagata ttagaataag 660  
 atgtggggaa tcatgaccgt gagacaga 688

<210> 7  
 <211> 552  
 <212> DNA  
 <213> Homo sapiens

<400> 7  
 aaagaaaaag aaagagtagt gtaacaattc cacttctgga ttaacattgt aaggagactg 60  
 tggacctgtt acagcagaaa acagatataa taggcaaaaa ttatttttta aaaaatctoc 120  
 agaaattggt ctaaaaacat acagcagact tttaaaaaac ttgtctgaga aaatgtacta 180  
 aatctctgta agacaaacaa gagtctgtgg cacgtgagca atgtttgcct cactctaacc 240  
 tctccctccc aggtcacctt cataaaaagt caactctggg aagggtgtgcc caaattgaga 300  
 ttacctgccc cattaatttc caatcaaagg atacagtata tcaccaggaa ggtagccacc 360  
 agcatttctc agcccctctt actccaagt gcagaggata aattcctggg gagtatggcc 420  
 aggaggccac gtggccacct ggccaccact aatagatcag aggattaatc tcacacatgg 480  
 aaggatgagc atactgggcc cctgattgcc ctgacccag cttacttata ggatggaagt 540  
 ttcacatcag ga 552

<210> 8  
 <211> 684  
 <212> DNA  
 <213> Homo sapiens

<400> 8



```

agttaacaaa aaaatactac ttaacctctg ctagaacata atgtgatata tttttgacac    60
ctcttagctt ctttagctga atttcagaaa tgcaaccatt agtattaaga agcagggtact    120
aaggattttc caaatcattt tgttattctt atcaatattt ctagtattct tttagatccc    180
ttcactcact ttctctattg ctttccattt cctgaagttt taaataaaat ttcccttctg    240
tttgtcttgt agggaaaaatc atcatgctta ccacatagaa tgtgagttgt aggagagaca    300
caatgggaga catcggttaa gggacaaaag acattaacat tttaggtgat tgtgagttca    360
taatttttcc agaacacaag cattgcatgg ctactctaata actagatt attaaaatag    420
atatatcttt gccctacctg ataaacacta ttgtataag tgaatatatt tttaatatta    480
atccaatata ttccataaga aatatttgat ttgcaaagta atctgagcat tacgatgatt    540
ccctatctaa atactggcat ggtgaaaatg aggacaaatc taccctttct ctaatgtagt    600
tacaggcaag ctatactcat ataataaaca tagaacgtac aatcaaaacc aatgcatgag    660
tgtaggatgc aactaaagtc aaga                                           684

```

<210> 9  
 <211> 641  
 <212> DNA  
 <213> Homo sapiens

```

<400> 9
atcattttga taaccagtct gatctgagaa aattaaacca tgtcattcaa aacatgtcct    60
ccccaaattt aagaaacatt aggtcaatct cctgggttaa taatagctgt atgttttagt    120
agattttgaa atattatgta atcatttgaa attataagct tctggccac aacttgactg    180
acaaatacct gtttcattat ttttaactag cttttgttgg actacatata tccaaagaca    240
aaagaaagat aaaagttgaa ataatccaac agttatccta cacaaaagta tgacaaaatt    300
accgttgtag aaattgaact catcaagcct gaacttttga ctttgaacaa ttacatggaa    360
gagtgccacc atggtgaact gtcagacctg tacagcatca cagccaactc tatacacaaa    420
caagggtggg ctgtattctg accattattg gaataaatta tcctgattac ctaatgtctc    480
ttcacacca ctaaattatt tattattatt atatttttac actgccatca aattaaagtt    540
gctaaaacac aactttgctc atgttcaaaa ttctatagt gtgcctcaac aatcactaac    600
taatcctcag aattaattac ctactaattt gtttttgaca t                                           641

```

<210> 10  
 <211> 520  
 <212> DNA  
 <213> Homo sapiens

```

<400> 10
tctagaatct atacatacta tgtccaatcc ctgttcaca agtagttatt tatatgtgcg    60
aaggttcata ctctgattt tccttttgct ccagggcaaa gaaaagatac tgaaatataa    120
ggtgagctta tatcagccca gtagtaagcc agtgagggt accacagttt ggaagaagca    180

```

ggggtgaaact ttacatgag attgggggga aaaaccatac tgaataataa aggggttttaa 240  
 ctgagattga aagatagtgc ttgagaagc acacaaaaga ttcaaaatgg gcgtataaag 300  
 aatgacctgt gctgaaaaac acatttttgc gctacaaggg acccaattga ctagatgaga 360  
 atttgtgtgg aaaaggagtt gataaggcag gctggcacat tgcagccaat ctgtgaaagg 420  
 cttttcatgt cctgtgaaca ggaaatcaca tatcacaaga gtggtctagg aatctgtgtc 480  
 tggcaaccct acagtggggc agactgaaga ggaataaacg 520

<210> 11  
 <211> 668  
 <212> DNA  
 <213> Homo sapiens

<400> 11  
 atgggcaccg ctctctttaa agtacacttt cctgactcag ctgtcctctt ttcctcctcc 60  
 attcccacca attctgggct acaggctttt cctctactct cccacagcat cctccctgag 120  
 ccctcaatca aagcacctac aatactgccc tcatagggag gtgctatctt tctgtctttc 180  
 cctgagcgct gggaccatt gcatttcacc catttatccc caaggcctag cacatgtcta 240  
 gcacaacaca gcaattaaat aaaccctgtg gaataaatta attgtggaat agcctgggtt 300  
 ccatggatgg ttatacaggt tgtgactgc acaaccatgt gcaacattcc tggaaaaaga 360  
 cagaaattta ttgattggtt gggggttttg aatagccaag gaaaactatt tgaccattgc 420  
 atgccctcta cctgggaaaa tcacataccc taacaacttc ttaggcctta ctgcatggtc 480  
 acatggggta acattcatat agtttctcca gctctctagt ctgccacaa aggtgatatt 540  
 gttcaaaggg gcaatctttc cttgccttcc accagtctat tcttaacttg acccaagtaa 600  
 tctctttcac tgcttaccaa agatatttca gcttcagcta tcctgtttgc agaattgtga 660  
 cgtatttcc 668

<210> 12  
 <211> 625  
 <212> DNA  
 <213> Homo sapiens

<400> 12  
 tcaatagcaa taaagcacac caagcacaca gatctcgact ttgaaatgcc acttctccat 60  
 cttaaaagac aaaaacagga catcttagac aaatggccaa ctccagggtg gttggggcaa 120  
 ggaaagaaga cgtgcttgtg cacatcttgg tacatcaggt ttaggaagct gtcactggtc 180  
 aaatctggga caacttgaac atcaaaataa ataatcattg taatggatta taactcatcg 240  
 atgtaagtct ctaagtacac acttatatca atacatatgt acatatacac atacatacat 300  
 ctttacatag tactgaatgg caactaataa tggcatttgg caaactgtta tgctaacaat 360  
 taactcaggc aagaaacatc aatggagggt aaaactggta gataaaattg ggatgagtag 420

attttacaca gtctccaagt gactttccac aaaataccca ttattacaaa ggaaaagata 480  
 gataggtttg cagcagaaaa aaaaatgtca gacatcatct taactagggg atcagtgtta 540  
 acttctccag catgagacaa gtagacaaac aactgccatc agagaggatg aagtaagaca 600  
 cagcatcact tctgtgaaat tctgg 625

<210> 13  
 <211> 616  
 <212> DNA  
 <213> Homo sapiens

<400> 13  
 tccgatgatg ttaacacccat attattttta aagaacatga agattacata agagtaggca 60  
 tttgcctatt tgtattttta agagtctgct cagctcttaa caaggaggc cctatgcaaa 120  
 atgagaaata aagtgtaaaa cgatttgctt gtcagtctga aataacttag gtgtcaaaaa 180  
 caagtaactt tcaccctcct tcaacctgtc ctcttgccat ttagcaatct aaaataatta 240  
 tccaatgtat ggttgccact caaaaatcat gttaaacttg agatattctg aattttgtgt 300  
 acaatttttg gtagagggta agagatagag aaaaatctta cattgtgttc agtgaattcc 360  
 cagacctcgg gggtaaaata agtgcaggaa gaatctcatc aggatatctc gggcaatttt 420  
 tcattagtac gcatgacaag ctgtttcacc acaggctatt gtttttatgg aaagttcaaa 480  
 tatagcagga tgggatgtat ggtgtgatat taacacatat gaacacaatt attacctatt 540  
 ttaggtatat acgacctttg tcacctagaa acattgatac tcttcattat gatgtacttt 600  
 tatagaataa gataaa 616

<210> 14  
 <211> 599  
 <212> DNA  
 <213> Homo sapiens

<400> 14  
 ggcctctctg aaggggaagc aagcttgcat ctagacttct ttctaaagat aacctagaca 60  
 ataataaata cagctgccac cagcctccta tgcactagag gcattattct aggagtttcc 120  
 gtgtattaag cttatcctga aattagttcc ttcttatgac tgagaggaga agtattacat 180  
 attgatttca ttgttagaaa tgggaaaatt ttaacaagt gtatttagag ggcaaccaca 240  
 ttttctgctc tgcaacctgc ttctccccct tcacgtcagg acatctagat gaaccctc 300  
 ttcggaaagg ctgcagagaa acatgtccta cagacctact atcatctggt taacaactcc 360  
 cagtgagcgg accaaaattc cagacgttc ccaactttct tccactgcac ggatgctgcc 420  
 acacatgctc atatacctct gaaccttcca gtgactacgg cacagcgaca gctgagttcc 480  
 tgggcgcaga accactgggg cacgttttgg cagctatgag caaatcactg tgcacaaagg 540  
 caatcccagt ttacacttcc acagagagga actgaatata ctgcccacc tcacctgac 599

<210> 15  
 <211> 617  
 <212> DNA  
 <213> Homo sapiens

<400> 15  
 aatactagat tccttcaggg actttttaga acaaggacaa gaataatcct ttctcgacaa 60  
 agtaaggagt gatctatctc aaggcagaag catattctcc tacaccagga ggaatttcca 120  
 ttaacataag caatgcccc aaggatgcttg ttatcatttt tattcaatgt tgttttctgt 180  
 gttctggcca atattatgac ttgagataca agtgagaaag atgactaaag gaattcatga 240  
 gacaagatga tcactattca ggagatggta tgattgtcta tctagaaaaa aagaaatgac 300  
 tccatcgtaa ttctgggaat ttactaacag tggctgggtc ctggacaac atttaaaaaa 360  
 tcaatcgttt ccttgtgtgg cagcaataac catttagaaa atggagtaaa tgcggagtta 420  
 aggggctgtg aatatataac agcaagaact cctgatctgc cgtcccgaca agtcgcctcc 480  
 ggagtggaca ccggccaggg aaggcaggtc tctggaggga aggtagagag aagatacggg 540  
 ggatctgccc ctccccagg aagctccccg agaaagggcc acaactgttt actccagcag 600  
 gctctggggg gattcag 617

<210> 16  
 <211> 518  
 <212> DNA  
 <213> Homo sapiens

<400> 16  
 gaaaaccttt gactactttc tctgtctcac ggtcatgatt cccacatctt tattctaata 60  
 tcttttcacc atgctcttaa cagccatcat gacaacataa tacttttaaa ggattgggct 120  
 taattggaca ctttggaatg agtgtcccct cctactcctc ctctagtgtg gttcaaaagt 180  
 cagtttagagg tgtggaaatg aagtgtattgt gaaaaactct ttcgagaagt tctgccatga 240  
 aaggaagtag agaaatgggg cagtagccaa gggagaaaaa gagcttgatg agttgcattt 300  
 gcctatttaa tagaagagag agcagagtat gtttgaatac cattgagaat gtttcaataa 360  
 aaagaaaaat ggaggggatg ggatctagag gacaaatgac agtgctagcc tttgaaaggg 420  
 gtttgaattc ttcagccatt gtaataggag ggaagcaga gcatggatat ggaatcaggg 480  
 ataaatacct attagagggc tgctgtgaag atttcagc 518

<210> 17  
 <211> 375  
 <212> DNA  
 <213> Homo sapiens

<400> 17  
 acagggcata ctcgccttcc acccacttta aaacagcctg caaggcagtg tgtgacctat -60  
 ggctttaact ctgatgagga ggattcctca tggcatgggt tgctgagaac cctgaatcac 120  
 aaggtataaa gcagggaccg aaggactgtg cccactgcag caaccccccg ctgggtttta 180

tgctctcctg ttgccaccct gaaattttta aagacttttt acgggggtctt gctctgtcat	240
ctaggctgga gtgcagtac atgatgtctt atacctcatc tggctgagac tcactagaga	300
aggtcactag ttagaactag agagggggct gggcacagtg gctcatgcc gcactttggg	360
aggctgaggc aggag	375

<210> 18  
 <211> 687  
 <212> DNA  
 <213> Homo sapiens

<400> 18	
ctttagcaa tataaagcct taaatttttt ttctgttaga aaatatacaca cagatggcta	60
attatatgcc atataaagcc attaaggaag aaaggatggc aaatgctcct tttagtgaga	120
cttctttgtt atgagatctg ggtataaaaa tgtgcagggtg tgtaaacaga ggaaggagaa	180
ttctgattaa gtccctcaag aattgaagaa aatgggtga gagacagaga acaactgtga	240
gctaggaaaag ctcaaggagt aaacctaaca agaaagtta agcaatggct acttttatac	300
agtttatttt agtaagtga aataactaaa atgaagttat ttataaagtt tatttgagtt	360
gttttctgat aattaaatag catgagaaat gggaggaatt tgagatattg cagttagaaa	420
gggagcagtg caccaactt attcttaact taaaagtca tactcttacc taaggtaagt	480
cctaattgga caccaactta aagctgaatt agacaggaat attgcaatga ataagcaatg	540
actattcaca atctactcag cataaaacag gttcattaag aaaggttctg caataacact	600
ctatgtaaga gtttatggaa caattaatag aataaaattg atgtacattt tatgtactac	660
tgcaattttac atattctaag gcacgag	687

<210> 19  
 <211> 546  
 <212> DNA  
 <213> Homo sapiens

<400> 19	
cccttgagca cacacagggc gatacttgcc acaggtggg actgaaggct tctttcttgg	60
cttatagttt ggaagcaatg ggagttggga gctccaaatc attcatggga caaatatcct	120
gtcttatatt gcttaaaaaa aaatcctatc taatttttaa gacaggggtg tttgctctta	180
aagcactttg catttaattg tgtaattac agaaattttc aatgctctct gaagaggtaa	240
ttgatattaa ccattggaat tctaatagct aacacatatt gggcatacgg tttttcacat	300
gtctaaacag tccatgtttc cttaaaaatg cagattgcag ggccccacac tggctgggga	360
attcgactt ccagtaaaca cttcagatga tttcatgat ctttcaagtt cggggaaaat	420
ggagctcgtt ttccactaga ttaaagcagt attccactgt atgcgttctc aggccctaaa	480
agaatcaaca ctctcaata agtaaacatt cacttaaaca tatccaggtg gatccaatga	540

tctacc 546

<210> 20  
 <211> 547  
 <212> DNA  
 <213> Homo sapiens

<400> 20  
 ctgctctata taaagatata gtccatgtat atggctgagt cttttatagt ccaaaatgta 60  
 tttttctgtg tactatgggtt tattaacttg acttattttt cttctttcag atttaaaaaa 120  
 tgttaactaa ttaaagtaac ttcccaagta cctaccaatg acattaatct tcctcttttt 180  
 gtcgtttggt ctttttacct ccaaatccta ttaatacagc aactttttta tatgattgtc 240  
 tactttttcag agtacttctt aacaacatag caaatgccaa aatgttaatg gaagtattaa 300  
 tgaaaacatg caaaaaatat ttctttatga ttctgataat tattgaaatt gccttagatt 360  
 aaacatgaat aaatttaatt attatatatg tattcaaata gttggatata tagtcctgag 420  
 aaagaatcct tcactacata tggtataaaa atgggaatga acacattacc taagaagtct 480  
 gcactagaaa taataagata ccttttcatt cttgacatct ttcttctttt tgaaccaagt 540  
 atctgta 547

<210> 21  
 <211> 731  
 <212> DNA  
 <213> Homo sapiens

<400> 21  
 tatcatgctg ccgcttccaa tgggcatctg ttccaccatg ttgtgggcat tcattggagc 60  
 tttgttgctt ctactgtta aaccctcctg tataattctg gggccccagc agaaaacagt 120  
 atgttaccct aaaatagggc aattgaagga tctttcaaga agggacaagt tgtaaagggtg 180  
 ggcagcacia agggaaacca acaaaaaatg aagacctggt gggacaggga cagagtgact 240  
 ggatgctgga gagacccaaa gctgcaaagg aaaggagcaa ggggaacaat accccaccct 300  
 ctccccctcc acctcccacc toccacctcc attcttctcc agtgggtgccg cccattgggc 360  
 aaaccagacc agaagccagg aagcatgaga gttagctga tgcagcccat acagatcaga 420  
 ctcttggtgact tcagagtggg ggagggtgag agggatgaag tctggaggca ccaattggga 480  
 aggccatcca gaatgctcct attctgtttg ggagctgggg atgggaatgt cccttctga 540  
 ggggtattta tggaataaat caaatcaaat cacagaaatc aaatcacaga aatcaaagct 600  
 ggagattctc tctccctcta cttgctggca gccaggatgt gggctcatga cctaaactca 660  
 gtcattcaga aattcccccg gggaatgcag tcttacagga gtagctcaag gccaggcagt 720  
 ggctcacacc t 731

<210> 22  
 <211> 462

<212> DNA  
<213> Homo sapiens

<400> 22  
cccatctgtc tgaatgcctc ctgtagtggg ggactcactc cctaataaat caatccctct 60  
tgtcttttga aaggctcttc aactgaactg gactccaaca tccagtgaag ctctccact 120  
catcctttta gctggaccct ctggggacca agacagcaga ccagctgcct cttctacagg 180  
gcagccctcc aaatggctgg ggccactgtc ttctctgcac tagaagacct ttctatggta 240  
gtatccttcc acataagcta tgacttctat tcccaggaaa gcctgatttg tctcctctaa 300  
atgcacttcc acttatctgt gaccctctta caatgaaatc agagagagat aaccctgatc 360  
ttctaactca gagcaagcaa gctcccagggt cttcagaggc cctgcagggc acacagatga 420  
cagcggatga ccagagggca catgccttgt ctaaagggga tg 462

<210> 23  
<211> 692  
<212> DNA  
<213> Homo sapiens

<400> 23  
tttcaccact atgtagccta aagttattcc gtcattcatg actatcctgt ctaaagagtc 60  
tgaagatctt tatttggtag ctatggcttc agctagtcca ttgctaagt tacctagagt 120  
ggttgacaga tttctaatta tacgttcatg agagggtact cccactatt gcaagagact 180  
tctgccaaac ataggccaaa attcatctcc ttggtttgca ggtacagttt gtctaactct 240  
ggaaaataat ttcaatgaac tacttcagcg ttcagaaaca ttggagttat aaatagaaag 300  
aggaagagcc acataaccta atagacaatt acctctcata tgccagtggg caacacattc 360  
ataagcccat gtgtgcttga tccagggacc acacagggtc cctgatggat tctgaaattt 420  
aaggctttgg attactggta acagagacat gttaaagtac atgtcttcag tcttgagtag 480  
agtgtaatca gtctgatttc tttttttttt taatgagaca aacatcagggt aaagaccttg 540  
acaagaagga agagaaatcc cgagattcta taatcataat aatcgaattg taattgctag 600  
tttaagtagt ccttcaaaaa tacatctcat tcctgacagg ataaaacaag ttttataaaa 660  
tatattatat tctgggttca ctagggaac ac 692

<210> 24  
<211> 669  
<212> DNA  
<213> Homo sapiens

<400> 24  
ccttcctcat ctttgctgct ctctgctgac aatttaaaaa cccgacatgt gttaactctc 60  
tccttgctctt ccaaccacc cacttatcac ctacagtcca tgctcccagg tggcaagcag 120  
agaggactgt ggtttgatga gtcatctcat gccgtggctt taattactga taagagcttg 180  
attatacaca ttctcaaaag cattggaaag ttaaaagaaa gtcccttttag gtagcagtc 240

atgacaaatg cagttcatga aatctgtgtc cttttcattc ctttctgagt aattcctctc 300  
 tgtctctatc aaagccttgg atactccatg gtttactagg cagaaactta tccatccaac 360  
 acagccacat ggatacagct ttgtgctttt agacaataac cacttgagaa aacctgacct 420  
 tttccccac tcttcattca gcttctgtcc tgctgaaaac aagaggacat cctgccacat 480  
 tgtcatctgc tctgccttac tcttgagaag tctagttggg aaaacaggcc ctataaagag 540  
 agacactgca atgccatggg gtgaggacaa taaaagtgtat ggcagcagag cactggagag 600  
 cagaggtggg gtcaccaact gcccaaatgg cactgtcccc tcagaactct tgcatttgct 660  
 tttaacgca 669

<210> 25  
 <211> 654  
 <212> DNA  
 <213> Homo sapiens

<400> 25  
 aatttatgac attatgacag tttgtcatta aagataacat tccaaagaga aatgggcatg 60  
 ggcatatatt taccactccc aaggaaatag ctaataaagt aatagagtac agattaaaat 120  
 aataaaatcc aaatttaatc catcacattg acaatgatta aaattaaatt taaagcagtg 180  
 ttgggaagaa tacagtgagc tgggtgtccat acacactgtg atgagagtgt agaaatctta 240  
 cagtcttacc agaaagcaaa tgtatcaaac actttcaaaa tgttcatact tcctaacct 300  
 gaaattccac ttttaagaat ttctcctaag aatatacttt tgtttaaaaa tattttacata 360  
 caaagatggt gatttttagta ttattttgaa agcaaaataa cccacagaat ctcaagtata 420  
 tgatccaaac aatggaatat cttatagcca ttaatttttag agatgaatat ttaataattt 480  
 aggaaaatac ctatgatact ttaaatttta aaaagttaca tagcagaaga ggccatattt 540  
 caatttttgc cttggaaaaa tatggtatca ctacagaaat gttgtagtgt tatcgctgac 600  
 aacactagtt atctaggata aagggatatt ctcatcttca tttcaccttt agta 654

<210> 26  
 <211> 687  
 <212> DNA  
 <213> Homo sapiens

<400> 26  
 ccaatatttg atcttttcta tctttaaaaa tggcagtttc atgtgtcttg atctaaaatc 60  
 ttaaaatcaa tctttcaatt ggataagagg cagggaaatt agcttggaag gtaaatctat 120  
 tatccagagg caaaatttca tgggctttga taaagggtgga tatttttcga taaggaggaa 180  
 agagtaaatt ttactaacat actttggctt ttgttcagtt ttcttaacct ctattttcgc 240  
 tttattattt atttttttgt tttactcttg ggaaagcaaa ttatttgttt tctcacatct 300  
 tttggggtcc aattttgatg attctgatct tttttagttg cttgacctgt agaccctcta 360



cagaacattg cagggcctct tctcagagga gcagcgggtga tgagcttagt ttcctaggct 420  
gggactgttg cgctggactt gacaggtgaa ctgaaaattg cagggataag tacacctatt 480  
gagaacaaac atccccatctc tttatcaaag ctcttcattg gctttggaaa actgctgtag 540  
gcctaaggaa actaaacttt ctagggatat tctagggttt aaacatatga gaaagagaaa 600  
gacgtcgggt cttatttaag agagtttatg agaccttatc cttgaaatag tcaaatttat 660  
aaatgacata aggctgtatg tgtagtt 687

<210> 27  
<211> 622  
<212> DNA  
<213> Homo sapiens

<400> 27  
ataaaataca gatctgattg tgtcactctc ctgcttaata tttgtagttg accctccac 60  
tgctctcatg aaagttcata atccttactg tggttgtaaa tgccctttta tgatctgtcc 120  
cttgcccatt tgtgtacact catcttgtgc tactctcttt cttcatcaat atgtccacc 180  
atactgtcat ctttctgctt atttttttta aaaaagtatg gaacatctct tccccttat 240  
gtgtcttatg caacctgtca gacaaaacca catgttatat tttctcaaca cacaatttta 300  
tttcagggtc ctgtgccctt tacaaatcta ctaatctttc tgtctggagt gttctttctt 360  
ctcctggcca aattctaato atttgtcaag agtgcaacag catcatttct tctgtgactc 420  
aattctccaa gcatcgtatc ctctgtgttc ctatagcact acattggatc ggtccataac 480  
aattctgtca gtgtattata agaacttatt tacaggtttt gtctcttcta ctatggcgtg 540  
agccttttag tcatatgaat tgtgattttg tatatttagc gcctaccatg gtgcttaatt 600  
cgtggtaggt gctcggtaaa tg 622

<210> 28  
<211> 684  
<212> DNA  
<213> Homo sapiens

<400> 28  
ctattggttt aataaattat ggtataatca aataatgaaa ctctatgcat ttgttaaagt 60  
aacttttcaa aagaatatct tgtaacatag aataacagat cctagtgcac taccactct 120  
ttgggcttta tcgcttttcc accatcatta tctgcatcac tgctgcagg ttttctacac 180  
ggccagggtt ggtctctgcc tgctcaatag tcaagtcaaa agaggcagga aattaacacc 240  
ctctggaggc agcctttgag gaatgatcca tgggaggtgg agtataaata cctcagctct 300  
gtttcctcta gagatataac taaggaatgg gttttacatt gtttctcaga gtttctcaa 360  
ggttttaaac ttcaatcacc cacagggtga tggggcttta tcatagtata catcctttgt 420  
ggcttccctt ccttcttgtc tcaattctcc attccaaact aggatttatt tcttttcct 480  
aaaacaaaac aaaatgttta acctgaaacc cttacaaaac acgtaaaatt tatatttaaa 540

aaatctaaat atttgaggag agaacgaaac ctaagtatat gccaggtat aacacgattg 600  
 gtggagatag ctttaaaaaa gttcctgaaa aatttagttt ttaaaagggt accctagtag 660  
 aaggtagactt aactgcctaa tttc 684

<210> 29  
 <211> 731  
 <212> DNA  
 <213> Homo sapiens

<400> 29  
 ctggcttctg agagcctcct ggtaggaag gaagtgttc tctttccact gcaagcttag 60  
 aaagccttcc aagttctctc cttctgcagc ataaagagac aataactcag aggaaggtat 120  
 ccccaggagt ttccagacag ctgcacagat ttaagtgcag aaatctgagc agaggtatag 180  
 tcctggcatt tacatgaaca cttttcagta gcaggaagaa taaatggaaa gagagctaca 240  
 gaaataccag gggcgaagtc ttcatctgaa agtccaatct ttgatcaaga gctggtagga 300  
 agtctgagaa tttgtatcag cagtgtattct aggtgtctg gtctgagtaa ttgggatcag 360  
 agcaacagct gatatcatgc ttacctgtg ccaggctccc ttctaagggc ttcttgaca 420  
 cctgctcgtg tcagtcctca cagcaatcac atgaggtatg ttctgttgtt gtctccttgt 480  
 gcggatgaag aactagga cagagaaaac tgccacagg tgtacagctg gggaggccag 540  
 agccagaatt cagacctggg gtgtcttggc tgatgtgagc tagtgtggc cagcatggga 600  
 cacagagga ggattagctg gagaagcagg acagaggga agagagcga gatctccgac 660  
 agtgctgggt cagagacact ttctgagcc atgattaac ctgattatgg gacatgtttt 720  
 agcctgtcag a 731

<210> 30  
 <211> 642  
 <212> DNA  
 <213> Homo sapiens

<400> 30  
 acagtgaagc gagatggagt cacacctttt caaaaattt aacaatcatc atcgatatgc 60  
 acagccttca tgtgtagtgt atgctcccag ctacagctgt agttaccaa tctcaaagca 120  
 agtaaacagc aagattccac actagctctt aactggccaa gctatatttc tataactaga 180  
 attgctatgt gtggatttcc ataagttata ataacacgat aagaccatt tatccatgta 240  
 ttctagtac tttttcttcc tatagcaaaa agaaaaatac atctttcacc atttacaagt 300  
 acaaatttca aggagaaatt ttaaaaggag agtaacaaac tgtcctgagt tgcagcaaga 360  
 ctctgagag ttccatttcc tgggcctct gctgctgtt tttggcattg aaccaggaa 420  
 tcttttctaa agcacacaga aatcttgcaa aagaggccat ttctagttag gcttttgtcc 480  
 aactgtctag ttaataaat taaattctta gattacaaaa tgtgttcaa aggtttaaca 540

aattgaaatg tccttaagta tttcaaataa attaaggaag aattcccatt cccatagtct 600  
tctactttcc tcttccacac ctatgatgaa tgtcctgaaa ag 642

<210> 31  
<211> 592  
<212> DNA  
<213> Homo sapiens

<400> 31  
cccttttttc tgctttcagt ttgatttgat tacaccttac aggcttgga tgataagttt 60  
aaaacatatt gaaggtttat gtacttataa aaacctcatc attccctaaa gaaaaaaaaat 120  
ctcaatttgg tttagtgtca ttgtagtctt gctttctaca tcttactaat gtctcattta 180  
tttattcatt ttgctctgtc acatttagaa tgattttgat gggcaaaaat catggtagtt 240  
acaaacagcc ctttaaaact attgttatac tttgttcagt ggattctggt agaggcttta 300  
aggtaattat ttcttttaaag catttgttaa atatacctcc tactgtagtg cccttgggaa 360  
caggcaaaat tcagaactgg cctgctagca gtcttaccag ggttataaaa gtaagattat 420  
tatatataaa acagcattaa ctcaatgcgt ggtgtgttgc agctggcaaa caacctcgt 480  
ccccaagctg ctaaattcgt ggtcttatga atgtctccat tgctgtgttt gctgtaacaa 540  
gaagtgggag ggtgttcccc agtagccttg actgtttacc aatgcacact cc 592

<210> 32  
<211> 485  
<212> DNA  
<213> Homo sapiens

<400> 32  
ttatttggtg ttaatttcat aggctcaaag gtctaagggt cccccctgtt gcggttgcc 60  
gtggttctct ttgctcctgt ctgccctctt gggccaata cctagtattg tgcttaggat 120  
tcacaaacgc acaaaatact tactgagcac ctactctgtg ccagggtgctg tgctatatgc 180  
tgagaaaaca atgttaaaca agatggataa ggttttcttc cttatgggtg ccatagtcta 240  
gtggcaaaga caggtaataa tgactcagtg tattctacta aggacaagca tatcgtgcta 300  
agaaaacctg tgtgggaatg ggtcagggaa ggtatccttg gagtagcccc gtttgaactg 360  
ggatctgaag actgagagtt atctaagtgg ggagagcatt gcaggcaggg ggatcagcat 420  
gtgcaagggt tctcagaaag gagggagaac aatgtgtaag aaatatcact gtagttgcaa 480  
cccag 485

<210> 33  
<211> 695  
<212> DNA  
<213> Homo sapiens

<400> 33  
tcattattat aagaattata agaattctga aatattagcc ttaaaataac caagttaata 60

```

aagcttaaac tttttatgga attatccatt tctgttttga aaaatactga actcttttca 120
aatactattg cttgttcact taacaatgat tacttgaaca tagttcagct aaagctttta 180
tgatattcac taatctagca tttattttcg cattgttttc caccatcact aaagtaatta 240
ctacatgttc accaactaat tattctgatg gtgcattaag aattgatctt taccttaata 300
ttttatggta tcaagtgttt ttgcattcat caagaatatt ccattttgct tataattttaa 360
tgatgagctc tagaatatca tctaacaat atctagcaaa ttataaatat gtcatttttt 420
aggtaaaata ttttaagagta tgtagtgcta tatatttagt tattttaaat caaatactta 480
atgtttatac tttttaattg atgtacaatt ttcaattctt tagaatgcg cttatgaaata 540
attgccctta ttatagtttt ataacaactt taatatatct tctgtatcta tagcagatga 600
tttataaaaa tgcttttctt tattaataac tgtctctatc tcaagttctt catagtgagc 660
tattttttct ttttgatatt ctgtagagat acata 695

```

```

<210> 34
<211> 655
<212> DNA
<213> Homo sapiens

```

```

<400> 34
aggcagtaat tccagtaatg tgatgaagta gcaagagata agtaagtcca ggtcagtgaa 60
gacttcgtgg ggctgacata tgaactgagg aaatgccac ttttgactt tcagttaaga 120
caaaaataaa ctacactctt ttttttttcc caggatatctg ttactttccc tattttgcaa 180
tacttaattg atacatacaa tctgtcaact cttctctctg gacctgcgca tacactgctc 240
catctgcctg aaacaatctt tccctggcca accgcctacc cactgccacc ttggagaaca 300
gtactcata gtcaccctca gattatatcg ttttctcacc catctcatcc tcttcttcc 360
cgtttcacca cctcccttca accttgggtg gctttgccc tctgtctgct tgacaggaca 420
cccctattgt tacctttgac tggactatta gatgacatct cagttactta ccttttatgt 480
gctagaatta atttcctagc tggagtgtgc cccatgacct gaagctgagt gcctgctcta 540
ccatgcaaga agctctattg ccgaggccta ggcctgtttt gggggcttct ctagccaatg 600
tgcaatgtcc cattcctagt tgcattctga aatataacat ctgagttcac agtat 655

```

```

<210> 35
<211> 506
<212> DNA
<213> Homo sapiens

```

```

<400> 35
tttcgaaaaa acgtatatga aagatttaaa atatgagtta tgatgtcttt ttttatccca 60
aatctgcttt aattatcatc ctatgagaac atttttggac atgcatgaac atacaagtgt 120
tctatgtacc ctccacagg aactattaga ggttaagcat cattcagcca aaaatgacta 180
gacaaacttc aatgagagga ctgatgtgaa catttaaata tatatcaaga tagatctaag 240

```

gttaaaaatt attgagaata aaattggaag aacaatgtat caacgttatg ctattcaaaa 300  
 ctagaataaa tgcattgtaa caatgggaga agaagggaaa gtaaaaaaga caattgtaaa 360  
 agcacgttat tggatagcaa atgtatggga agtaaagtac acacattaaa cttggcaaac 420  
 cagcagataa gaagttacat aagaatatag atggctaata acattttata gtataaatag 480  
 gccttaaaac aaatattaaa accttt 506

<210> 36  
 <211> 645  
 <212> DNA  
 <213> Homo sapiens

<400> 36  
 ggccgcccag gtcagggaaac cgtgggtctaa gtcccagctt tattcttagt tggaggagtg 60  
 gccttaggta tgtcacaggg ccccttaggc cttttggttg tcgttttcat aaaaggcagc 120  
 ttgtcttgct gctgacaatc atctttgaga gtgttagact taaatgagat cctgcagtag 180  
 ttttcaccct ccacaggtag caaaatcttt actctaaaca aattgtactt gattccttga 240  
 tgctaaaaca aaagaaaaac ctggaatctt attactacaa acatattcta taagccctca 300  
 tgtataatct ttacttttct tggagcccct cagtaagaaa aacaaaacag cttttaatac 360  
 aatgttttca caatggcaaa gttcaaacac agacaaaggt agaggcaatg gtatgataaa 420  
 gccccaggca ttcacacccc agattcaata attaccaatt cataatcaac ccaatttcag 480  
 ctctccacct cacacctcac tttttaaaag acagatcctc cctcattaga ttagttcatt 540  
 cacaaatatt ttatatgac ttgaaaatat aagtgtcctt ttaatcattg tgatatcaaa 600  
 ttcaaaatta acattaattc tcaaataaat agggctatct tgatg 645

<210> 37  
 <211> 563  
 <212> DNA  
 <213> Homo sapiens

<400> 37  
 ttgaaagta catgtataac taatctacat ctagcatcaa ataactacca cttcttctt 60  
 cctgtttata tcattactgc cttttatttc atttatccac atgctataat cactcaatac 120  
 tttgttacta ttattgtaaa cagttatctt tcagatcagt taagaaaaat aaaacttaat 180  
 tttaccttaa tatagtactt ttctaagtct cttccttttt tatgcagttc tttttgacat 240  
 ttctcatagg gcaggtcagc tggcaatgaa ttattccagt tttgtttgtc agaaaatata 300  
 cttatttctt tgaatttgaa ggataatctt gctgaatgca gaataatagt ttggtagctt 360  
 tttgtttgca acacttcacg tattctcctt tctttgtgtt tgcattggtt ctgaagagaa 420  
 agataatgta attcttatcc tttttcctct atggataagg tgttggtctt tccccctctc 480  
 tagcttcttt caagattttc tcttctcttt ggttttttgc agtttaaata tgatatgcct 540

gggtggagat ttggatttat tat

563

<210> 38  
<211> 604  
<212> DNA  
<213> Homo sapiens

<400> 38  
acttctaact gctggcttta atttaattta atttaattac agcattttcc acacatgccc 60  
acaggctctt ggtaatagtt gcatttttaa taaatctaata atataataat gactttgttt 120  
ttaattttcc actgagagtt ggatcctgag ttgaacacag agctccagac aggggcgtct 180  
ggttcactcc atgtgattgg atttcagga accaaggggc tcctaattgg aaaatagctg 240  
tgctttcacc ccctatcccc acacacctgt gtttaatgtc ctcagcaagc atcccatagg 300  
acatgaaatg accgcttggt tcagtcaaaa tgatcaaacc agttgagcag gcattcctca 360  
ggctggactg tgaaaggaaa atggaggtaa gcgagcaatg cctggccaag accattatac 420  
aaagagactc tatggacagc actctggtgg tggcctttac ggagtgacct actgctctct 480  
gcctttatcc acaagtcact gggccaactt agaactgtaa tcaaacatag ttcaacaaa 540  
ggatgaattt tatgactact gatttctcct ttgcaaagac cgtgggtgat attcatcggt 600  
aggc 604

<210> 39  
<211> 687  
<212> DNA  
<213> Homo sapiens

<400> 39  
ctcgagcagt aacctgtgct tctacaatta tgacaccac tccagggata gtcactgcca 60  
aagggtagaa ctgctggggg ctcatcgac tcacacagac taagagtgtg gcatctccca 120  
gttatgcggg catcagggca acatggggag aacagtggca ggcacataag gccaccccca 180  
ggtacaatgt ccagtgcagt tcacgggtag gtaaatctac tctgtgtccc cacagacca 240  
tagactccca gggggcaca agtcaatcag ggcctgacct tggtagtgac atgtgttatg 300  
tttgcaagg ctgtgacagg taccatccc acagtgggtg taccaccaatg ttgctctatg 360  
cactgtggca cttgggctgg gagtactaca tgttccccac tagccagccc catcataaac 420  
gctatgggcc agccaggggt tgggcacacc atgtgtcttg cagcatcctt tgtccaaagc 480  
tgccatgttg cattccaggc atcagccatg ggaccccaa gtctccaacc atgtccagtt 540  
ctctgcagac acaagatgta tgtgccaagg caagccatcc gcagccctgc tggaagggca 600  
gtgcatatcc aatagttgga aacattgggt acctagtgtg aggtgtgggc ccagtccaca 660  
atgcaattgg agtatgttaa cctctgg 687

<210> 40  
<211> 550

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 40

aatttttttt cactacggaa actcgtttgc taatataaat gcagactttt tttaaaaaaa 60  
agcttttattt ggaaacatga tgaaaaatgt gatgtattaa tacttactga tactccaaga 120  
aaaaaataat aaaatatatta gaaagctcct cccatcatctt cctttggctt tttaactcta 180  
ccagatcttt gagaatgcat attgttgctg gttaaccaga tgaaccaccc tttccttact 240  
agttctgcaa gattcaatat cattcatagt ctccagcact ctagagtaat cattactagc 300  
tgttaggaaa attatggtat ttcctaaaaa ctttctttgt gacaagtga taaacaaaa 360  
ggattaaaaa aaagatgttc cagtttggga aaaataatgc aatgaatact gcatctgatg 420  
caccatttaa gaaagagaga aaataaaaat gctcatttct aattgtcctc atttcagcag 480  
cttcccaaat attcttctat ttctttcttt ttaagtaatt accacatttt catatttgct 540  
gaatcatgaa 550

&lt;210&gt; 41

&lt;211&gt; 617

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 41

cccgagtgac agaagccatt tcactgccag agactcttag cggccttcag ttctcttgag 60  
ctggagccac tgggtcttgt atgaaagctc accaggacat ctcatgtgga cctcgggcat 120  
ctgagccggg accatcctat tacaagtgcg gaaaccagat cattaatgca gagctgaatt 180  
caaattgtta cttgctagct taggaaagaa tccttgga tccaacatat tgtctaaatg 240  
gatcagttaa tcttactatg tgcattctac atacccttc attgtttggg cttaaataac 300  
ttttctgctt tgtctggttt aatttcattc aatgtggatc gctggaagaa tatgatgtat 360  
gtttttagaat agaaacagtt ctgagatgaa gttgagcaca atttcctgtt ctagttgcaa 420  
ttaaataata atatagcatt tgacataaaa tagctggccc gatataattta gagtacaagt 480  
taagtgtcat ccccttagaa ttgggcattg actccgtaga attccccttt gtacaagggtg 540  
agcaaatgta tattttgtta aaaataagta tctgactgcc aaaacggaca gaaagctctt 600  
tgccatatgt gttttca 617

&lt;210&gt; 42

&lt;211&gt; 653

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 42

cttttttaatt ttgtttttgt agcagttgtt tgtatccatg tgtgttggtg cccatatgta 60  
ttgtttgggg tttggttatt ctctcaaaac caagttaccg taaaagttt gaattttagt 120  
atttccttat tgagtagtgg gaccgtctag actgtgtgct gactcttact aaagtcattt 180

gtttttctta cccgtggaga ggtgtattct tgaacccttt aaacgggtct ctactttggc 240  
ctaagaccat attagaaaac ttttttgaag tcacttatta tatgccatat aattaaag 300  
ttatatggta tattctccca ttacatttta gccacaatgc ccgtatatta aataagcaaa 360  
caaactatat gtggcaatta aaacttaaaa aaaaaagcct gaattggctc ttagaaatat 420  
ttaatcaagt agtatccact agaacttaac atttcacctc gtggatcatc acacacaaaa 480  
taccacaacc tgctgtcatt cagggtccta gcaggaacag gtagcatcaa ataggataat 540  
tgatgagagc ttaagaaaag aactatttac aaatatgtgg ccagattagg ggaaaccagt 600  
aaggttggga atgccgcca ggattctaac aagagtgaga atctatttct act 653

<210> 43  
<211> 642  
<212> DNA  
<213> Homo sapiens

<400> 43  
tccatgtaac attgatgagc acagttttct cttctgtagc aagcactcct ctgcctaatt 60  
catatgacta aaacagtgtc tctcaaacat atgggtctcag gaacccttta aaatcttaac 120  
aactagtaat gaccccaaaa aggtttttta taatatgaat tttatatata aatattttat 180  
tggaagtcca cttttatgaa aataaccttt tttcaaaaat ttcataagaa aaaaatagta 240  
ttattttaca tatttgaggc atctttttaa tgcctggttt aatagaagac aattgaatat 300  
tcatgtcaac ttctggattc gatctgtttc aatatgtgtc tttggttgaa atacatgaag 360  
gaaacttggg atcatcagac atatagttag aaaagggtgg agtattttaa cagccttttt 420  
ggacaactgt ggacattgtg ctttgatatt acaacaaaac tggagaagtg gtaggttcta 480  
aatgattagt tgcaacatgg aatctgaaac cacatcatga actatttgta atctggcata 540  
ttaagatcta tttatctatc ttgcactttg aatgggatcc tttgctcatg catctttttg 600  
taacatgaat catctcaaac acgttggttc attgagttat gc 642

<210> 44  
<211> 674  
<212> DNA  
<213> Homo sapiens

<400> 44  
aattaaaatc cctgcagtca aattagactc tgcattgtctg gggatattta aaaggataat 60  
gtataggggt tgccatggta actcatcaag tggtaattct gtacctttct gaggtaaac 120  
cttgaaagga gaagacaagc aatttgggga gataacagca ccagaaattg agttcatctg 180  
taacttaggc tctctgtgag tttgtttacc agctattcac catgtggatg aaaaacagta 240  
aaaagacaaa aaagattcac atttcaaggc tccctaaaat tgccaattcc actctatagc 300  
tgattctcag cacaggagga aatgggacta gaatgctggg agatgacact atcatcgaac 360



```

agtgagctcc aaggagaagc ctaattgtta cttctcaatg gcagaaggcg ggtgcttccc 420
ccggggcagg attctgttta atccttaggt tagagcccag cttcaaccca gtgtcacagg 480
tcaattacca ccctccaacc ctgaggggcg acatgaacca tactcacgca ccggcgcatg 540
ctccctcttc agcacctctt gtacattcag agctcctgca tgggatgccg agaactcaca 600
cccttcagg gctgctgaag atcatatgac tgatcatcaa ctttgatttt tgacccatct 660
gtcaacaacg acac 674

```

<210> 45  
 <211> 609  
 <212> DNA  
 <213> Homo sapiens

```

<400> 45
gcctaactga attataaccg cgagtttgca cagtggtag catagctgat gagatgcaag 60
caaaaaaaga gtattgctga cctaggacca tgaggaaaaa ccaaatacaa attagtcaag 120
ttggaggaca tttgttgaaa actccacact tccatgaggt ctgtagcctt gaggctatca 180
gtgccgacac agaacattct gaatagttca atgcctcttt ctgttaaaga ggagacgct 240
cactctgccg ctcaatcttg gacttgtttg tgcacagagg tccttgctta tgtaacactc 300
gcttttaact ataattcaca gagtcctttg aacacataaa gggaaagcca ctttcgctcc 360
tgttaaggat gtataagcac aaaaaatgaa cagtgaatta atcctagtgt tttatacatt 420
tttttttaaa aaaagaatct aagccagaat gaggttactg cctaggcaaa gaagaagaca 480
gctcatcaca ggtgagtgtg acacgttttt catatgtaca aattaagcag cctgaaacaa 540
aaggcactca aaaggtaaaa gaataccagt ccaccctct gatttgtaa atcaaagttc 600
tgtcaactg 609

```

<210> 46  
 <211> 522  
 <212> DNA  
 <213> Homo sapiens

```

<400> 46
aaaaaaaaaa aaattcaggg gaaaaaagca attaaaaaaa cataactata aaaataatac 60
aaattacaaa acaaccattt acatagcatt tacattatat tagttataag taatctagag 120
atgattaaag tgtacggagg aatgtgcata ggttatatgc caatactgcc tcattttata 180
tgagggactt gaacatagaa gggttttgga gtccacagag gtcctgaaac caatttcccc 240
ttcccatgcc tgggatgact gaattatata gcagcaaaaa tgaatatact caagctatat 300
gcatgagtct cataaatata atgctcacag aaaaaagcaa gttgcagaag ggtaaatacg 360
gttgatatat aaagggtgta aacacagaac tatttaatga tatacggatg cagtaaaagt 420
ataagaaatg tatgcaaact tacttaaatt caggggtgtg gttacttga gtaaggcgaa 480
tgtttgggat gtcagtaggt acctgacaaa tggcaactta ac 522

```

<210> 47  
<211> 681  
<212> DNA  
<213> Homo sapiens

<400> 47  
agctaggggtg ggcaggagtg gtctctgaga ggtgacattt gagctgagac ctgaatgaca 60  
agagaccaat gtcagctctc ttttaagaaag ttttcctttg ttttagtggc tctctccata 120  
ctcttatttt aaactcactt aacatcaata taaaagtgtc ctttgcagca ggacactttt 180  
aggagggtctt gagcccctct cccaccagca ctcatctgtg tacaacaag ttgttgctag 240  
tgggtgttga gctcgttttt cccaagcttc accttggcat taccagatc tgttcaacct 300  
tgggcatctc ttctccag ctggatgtc acccaacttg ttctgcctca gtttctggag 360  
gagcctgact ctatttttgc ccccttgaa agaaagtaca ggactgggtt gaggcagctg 420  
ctcacactca ccagaggcct ccatatcttg tagccacac tggtgccat caagagctgg 480  
cagtcttgag aaagcagaaa gcagatggtg aggtagaagg agcagtgat atggaagggc 540  
acaaaacaga gggatgaagag gccacacacc agtaggatgg tccggatgga cctggctcgg 600  
gctgtgttgc ctgtcctcat gaggttctcc tctggcttga tcaggctcct gaccatcagt 660  
gaatagcaca ccaaagtgac c 681

<210> 48  
<211> 548  
<212> DNA  
<213> Homo sapiens

<400> 48  
ccagggggag gggggcacgg gctataaacg ctcgcccgca gcggcgccgg cagagagccg 60  
ccgagcccag cacagctgcc ctctggaccc tgcggacccc agccgagccc cttcctgagt 120  
tccacaggcg cagcccccg gcggtcgggc ggaggggtcc ccggggcggt gccagggcgc 180  
aatcctggag ggcgccggg agggaggagt gcgcgcggcc atgcacaccg tggctacgtc 240  
cggacccaac gcgtcctggg gggcaccggc caacgcctcc ggctgcccgg gctgtggcgc 300  
caacgcctcg gacggcccag tcccttcgcc gcgggcccgt gacgcctggc tcgtgccgct 360  
cttcttcgcg gcgtgatgc tgctgggcct ggtggggaac tcgtggtca tctacgtcat 420  
ctgccgccac aagccgatgc ggaccgtgac caacttctac atcggtgagt gcgggccgct 480  
gcgcgcacc tgctgccgtc ccggggggct ccgaggccg agcggcctgg ggcgcctct 540  
cgcgacgc 548

<210> 49  
<211> 695  
<212> DNA  
<213> Homo sapiens

<400> 49  
aagtcgcctg tctttgatct ggtagccagg ctgtgatggc tagctttagg atattttccc 60  
tatattttctc ttgctgtcag gttaccctt ggtatactg taattgattt cccagttag 120  
agagtttaga tgtggacagg ggaagtacaa actacagctt agtgcaagat aaaccaaggg 180  
tgtaattatc aagttgtact tgaacagaaa tattacccaa taggatttcc aaatgaacag 240  
gatggcaaag agttctgggg tgtggaagtc agagtaggtg ccaaaggatc tagatcaaag 300  
gggttggtag atgagcaggg atgggtcaga gaaatctagg actgttaaag caagcatgac 360  
ccaggccatg ttctgagggtt ggtaaagtga attatagaag gtgagaccaa atgtgagatt 420  
gtgagatttt aaccacccca aagaggaggt atgtgcctca ggcaaagaaa atgggaaaaa 480  
aaaaacatgg tatatggcat atttgaggag caaagataag ttcattgtca ctagggcaga 540  
gcaagggata agtgaatggg gtgagacaag attggagagg ttaacagtgg ccaataacaa 600  
gtgataaaaa taattttcaa atgagagcag cccagcactt ataaagtggg taatgtgcac 660  
caagtactgc tttaagttat cctgcagtat tattg 695

<210> 50  
<211> 586  
<212> DNA  
<213> Homo sapiens

<400> 50  
gcctccaacc gatatttctg tctgttgctc tgaccaggta ctgggccatc accaatgccc 60  
tgtagtatag taaatgggcc atctcaaatt gtatctctat cccagtgtc tctcctaga 120  
cctcttgac cacctactcc acatgtaaga cctctacat tttggttggt ttgttcatca 180  
tcttcacaca ttgcccaaca agaacatcca gaagccatca tcacagcacc actgccagg 240  
tcacacagc tctctttct tcctcaacc cagcctccat gagaggcaaa ggcgctaac 300  
tggctctcct ctgcttggtt atcacatgaa aatcaagcat gcttatagtg tcctagtaca 360  
acaggaaatt tactttcaaa caaggaaagc cacagaaacc ctggggatca ttttaggggc 420  
ttttatcatc tgctggctgc ctctctttat tgtttctctg ccagccaaga taccaccata 480  
ttaagacatc ttcatcttgc tgagcttttt ttttttttc tttttgatac caagtctcac 540  
tcttgtctcc caggctagaa tgcaatggta caatctcagc tcactg 586

<210> 51  
<211> 234  
<212> DNA  
<213> Homo sapiens

<400> 51  
caggcgcctc aactgttcca caaaccaagc ctgaaaccag aactccaact tctagtctga 60  
aaagcaaagt ggcaacctgc aaacacctg tggcccaag tagtctcacc caaccttggg 120  
gaagaagcag aattcaagct gtaactgcct gttggagaga gccaacctc ggcctctgtc 180

ctcgaaaggc agcaccaaag tttccaagt ggaatcaa at gtcagggag gatc 234

<210> 52  
<211> 308  
<212> DNA  
<213> Homo sapiens

<400> 52  
ctgtacctgt cacagttatc aaaaatttat tcattcagaa gtctttgttg aacacctgtt 60  
acgtgtactg agcattgtcc taggtatttg agatacatca gtgaacagag gatccttaac 120  
agacaatata cataataagt tatgtaatag cttacaaagt gacagtacct ttgggaaaaa 180  
ggaaaggtat tataggataa agatgatcaa tgaacaggaa gtttgcagtt ttaaattgag 240  
tggctctgggt aaggaagatc atacctgaac caagacacaa aggaggtag ggaatgatga 300  
gccctgca 308

<210> 53  
<211> 584  
<212> DNA  
<213> Homo sapiens

<400> 53  
tagcagagca ggtgctagtg atatttgtag aacagggtct gaatgaatgc atgaacaaat 60  
gcatgaatgt ggaaatgaaa ggggatgcag atggagatga tgcagatgga gatgatgatg 120  
cagatggaga tgatgcagat ggagatgatg cagatggaga gcagtggcca tgcagagtct 180  
ttgcagacct tggcttggct tcaggctgtg ggggctctgc aagccaaggg tttgagttcc 240  
acctccagtg cttgccagca atgccacctt gggtagacct tatcttgcta cctggaaaagt 300  
ggggatgctg gcagccctc cctcctggca tctactgacac tgcattgcca ggggtgtgatc 360  
cctttgggta caggcggggg tggtagacct cccagggtgg caggtagcatt ttggatgaaa 420  
ggccaaggac gattcatagg agagcacagg agtcttctg tagccccagc aattccacag 480  
aacctgctgt gaactgctgg ctgctgccc taacttttcc ctgtccctat ttccactcct 540  
tggaggccgc aagaacaact gctggctggc cttggccact gcct 584

<210> 54  
<211> 560  
<212> DNA  
<213> Homo sapiens

<400> 54  
agtctttttc tttagggaac tttgttgttg cttcactata tagttgttgt ttcaacaatt 60  
ttgtgttgtt tcacagtttc actgtgacag tttgatgtta ggttgattct ttttcctcct 120  
ctgtataaaa gattatgtca ccagaatctt ctttcattac tttggatagg acctaaagga 180  
ccctctcaat ctgaaaatct atgctatttg ttatcacaga gcagttttct gctgtcattt 240  
ctttgattgt tacttttcta tttattcctt tttctcttcc taaaatgcca ttatttgtat 300

attggagtca tagatctgag atctgtgaat ttgctattca tgtctcatat ctttttgcaa 360  
 atggtttcca tgtctccaag tctttgttct ctattgtgag atattatttg tattgttttg 420  
 tccagaatat taatttagtt ctattcattg actattcttt ggttttgctg ttgaattttt 480  
 aaattcagga atagtgtgtt tttctttcag attatttttt tctgtgacct aattgcatct 540  
 tcttacgggg tcttattata 560

<210> 55  
 <211> 234  
 <212> DNA  
 <213> Homo sapiens

<400> 55  
 gccagggaa gccaaaagat tggacatcca tgcctccctc ctctccctc ccgactgcc 60  
 tctcttgatg gcggccagtg tggcctacaa gatatggagg cctctgggga gtgtgagcaa 120  
 ctgcctaaac ccactcctgt actttctttc aaggggggca aaatttgagt caggctcctc 180  
 cagaaactga ggcagaacaa gttgggtgag catccagctg ggaggaagag atgc 234

<210> 56  
 <211> 585  
 <212> DNA  
 <213> Homo sapiens

<400> 56  
 tccttggtca ttttggtgtg ctattcactg atggtcagga gcctgatcaa gccagaggag 60  
 taacctcatg aggtacaggc aacacagccc gagccaggtc catccgggac catcctactg 120  
 gtgtgtggcc tcttcaccct ctgttttggt cccttcata tcactcgctc cttctacctc 180  
 accatctgct ttctgctttc tcaggactgc cagctcttga tggcagccag tgtggcctac 240  
 aagatatgga ggcctctggt gagtgtgagc agctgcctca acccagtcct gtactttctt 300  
 tcaagggggg caaaaataga gtcaggctcc tccagaaact gaggcagaac aagttgggtg 360  
 agcatccagc tgggaggaag agatgccag ggttgaaacag atctgggtaa tgccaagggtg 420  
 aagcttggtga aaaacgagct ccaacaccac tagcaacaac ttgtttgtac acagatgagt 480  
 gctgggtgga gaggggctca agacctcta aaagtgcct gctgcaaagg acacttttat 540  
 attgatgtta agtgagtta aaataagagt atggagagag ccact 585

<210> 57  
 <211> 660  
 <212> DNA  
 <213> Homo sapiens

<400> 57  
 gtcacactga attagggacc acccttgtaa ctccatttta actcgattgt ctctgtaaag 60  
 gccagtcctc caagtacagt cacattctga ggtactgagg gttaggactc caatgtatct 120  
 ttttgagggg acacaattta accctaatag accacaatta aaatggaatg caataataaa 180

aactaacttt tattgagcat tcgtagtctg agtttggcat tgctcaagag tgccttacat 240  
taattaatgt aatcttcaca atcctatgaa ctcagtatca ttattaccca catcttacaa 300  
atgagtgggt ggagtccatg gcaagagtaa cttgccaag gtcacgctgc tggtaagatc 360  
agaaccagac tcaaaaacag tagtctaatt ccacagcaga ttccgtcaac aactattcta 420  
cacagtctct actttatggg gttcaacata gagactatct tgatgtctgc ggtagctgtg 480  
agaatgtggc tcagagactt ccatctatgg ggaactcaat caaccaaagg cccagctcc 540  
tgcactttga gacctgtcac tatgttatca ccgagccac atttcccatg ggctgcttcc 600  
agccaatgcc caaacaatgg caggagact aaggcatcct gttcctgggg agatgtggga 660

<210> 58  
<211> 643  
<212> DNA  
<213> Homo sapiens

<400> 58  
attctgtcct cttctctctg cctgcggccc ccatctcctg agcccagcga gctcagtgtc 60  
agttcactgt ttgctcctcc ttgctgcaga cacagaagat ttgggagcgt tcctgccgag 120  
gttggttaagg atacctggaa cagtgggcgg cctctttgct cccacttgc taggagtaaa 180  
gccgtttaaa aagacacctg agcctctccg ggttcctgct cctcactcaa cccacagta 240  
gatctggtgg ggaggttagg ggctcagtga atctgcaggt gcagcatcgt gtectcagtg 300  
tcctgcccc tgcttcacc cggtgtcgac agtgcacgg tccacccac gcctgccttt 360  
ccatcgcttc tcacagccc tgtgatcttt cctgtggccc tgctgtgctg gtgccctgtg 420  
aggtcctgtg gacacaagag actgcacggg ccacacccc agctgggtga gtectctccc 480  
tcctgggtac tctggacagt aaagaaagat ggacacgtgg gctccgtgga gcatgaggta 540  
gtccaggacc tggcgggcca caggtcctgc ctcctgctt ctcgtgccct cctcccttt 600  
gggtctctgc tccacctcgg taaacgcttc gttcccaccc etc 643

<210> 59  
<211> 670  
<212> DNA  
<213> Homo sapiens

<400> 59  
aatatgtcct aatattctag tagggtaaat tctttattgc tttttcttt ctagaatttt 60  
tcttatatta ttttcatat aaattttaga ataagtctgg tttgggggt catatagcaa 120  
taggtaaatt gattaataaa gtgatttggg gaaggtttca caatacattt atgaatcaac 180  
ttcgggagag tggttatgct tatgtttagt cattatattt taaaatgtga catatctttc 240  
catttgtttt aagtccttga tcaagcatta gttgcctcct ctgagaatct ataattaaat 300  
tcaagataaa ataatttttt ccatttattg acccattttt agcttacaat ttgttttcta 360  
cccttgtaag tattatgttt ggtaaattat tttttattaa tatctccctt acagatatta 420

tacgccataa ggaaaggagt cacagatttg gtaatagaga ctcaatacac gtttggttga 480  
 atgatgaaag cattatgagg catattttct tactatgttc acctaataat cttaaagtta 540  
 tcaagttatt aagtagagcc cattcacaag tccagatcct ttgattttta atcctgtatt 600  
 ttcccatatt ttcaatatct aataggggaa gtaacatgct aaaatgctat agttttgcaa 660  
 ttttatatct 670

<210> 60  
 <211> 662  
 <212> DNA  
 <213> Homo sapiens

<400> 60  
 aaggaaaatg gaaactagat gaacgtgaca atataagact tccaaatcca cgtggttcca 60  
 tgaaaatagg aaaaaccgaa tgccaaaggg caggccacag aaggaggaag accagcgcta 120  
 tgagcaggat ggtcacgtac agcctgggtca gtggcatcct ccgggaccca caaaggatcc 180  
 tgaccagcag gaccaggctg gaaccacaga gagccacaca taaaaaaatc agcccccta 240  
 ctatgatgaa atctgatgtt taacaccaaa cagaatcagc accactaaac aggaagtcac 300  
 agaaactcca ctccaggagc ctccgcagca gagacagggc ccagagcatg acacacacga 360  
 ccgctgacag gtgtaggggc gggggcggca gcgctaccag atggggcaca ggacgtacag 420  
 gcagcgctcg gtgctcatgg cgcttagaaa gctcaggctt gcaaggtagg aaaacatcat 480  
 cacagggctg aggattttga agatgggatg gaggatattg atgaggctta acggaaaacg 540  
 tataatgtgg cttctgagaa agaggaagtc ggccatggac aggttgaaga thtagatgga 600  
 gaaagcgttc ctgcgcatgc ggaagcccag gagccagagc acgactgcat ttctgtcat 660  
 cc 662

<210> 61  
 <211> 603  
 <212> DNA  
 <213> Homo sapiens

<400> 61  
 cacacacaca cacacacaca cacacacaca cacacacaca cacacgcag caccattta 60  
 atgggttccc tgggggcagg gcatcagtc cactcactgc tgggcctcca gggcctgcca 120  
 aaggggcaaa gtcacactca gacataaact cttggtttta gcaatccaat aaacagtcac 180  
 gaaactaagt gaggaaagt attagattga agggatttga gggaaagtcc catcaaaaag 240  
 taaaacttga tcccactcc acttcttgga tgagttactt aatctctctg gcctcagttt 300  
 ttacacat aaatagaaa ccatgagagg acctaccta ccaggctgtt cttaaagtta 360  
 atgagttaat tcctgtacaa gctgagaaca gcatctgata cagtatctaa taaagtcagt 420  
 tattattact ttattatta ttatgtactt ggttatcatt attttcattc atcaattatt 480

attctcttca cctcttttget gccacctgga gttcctggaa ccccttcacg gcgtacagca 540  
gggagacagg ggagggcaga tgccatttgc acagccattg ggactaataa gccccagcac 600  
ccc 603

<210> 62  
<211> 427  
<212> DNA  
<213> Homo sapiens

<400> 62  
taatgtggga ctaaaaaact attaaaaaat aatgacttca accttcccaa attaggatgg 60  
aagaacataa acctaaatat tcaaggaaac aggagcaaac cctaaataga atacacccaa 120  
atacattcaa tttctggaaa tgaaaaaaa aaattaaaaa tcttgaaagc aaacagagga 180  
aaaatggcac atttcttaca gaaaaacaat aatgtaaacc acagcagatt ttccatctga 240  
aaccatgaag gttggaagga aacagataat atttttgaag tactgaaaga acagaactgt 300  
gaactgtaaa ttcaataccc agcaataata ttcttcaggc actaaagtga catagaaaac 360  
attgtctaata gaaagaatgc taaggtaatg tgttgctaac aaacttacct ttaaagaata 420  
agttctc 427

<210> 63  
<211> 550  
<212> DNA  
<213> Homo sapiens

<400> 63  
acctctaact ttcttcaact atgtgttgat gtctgtcact gcttaacaag caaaatggca 60  
tcagaaagag ggtgaacaaa taaaggataa ttagggcta atgatgaatt cgaggtaaag 120  
cacatcaatg tttccaccaa ggtttttgc tccagtgtgg tagggcaaaa agatgtgaac 180  
tgaattattg gtactctcaa attaaatgta ttcatcttat taattcattt agcaacagac 240  
atacacaggt acatatatccc atatccgtag ttctacttat aaagaaaaat taaatccacc 300  
caactgtttt gttttctgca atatttttaa cttctgtgac tttttgtttt ttccattgct 360  
ttgaatccac aataggtagg taggagaatt tgaagcacca ttgaaatgaa gtattctaga 420  
aaagtatgca gaaagataaa gaaaatgcat ccatctctag aagtgcctac atctacttag 480  
caagtgtgaa actcacaatg aggatttagc ctgttagtat ggcacagatt ataaatagga 540  
gagtcgctgt 550

<210> 64  
<211> 556  
<212> DNA  
<213> Homo sapiens

<400> 64  
accttggcct cccaaagtgc ggggattaca ggcgtgagcc accgcgcccgc gcctaatttt 60



gtattttctta ttctgtattc ttttccttaa aaaacctttt gcccaaattg tatcaacttc 120  
aataccccaa cgctggaccc ctccctagat acagtcataa agcaaatgac acgttagacc 180  
acgtgctccg ctaagaacat agaacctctg gcctgggtga tacttggtgt ttctgaagaa 240  
gcttttcctg ggggtggagga ggaggaggag gaggaggaag accctttgag ctttaaaatg 300  
cccaggagcc atttcctgta atgggtggat gcaaagaagt aatgatggg gtaatgccac 360  
agttcatgtt catgagggcc acggtggcct gaaggacag taagaaagcc ctccgctcgg 420  
cacaggatgg cagggtggagc atacctctcg ccatgaactg cttgatgttg aggtggtagg 480  
ggctgaagca gaccaccacg gccaccagca tcagcagcgt aagcaggcag cctcgccagt 540  
ggcgtccttt cctgct 556

<210> 65  
<211> 600  
<212> DNA  
<213> Homo sapiens

<400> 65  
catacccact gagggagaat ggagaagagg gtggggttct gcttgcaagg ccctttgcac 60  
ttcaaatatt ttacagggaa ggggatggca gatgcaccct ctgccaaggg aagctttgag 120  
ggccagcatc acatagccct gtggtgaatg agagctggca gggtgacagt ctgcgaggaa 180  
ggaaggatgg agctccgacc cctttgcttt ctgaaactcc tgctgagaga gttggctcca 240  
cagccctggt agggctcggg tagctgctgt ggctgaatca gtcctctgtt atcaccgct 300  
cggtgccatg aagtggaaaa gcagtctctg ccctcctcgt tcctccaata agcccatcct 360  
aatcacctt atcatgctcc ttccacaccc tgagaaaaaa tggcctcgca gcagacgttt 420  
gaagtcaccg ggactggaaa agtctttcaa atggcacctg atttggtac atgcctgcag 480  
acaggtgaaa gttagtgcc ccatttcaca ggtgaggcca ctgaggttca gagaagtcaa 540  
tcaatgatgt gatcatgctc acacatccca gcagtgacca aatatgtaac attcatacac 600

<210> 66  
<211> 549  
<212> DNA  
<213> Homo sapiens

<400> 66  
cctgccccca ccaccaatac tgggtgccac gtaagttgtc tagtgaactg aggaaatatt 60  
ctctcatca actgccactc tcaaggcccc aagtgtacca tttggaggct taggtattga 120  
tctgccccac cggtcacac tggcacccat gcacaccttc agggacctaa ggacaggccc 180  
actctgcctg ccaactgtcat tactggtacg caaggactgg cctgcctagt gtctccatcc 240  
acagcaaagc attgccacag cccttagttg ttaagccact gaggagctca cagacaccac 300  
tcacactgtt tacagcagga gaaatcctat ggggcctata atactgtgcc caccttgat 360  
caaaacaaaa gtactctatg caactaacac tacagctata tctacaggaa aaagcctctc 420

cctacaaaag ccaatccaaa aacctaggag aagcaactgt cacaccaa at acacagatac 480  
 caacttaaga acataagaaa catgagaaaa caaggaaaca tggcattttc taaaggagca 540  
 caataactc 549

<210> 67  
 <211> 550  
 <212> DNA  
 <213> Homo sapiens

<400> 67  
 agctgggatt tctgctaact gatgtccagt cggatatttg atatctcaa tgacatgaaa 60  
 ctactactg ctgagcaacc ataggaagac actggccagc ccatccactc atgcggtgct 120  
 ggaacccttt ttttatttta aaatatttaa ttgacaaaaa ttgcgtctgt tcaagggtgcg 180  
 atgtgatgct tcgatctaga tatatacagg tatattgatt accacagtca aattaactaa 240  
 caaatctatc accacccatg attaccatca tgttgagggg atgaggcagt gaagacacta 300  
 aagatctgct gtcttatcaa atttcaagtc aacaatacag tattattaac acagtcacca 360  
 tgctgtgcat taggtcccca gaacatgtaa ctgaagggtt gtatcttttg accaacatct 420  
 cccagctct gcgatgagtg atggtcagca ttttccaaac ccactctgaa gactttgcct 480  
 gggttggtac atcaatatct cctgagaaag tacaaaagtc caggcccagt cacagaaatt 540  
 ctgatgcata 550

<210> 68  
 <211> 605  
 <212> DNA  
 <213> Homo sapiens

<400> 68  
 caaaatatac atgcatgtac atactatgaa atatgtatta tgtaattttt gtgattctat 60  
 gtataagtta aatgctttta tatttgcat ttaaattgat actgcacaac ataaaaatga 120  
 atgtgaaaat ttattgtggt aatttagatt tttaattttt ttacataaaa ggacatagaa 180  
 tagcaaagga aaaacaaaac aaacaaactg aaagacgtaa caagttgaaa aatagatcac 240  
 agataaagga aacattttat actttgatac acttaataga accttttgct tatattttga 300  
 actagagccc cacactttca ttttgacta gaccttaca attatataat caaccctgga 360  
 cactgaatta agacaaaagc caatatttac aaaaatggc accatagccc aagctattgc 420  
 tttgaagcta cattagtcc tgtttccagc tgtgagcctg aactccattt taggaagtga 480  
 gactggccag ggtttctgtg tagagtttg catttttatt ctctaggacc ctgcaagagt 540  
 ctacagtaat tgtagactca aaaatgtcag agattgctgc ttgtatttat ataatgcccc 600  
 atact 605

<210> 69

<211> 669  
<212> DNA  
<213> Homo sapiens

<400> 69  
tattttccta tctaccacat ggaatcagaa ctgtcttgga gatttatgca tctgaacaat 60  
aatattttaga acatcatctc gtctttgaca ccactttggt caacacaaaa tggctattca 120  
aactactctg gaaccctgtc ttgtcaacca atgcaggaat cttagttaat gtattccata 180  
aacacacgca ggtttccctt aagcacagac tccatgtaag acaagtttca tactttttca 240  
ttgtgaaaga tgcagggtact attggatgga tctgaagagt tggcaaaatg acaggaagat 300  
caggcaggct gcctgttttt aactttatga aatttttcat gttttattat ctatctactc 360  
agataaaatt aggtgggaca cttttttaat gcttccaata aataagaaaa atgtgcctgc 420  
agcatgaaaa atcctttgac tgccttgtgt tatttgcaac agatgaatct aatttgatt 480  
cagacatcag tgctataact aactagagaa ataaaatgga tgtctatgat ctctcttcaa 540  
ttatttagta aggatgaagt gtcaattggc taaaagtaat aacaccatgg ctgtacttag 600  
tgttacacct attaggtaga aatatacaca catacacgca tatatacaac agattaataa 660  
caccagaag 669

<210> 70  
<211> 537  
<212> DNA  
<213> Homo sapiens

<400> 70  
tcctgaagtc agatagtagg agtcttctaa atttgttctc tttcagaagt attttggctt 60  
ttttattctt atgaattttc gtgtgaattt agaaacagct tgtggatttt aaaaggaaat 120  
gtctgcttgg atttgaatgg aattgcgttg catccagatc actttgagga aatttgatc 180  
ttaattctat tgaattttcc aacaatagac atgatgtagc tctctgttca gctcttcttt 240  
gattttttta atagacattt acagtttttg gcacagaatc tgtatatggt ttgtagatt 300  
tatagctaag cattttatgt ttttgatgct gttttaaaat tttaatttcc aactggatc 360  
tgctgccata cagaaataaa acagaaatac agaaatacag ggtacaaaat aaacttgacc 420  
ttgtttcttt cactctagat agtattgctt attagtctca ctaagttttt ggtaagttct 480  
ttgagatttt tctccacaag caatcatgct aactaaaaat aaaaacaatt ttgtttt 537

<210> 71  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 71  
aaaaataaaa gttatggatc acagcagatc ataatagaga atagtccatc tctcctagaa 60  
aattttttaa aataaatctt agaaactgca tgggaaatac tgtaaaaaca aaggttattg 120

tcctcagcta tgaattagaa taaatttggc actagattat ggggtattcc cacaggaaag	180
taccttactg attttccctc tacccttctt gatacattat ggttgaaccc actgttatgc	240
aacacctgct tacttttggc ttaagggtca tagtgacaaa agagaaacct ttaaagaagt	300
catagtaaat gttagggaaa gggattttca atgcatggat atatttggca aggtaaacia	360
aaagttgcct gatagcaagg gaggaggcag gccactgtga atagcaactt atactagtca	420
atattgaaaa gtaaaagcag ttgaatggtt tcaaagtata taagaatata aactgattgc	480
ttataaaatg ttttttaagt agagactgca ctttaatgtg agatgaggcg gatctatata	540
ttaattttat atacgcaaatt gatcctactt acattcttga aaataatttg actctttagg	600
tgaaccaact gaaatctcat ttacactggt gatttgccta gtaaataatt ctcttttagta	660
tgagaaaatc aaagaagttt gaagtggac aaattctaaa ttactagaat atgatttaaa	720
tggctaggag aatattataa ggggtataaa acagaatatt aatccaaata ttttaagatgc	780
taattctggg taaaagctat ttttgagatg acatgaattt tcaaaatact aaaattttta	840
aaataatcat ttccacaaac ttatttaagc tgtgtgtaat gtatgtaaat actaagtaat	900
atgttattca attttaggaa ctttatgtat gttttcatac tagtattaga aaataattct	960
gaaaggaaga tgaaaatgaa aatattcatt taggttaaac	1000

<210> 72  
 <211> 1000  
 <212> DNA  
 <213> Homo sapiens

<400> 72	
atgatattcc tattggatgg tgctaactctg gtgcagggtt tcttaacctc aggactactg	60
gcattttggg tcaggtcatt ctttattgtg tagggctgtt ctgtggattg tagaatggta	120
agcagcctcc ctggcctcta tccactggat gccagtata cccgctccag ttgtgaccat	180
cagaaatatt tccagataaa ataccaaatg tcccttgggg gagaaatcgc cccagttgg	240
gaaccgctag tctggagaaa ctccaagatt taaagggtgt agaagagaaa gagctgccag	300
agaagactga aagggcagtg gaggagagtg ggggtgtgtg gggggggtgt gggcaggagc	360
caaaagagtg tttcaaggac ttggtcatga tccttttaaa atgccagtca gatcatgtca	420
cttcctgctc aaaaccatcc acacgcttca catccattt gaaataaaat gccaaactgct	480
taccatgccc tatacacaga acaactgtaa taacctgggc acctttgaga gtgaaaggag	540
gcaatactaa taatcatgcc agggcagttc agggcacact ggaggtacca tctcctaagc	600
tcaggccct gccatctct ccagcttcat cccaaccac tttctgcctt gtccactcac	660
ccacgacagc cttcttgcca tttgtattgg gccattctca cattgcaggg gccagagctt	720
aggatgacaa acatatagca acacatataa tgtaatgtca gtgatattaa tagatgctgt	780
gaaataagat aaagtgaggt ggagacatag ggtgactggg ggattggtg ctattttact	840

taggggtcag gagatcgtct ctgaggatga atcacttatg cagagacccg aatggagaga 900  
 gggaatctaa gaagatctgg ggaagaggat tccaggcaga aggaacagca agtggaaagc 960  
 cctgaggtag gaacaagcat ggaatatcaa tagaatggtg 1000

<210> 73  
 <211> 1000  
 <212> DNA  
 <213> Homo sapiens

<400> 73  
 ttcctattgg atggtgctaa tctggtgcag ggtttcttaa cctcaggact actggcattt 60  
 tgggtcaggt cattctttat tgtgtagggc tgttctgtgg attgtagaat ggtaagcagc 120  
 ctccctggcc tctatccact ggatgccagt tatacccgtt ccagttgtga ccatcagaaa 180  
 tatctccaga taaaatacca aatgtccctt gggggagaaa tcgccccag ttgggaaccg 240  
 ctagtctgga gaaactccaa gatttaaagg ttgtagaaga gaaagagctg ccagagaaga 300  
 ctgaaagggc agtggaggag agtgggggtg gtgtgggggg gtgtgggcag gagccaaaag 360  
 agtgtttcaa ggacttggtc atgatccttt taaaatgcc a gtcagatcat gtcacttcct 420  
 gctcaaaacc atccacacgc ttcacatccc atttgaaata aaatgccaac tgcttaccat 480  
 gccctataca cagaacaact gtaataacct gggcaccttt gagagtgaag ggaggcaata 540  
 ctaataatca tgccagggca gttcagggca cactggaggc accatctcct aagctcaggc 600  
 ccctgcccat ctctccagct tcatcccca ccactttctg ccttgctccac tcaccacga 660  
 cagccttctt gccatttgta ttgggccatt ctcacattgc aggggccaga gcttaggatg 720  
 acaaacatat agcaacacat ataatgtaat gtcagtgaata ttaatagatg ctgtgaaata 780  
 agataaagtg aggtggagac atagggtgac tgggggattg gtggctatct tacttagggg 840  
 tcaggagatc gtctctgagg atgaatcact tatgcagaga cccgaatgga gagagggaat 900  
 ctaagaagat ctggggaaga ggattccagg cagaaggaac agcaagtgga aagccctgag 960  
 gtaggaacaa gcatggaata tcaatagaat ggtgatatgg 1000

<210> 74  
 <211> 1000  
 <212> DNA  
 <213> Homo sapiens

<400> 74  
 aagcttacct tggctgctta cactcttata caatgccatt taccttgtgt gatacataat 60  
 atcttgtatg aatcctatct tctctgtgtt tgtgtacctt tctttgaaga atatgacctg 120  
 tctcaataat tcttttaatg tttttctctt agtcctttta acatcagcag ggcatttgta 180  
 gtggtgacag gagaaacata aacatatacc tcttttctat tgcttttctg ctatttacia 240  
 taattctgta tgactctgaa acaaaagaac aattacctga caatttcttt ctgagtccta 300  
 tattctggct ttcatatcca atctcctttt atcatgctat tacctctctt ttcttctgtc 360

```

tttgaggatg ggaaaattca tcaacaccct aaataccagc cagagaggaa aaaagagtct 420
ggatggaggc aggactcctt tcaaagctga atctcaagca ctgatcacgg agcagcagca 480
aagagacact caaaaagagt ggagagagga aaaactagct gatctctaag gtgtcttcca 540
ttcaaattca ctataattat aagaatgtga ttactggagg aagaacaagg gcaggggcat 600
ttctgcaaca tgacgcaaaa aaatattgac cttaaatttg atacatatga actttctaaa 660
tgtagagaga agctacctcc ttgctgcaact tgtatgtgtg ccattcattt cattttaata 720
aaagtttgta aacatgaatg aatgcagggg acagaccacc tctttatgag aatgcagcat 780
agttcagaga aagtctaatt accaaaaact gaatacatgt ttatactgaa attttaattt 840
tttctatttt tatttttaat tgtgataaaa tataaataac ataaatttac catcttaatc 900
atttttaagt atacagttca atagtattaa gtccattcgc attattgtgc aaccaatttc 960
cagaactctt tttatcttgc aaaaatgaaa ctctataccc 1000

```

```

<210> 75
<211> 1000
<212> DNA
<213> Homo sapiens

```

```

<400> 75
accacaaagg ctgagggcat ggattattgg aaactctctt ctgaaaaatt ttttactaat 60
ttgggagatt aacagtcaga atcaatgggt gatggtttat agagtgatac caaccttgtc 120
cagtctctgt catcatttcc aatcaacaaa atgaataaag atgaagagag tatgcttatg 180
acatcagtga atagtacaga tctcagactg ctgaagaatg tacaagatga cttagcctgg 240
atccaaaaag ccaagctgga gaggtagggt ggttccaaca agacaaaatg taaaaacgaa 300
gaccaatact taagaccaa aagtcaagcc aaacaaaaca tgctgatgtg gctaaacagc 360
aagttgtgct aaaaaataag actcaagaag tcaaaggcca gttttatatg aatccaaaaa 420
gccaatgcaa ttttaatttg ctttaataaa tatgtattat ctggaaaaaa acacatacta 480
cagtgaagtt tctgtggaat gaaatactaa agcatgtttt cttggagaaa gagtttccat 540
gaccaaataa gttgggggat actccaagtt gatataaaca ggtttatttt ctacaggaat 600
actcaaagtc gatatggtga ctattgcttc tcaaagttat ttgaacatgg aacacttctt 660
ttttagtagc ctcttgaggc tgggtgtaaa gagaacactc ttgagaaaac actgaacaag 720
ggctgtctca ggaggcagtt ctctgtaagt gggactcttt ttaaaaacag aagagatcca 780
aacatcagat gagtgttggt ctaaagacc ataaggttcc ctcctaccct cgaagtctgt 840
aatacttggt tatccagacc taacaaacaa tcctaattcc ccatgacacc tggaccagag 900
tttctgatga gagaaactct agagaaatac tagtagcaga gtaatgattt aaaaaaaaaa 960
aaaacttttc ctccaatgag tgcatgcttc aaaagggtgt 1000

```

<210> 76  
 <211> 1000  
 <212> DNA  
 <213> Homo sapiens

<400> 76  
 ctccaggatg cgccccttcc cggcacagcc cactgccata tcttgctgga acctgggtca 60  
 tcgtccatcg tctatcacag gctccgccag ccttcgtgga tgccatctat gtccgtgggt 120  
 ctcacccgtc tcgccaccag cttccactac gacgctggac agtacacagg gagcagacgg 180  
 ggattccagg aggaagccac tgcaaatagg gcctgcagct gccctctctc cttctgaaat 240  
 cctagcatag tccaggacac agcacctccc tggctgagca gctgaactgc caagctcaac 300  
 tccctgattg agcagatatt ctgcagaaat agaaaaggat ggaggggaagg cttcttccca 360  
 cacaatgaac atcaaaccce cccaaggggc agtggtctggg gcctcccttc ccaaacagct 420  
 ggctcaaaac atgcacaaaa ttttccaaa gtgggctggg agcagggcag ctggcttcca 480  
 ctttcatatt actgatgcat ccagacatac ttccatagtg tttaaaaatt tttggatgta 540  
 tgtcaaatgc tcttaagagt gcgatcttag gcatgtggta aataaatatg atgtaatcct 600  
 cccgtctcca aggggtgctg tgccctctcc ctccctccct cactggtcct gggcaagccc 660  
 ttgacctcca cgatctctct gcgcctctcg tgacgcccac aacaaggggc tgtgccaag 720  
 ggaaaggtag aaagaaaaga ggatgtgctg tgtgctgtca tcatccctgt gccagagaca 780  
 gggcacaggg tgggtggcctt gcaccaccgg cgcattcccc acatggggaa gctgggggtca 840  
 ccctgcacca caggcatccc atcagcctct gtgacactga caatgattct cgtgaatgga 900  
 caggctgaat ggtcctcagc cctctcttcc tatgctggct gaactctgag gcgggaacag 960  
 gacagacagt ggctggaggc cctggcaggg agggcacccct 1000

<210> 77  
 <211> 1000  
 <212> DNA  
 <213> Homo sapiens

<400> 77  
 ctgtcagttt ggtgccctcg gctacgcagg gcctgttaga aggggtgccct ccctgccagg 60  
 gcctccagcc actgtctgtc ctgttcccgc ctccaggttc agccagcata gaaagagagg 120  
 gctgaggacc attcagcctg tccattcacg agaatcattg tcagtgtcac agaggctgat 180  
 gggatgcctg tgggtgcagg tgaccccagc ttccccatgt gggggatgcg ccgggtggtg 240  
 aaggccacca ccctgtgccc tgtctctggc acagggatga tgacagcaca cagcacatcc 300  
 tcttttcttt ctaccttcc ctttggcaca gccccttgtt gtgggcgtca cgagaggcgc 360  
 agagagatcg tggaggtcaa gggcttgccc aggaccagtg agggaggag ggagagggca 420  
 gcagcacccct tggagacggg aggattacat catatttatt taccacatgc ctaagatcgc 480  
 actcttaaga gcatttgaca tacatccaaa aatttttaaa cactatggaa gtatgtcttg 540

atgcatcagt aatatgaaag tggaagccag ctgccctgct cccagcccac ttgggaaaa 600  
ttttgtgcat gttttgagcc agctgtttgg gaagggaggc cccagccact gcccttggg 660  
tggtttgat gttcattgtg tggaagaag ccttcctcc atccttttct atttctgcag 720  
aatatctgct caatcagga gttgagcttg gcagttcagc tgctcagcca gggaggtgct 780  
gtgtcctgga ctatgctagg atttcagaag gagagagggc agctgcaggc cctatttgca 840  
gtggcttct cctggaatcc ccgtctgctc cctgtgtact gtccagcgct gtagtggaag 900  
ctggtggcga gacgggtgag acccacggac atagatggca tccacgaagg ctggcggagc 960  
ctgtgataga cgatggacga tgaccaggt tccagcaaga 1000

<210> 78  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 78  
tatattttct ggattttacat gccaggttac aaaaggagac ccacacgaaa tccctgaact 60  
cctgtgccca cccagagatt aacatggaga ggtcaggggc tgttttctct ccataggctt 120  
cagtggcctg gatgtctgag ttttcagaga caggataagt ccacatatta tttttaaaca 180  
aatctcttac aactcaaaag ctttcatatc ttactttctt ggtaagagtc aagtttatta 240  
tccacgtcca tacaacaca gctggctaca caaactgac taggacaaaa agtcagaaac 300  
atggggccat aggattctgg gtaaatgtgc tttctaaca aaactatcat atttacagaa 360  
aagcagacaa agtgatgaga gtcttctgcc tttagaatta gctgacttta aaaattaatt 420  
taactctgac atgtgacaag aattttatac atcattgcaa aattaaaaag gcactttgga 480  
gtggaagtac tgattacagc atatttttga tagagataat ggactttatt taaaacacat 540  
tctaccattt tctcctgtgt ttttcttga gtccacagag gaaagttact acacaaattc 600  
aggttatttt tattgacggt tatgttatgg tgaagctaga tgaatagagt ttaaagttaa 660  
gttttgttgg gtatttccag gccacttggc acatcaaca ggtaagcact ttttctcaa 720  
gaaaagtgtg ttgtattgat cttgctttgc tctagtattg acaattatat gaaatttta 780  
gcatctcctt agaattccca gctttttgag ggccaatttc tattcagggt tttatggcta 840  
atctcttatg acatctgtca ttccaagtat ttaaaactctc atatgtttct ttggtgtgca 900  
tttttccatt tgtttaagct cgtttcttag gtcagtgagg gtgtgtgttc tttcttttat 960  
atcacagggc tttgtccaca gggtagactc agctcatgtt 1000

<210> 79  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 79  
gaaagctgac aaaattacat ttcttgagtc cagtatctat tctttaattg tcttccttta 60



tatttgaact cttagtcaac tgtggtccaa agagcattca actgaggagg gaggctcgct 120  
aattttccct cacctagtga cgcccatgct tgagcttcat gaaatttaag ataattatta 180  
ttatatagtt atataatcat ttcatgtact atctttttct tcttctttac ttttattttt 240  
taaaagcaga aaacaataaa atggccatca attgcatgaa cactgctcta aaaagataac 300  
agtaagaccg aacctgaact gttggctacc tggccgtgcc atattaatag cttacaagga 360  
tcagatatag aaatatcaat cacagggtgt gtagagggtg ccatgtacag agcacaacat 420  
tgtatattaa aaggatgttg agcttttata attattgcta tggttttata cagtgttaata 480  
agcccatgat aaataggagc tcataattta tcttaatgaa gtgctatttt atattactta 540  
ttgatttatg tttttccccc aagaaagttt taaccttctg agacttagag actcatttaa 600  
atgctttgac ccccataccc tctttgcagg gtgcaggagg atgtgtatga tcttaacctt 660  
tacagcaaat ctcttctttt ggatggggta ttgcaatttt cttttagagg atcacactta 720  
gtccagttca atgtagttta gaaggggctg acttcatctc tggttccatg ggtggacgct 780  
tgatccactc tggttaagca aaatactgca tcagtgtaac tcatttgtga atgggtacat 840  
gatccaagct ggaccaataa gagccctacc tagagttttg cttgaattgt taggataaag 900  
ggaaattctt tcctgaagca ccaaggttat tttctggaga aatcatgacc aagagtgaag 960  
ccaatgcatg gaaaacaaaa gccgtgagta aaaaaaaaag 1000

<210> 80  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 80  
atgcatcatg tcttcatttt gtggcctcta atagattctt gggatgtaaa agaactcatt 60  
ttatatacat atgcaaatth aaaaccttct ataataagtc tgaatcacc tgtgtcctct 120  
ctgtgtttgt gttatcagca agtgaatttc tcagtactcc cacatcaca accccaatta 180  
ccactccata tgtttcccaa attagtagct aatagcgttt tcccaggcga atgtatctag 240  
aaataccag ggattcactg ctatacctaa gtcagcaatg gttcatcttt ctcttgctg 300  
tggaggagaa cttgaccaga ggagtccact tcccctggcc cggcagcttc ttgcatggga 360  
aactagctgc tcctgctgct acttggtgta tgatttacc tatagcacat tttatcttta 420  
cgtaaacaca caaagtcctt tcacgtcttt gttcctgttc ccatgccatg actccttctt 480  
ggaataccat tcttttattc ttactcacta aataagctct tctactcctt ttcttcgggc 540  
ccccttctc tgattcagct gagaaacaac tactgtctgt ctccatcaaa gctaattttc 600  
tgcctcctgt tttccacca tactttgcca tcttagacat ctgttgcata tcattttttc 660  
tgttacttaa ctaatgcatc agtcttcatt cattctctc ccagactata ctctcctgg 720  
gttcagagca tatctcattc atttctgtgt tacctttgct tatctcagt ctggcttcag 780

agtagatact tcagagatgc tatttaaatac agagttaggg tagttagaat aggagagaat 840  
gaggactcta tgggtgctcag gtgccatgca tcctgcaaag agaacatgaa aggacatttt 900  
tttttccttc aataattaca tggactcctt cagtgatccc tgtgtctgtt gggccttgag 960  
taattacctg caatctctgt ctttgtgagg ctattaatta 1000

<210> 81  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 81  
gccagtcaat gccaaagaca ttctgttcgg ttggaatga ataaaacttc tgatgcccac 60  
atggtaacct tatgctttga gaactcttct atagcacaat aaaatctgag ccgtcagagt 120  
aactaagtga tggaaaatga ataactaaat gtatagggaa agaatccaga aaagaaattt 180  
gtattttatt ttttctaagt aacttcacac gatatgtttg agaaaactgt atgatctagt 240  
gaatagaata ctcaaaaactc taatatacaa gtcacaggta tgggcccctag ttacttcact 300  
aaatgactgg ctttaggcag ataactgtgc tgggtccagt tactaactat gagaaataga 360  
aaatacatca ttacctttct ataatagtcc acaactatct cagcacaccc aatgtgacaa 420  
aaaaccgtct caagcccact tcagtaacaa ctgagaatct gtgggttcat ttaaatgtca 480  
aggccagcag taagtgaggg ctggttctga ggctgacata ttctgaggag aacatggtct 540  
tgctttctct tttctgggca cttttgtcct ctggatggaa tccattcttg ggcaggctga 600  
agtccttctc tcatggtggc aagatggata tgccaggcaa ccatcctgtc tgcagagagc 660  
ctgcctagtg agaagttttg ggattagttc tgacttgatg aatttggggc tcatgtttat 720  
ccctggatat atctcttttg ctgaggtgaa tggatatgtt gactgccaca cctgggtttc 780  
tgtgactact cctggattca gtgatggagt cagccccaag taaggcccat aaacaagggt 840  
ggaggagagt ggttcctgga aagaaagtca gggtaaaggc aaggggacaa atgccagatg 900  
ggcagtaaat ggcagctgtc caaattttat gcctgaacca ctgaaaggaa tcttcactct 960  
cactgtgggt attaacatag gacgcggtga tgcttaatgg 1000

<210> 82  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 82  
actagcttgg atgcacaagg attcaaggat gcatagttag caagtagcaa agtagttatc 60  
aagcctaggc gggcgctgac tccagaattc aagcccaagg tcacttctct atactatttt 120  
acattgtatt taagaactac atgaacatga atgcatggtg tgatgcttat agtttcctga 180  
tgcttatagt gtcctgatcc tacttctgca taagccatgc aaaggtagtg acccagactg 240

tagaaatgcg tcagagtgcg atataccaac aaaatgaaac gagtgaaagt agtataattt	300
tccaacatgt atacactctc tcacacacac atacacgtga gaggagaact aaagattagt	360
gacaggggat ttataacatt ataaaatctg agagcctgaa aacaaagatc caaggcagag	420
ctagagggaac acaggtatgg gtcagtcagg tgcaagttga gaacacagtgc ataggggttca	480
gaatgggttaa gtataaacag aactagtgtg acagaagtca ttcttacata atattttttt	540
agttggtacc aagatggagt agatgcagta tgtggtagta aaatcacagg taattaacta	600
aattgttaaa aattgaaata ttgtgctcat tactgatttg tctccaatat ttatctctga	660
tagtcaataa atcaaaatat atcaaagctt aaattgtcag aataaaaccc atgtttgtat	720
aattgcagaa aaattattga aaagcaaac ttgtcaggga atccacgtgt tatcattgca	780
cagctcatat gaatctgaaa agtcacaaat aaattagcaa catggagtta attggttttt	840
ctttttttgc ttactgtta tttttcttta ccacatgcaa tttcttttct ggtttttggt	900
ttattatgga aacaatacac tcttttttcc taatatttat gcttctgcat ccttgcttat	960
gagtttcttc ttacatgaat gctgtcgtcc ttcttctccc	1000

<210> 83  
 <211> 1000  
 <212> DNA  
 <213> Homo sapiens

<400> 83	
catggcccca aattagtttc ccaccttatg ttccactagt ttcatagaca aacctcttcc	60
tgccatactg gtctggtcag tgccctccag acactgcagt actgccttga actgggttgc	120
tgtcatcttt tctctctgtc atctaaattc tagcctgtct ttgatggcta aaagcctaac	180
atctctgtgg gcctcagaga aattatcttc ctctgcattc ctccagtgg catctctcac	240
taatggatta atcatattac cctctcctat tgttatgtgc ttttatgcat ataactttag	300
ccccccata ggaccaactg taatcccttt gaggacaggg gtttgatctt gtacctattt	360
atagttcccc acgtgcctag agcctcttgc acactgtagg ctgggggaaa atatttgctt	420
atgctgatga tctgagaaag ataatactgc aaacaggaga agtaaagatt tctttgtctt	480
gttcatttg gaatgaatta gtggcaggta atcagttaga ggtcagttca gaagggttaa	540
atacgtggac ttatcccctg ttacaggtct cttatcttta caaagattgt gttcctgtta	600
ctaacctctt tctaaatcat tgggtgtgtt atttacaaga aggactgggc caaatatgtg	660
aggaaacatc aatgtatact catccctacc atttgaaaa caagttttta gtgtgtgtac	720
cactgatgaa gtatgaagaa taacgttccc attcattcca gagtactcag gccctttgcc	780
tgggactgct agctacacat gcaaagtga ttctatatca gcattttgta aagcccacta	840
ttctcaccgt accagcttaa ctgcaaccag ttatttaata ggattctaata taatttaatt	900
ctccactggt agcaattttt gatgcacaa gtctgtgcct ttacacctt tgcacccctt	960

ccccagcact taactcagca ggttgcatat agcaggaacc 1000

<210> 84  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 84  
taacttgttc cagcacagat tcaaaagtct aaattctgaa gtctcaacta aatgtcatct 60  
aaaccagatg taggtgagac tcaaggatg tttattctga gagaaattgc tctccatctg 120  
tgattctgtg aatcaaatag gtaaagagct tccaaaatgc aatgggtgga cagacataga 180  
atcgacattc ccattccaaa agggagaagt aggaaggaat actacaacaa caacaaagta 240  
aacgataaat cttaaggctc cagaataatc tccttttgat gccccatctt ccaatcttcc 300  
aggcacactt gggcaggcgt tgggccccca aggctctggg tgtcccagtc ccagcccaca 360  
tgacagcact tacatattag agccacatgc caggctggaa atgccctcta gtggctctac 420  
tggtctatgg tcagagggta ggctgtctcc tatgactctg ccaagcacag ccttagtgga 480  
ggctttttgt ggtggcccca cccctatgtc aattctttgc ctgagcctca agactttcca 540  
gggcacccct tgaatctgt gtggagtcag ctttccctct atggtattgc actgtgtgtc 600  
ctggtggaga tgatacttag agaacattac caacgtttat catctgtgcc ctccagaaag 660  
gtggccactg gagcccacac cacacttgga cctcttgagg ccatgcctgg aatgactgag 720  
cagtgtctgt tcagaaagca gggagcagag atgaggtagc atagggcagg aagtgtctgag 780  
ctccagtggg catcctgggc ccctcttttg acctgtttct gtcccctagg ccttggcacg 840  
ctgggcctgt gatgggagca gcagccgtca tgatgtctga aatgctttta gtgggggtca 900  
ttcctccatt gccttgatga aaagcacctg gcttctgcag ttccatgtta atctgatcaa 960  
atggttgctg ggccacatcc ttggtattct ctcccaaaca 1000

<210> 85  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 85  
ccacagaaac attcttcagt agaactttaa tattactgtc ttataaaatt ctgtcaaatg 60  
aacaaaaagat aacccataat tacaccctaa tatgactgct ttaaacattt tactgtattt 120  
cagccttttt gctatgtata taattttaca gagttgtaat catacccagt atatgatttt 180  
atcatgtttt cccacttacc attataggta tttttaatat tgctacatag tcttcatggt 240  
tgtcattgtt aatagctatg ctgtaatagt tcaactgaatt gaagtgtttt atttacttag 300  
ctaccctatt atctttaaac aattttcta ttttttttat aataaacatg gacatatattc 360  
tgacaggggt gttctttttc acatcttgac ctacttttca catagtgtta caattacctg 420  
accaaagaat acaaaactttt tgtctcttga cgtatatttc caaaagattt ttaaaagggtg 480

cattaattta ctctgcagct ggtgtaaatg aagaccattt tgtcattggt ttcttgagag 540  
tagagcttcc aaaagtaggg atatgtggct aggaggaaga aatccagcct ggggcaggca 600  
ttctgtaaag aactccagtt ctccactggta cactggtttt atttttctct gtttcttgca 660  
gactgagcaa ttgataactc tgtgggtcct ctttggtttt accattgttg gaaactccgt 720  
tgtgcttttt tccacatgga ggagaaagaa gaagtcaaga atgaccttct ttgtgactca 780  
gctggccatc acaggtgaagt aactatgcaa gtgagaggca ggaagctata tgtgaagtcc 840  
ctatggcttc ctgcttttaa tgaattttat caaaaaaaaa aaaatgtaac gcacgggtca 900  
atttggaat aatttctgaa agaataataa acctatattt gaatatttcc tctggcatac 960  
ttaacacata tgaatgcctc taagatttca ttataaaagt 1000

<210> 86  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 86  
aataagcaaa tctattttga cagaaagatt catgattgct cctggcagca gggggtgagg 60  
aagttggtgg gaaatgggta cagagattct tttggcgatg atgaagacgt tgtaacagct 120  
tttgaatttt acaatccaga attctattct ctgctaatta gtcaataaaa gggcagaaaa 180  
tatacathtt aaaacacaaa gatgcagaca ttacattcca catacaagag gatgtacccc 240  
agcaaaacaa ggtgataaac caagaaagag aaagaatggg atccaggaac aacagettca 300  
accagggata acaacaaagg gaactactcc agtgtaaca gctgggcagc cagagagaca 360  
gcatgtagtc ctcatgaaag cagaaagaca gaggggttctg agacagaggt ctccaggaaa 420  
aaaaaaaaaga acctgactta ctggataaac aagtcttttag tttaaaaaac acaaaaaaac 480  
tgtatacaca tatatatata aaatcaggta gtataaagaa aaacagaact ccagagattc 540  
ctgggtcaca gaaggggaaa gggctgttca agaaagtga attgaactaa ctgaaaatac 600  
agctatcttt atattggaag gacagtcagg aagtcaacag ataaggccta aactgcataa 660  
agcaggaaac agcagactaa agacattatt aagaaatatg gaacacaacc aaaagaaata 720  
gcaaaaacaa tgaaaagtga ctgtttttca taagtggagc aggggaagag aaggggttat 780  
ttttttcccc atttatgtgc ttttaagaact acttgctaaa aatattgggc acatatgaat 840  
ttgataaaaag cgaaaaactt tttacttcac aagtcagct ttaacatacg ttgattacag 900  
tgaagttttt gttctgttaa ccactttagt aggatttgc taaatttagt gatttacaat 960  
gcctgcagta gaatcagaag atttactg aagggttat 1000

<210> 87  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 87  
 ccttctcttt cgggtatttt agtcagcctc tttttatcgc tgttatcaca gatatcccca 60  
 gagaccactt gttatcataa tttgctaata tttcacaaaa gatgaccatt tagtttttaa 120  
 ttaaacttta taggacttac actctcattt gttaggcaag gaaattgagc cagggtcaaat 180  
 taagtaaatt gcccaaaatt ctactgttt ttccaagtaa ttttaaagag tgacatccag 240  
 aaaatctgtg acttctagga atacatttag aaaaacatat accagagggt ttaattgcag 300  
 cattgttttt aacagcaaaa attggaacta aatacacatc aattggatac agataaataa 360  
 agtatgagat attcatggac cagaatcctg tgctgtaatt gaagtgaatg aactggcaat 420  
 gtgtgcacca gtatcccaa attataatat ttactaaaa aagcaaatg ctgaatgatt 480  
 catgctgtat gataacatta tataaagtct gagaacatga aaagcaactg caaacataga 540  
 ttatagctgc ataaataaat aataatagta taataaacat ttgtaggaat ggaatagaga 600  
 aaaacattat gagatccaga gtgccccaaa aaaacctgcc cccatatttt aaatcaacca 660  
 ttttctcatt taaccccat tttcctcctc acttactatg tgactagatg ttctttgggt 720  
 ttgttaaaaa aacatttccg attccttaac atacctaaaa atataataaa ttattctctc 780  
 attattttct tctacataat atacaaatta cttcaaaata cgtacacaac ttactttcac 840  
 ataataaat ctaacacagt ggcttttctt aggtatgcat tctactaaa tcatatatcc 900  
 ctttctctaa taataaaaag atttatatgac ttataattat atactaccat agctgggcta 960  
 tcatagtagc ctttctttt aatataaata ctttgatata 1000

<210> 88  
 <211> 1000  
 <212> DNA  
 <213> Homo sapiens

<400> 88  
 gggacatttc atgctgggga acatttttagg caaatgtgcc ccaagacctt ttcgataagg 60  
 atactccagc gaaacaaatg agactgttac aggaggcagc actgaggcag ggcagggtggc 120  
 attggagaac atgcacacca ccccatgggc accgtgcaac accacccacc acccatggaa 180  
 gtggtgacaa cagtggggag gggaagcctg tcaagcagat gtcaccaggt gcttcaagca 240  
 gtgtgttagg tcctgtctta taggtgccag gccaaactcac ccaccttctc tcgactcttg 300  
 gaaagaaaat agtggaggtc tttctaaatc atgtgagaca ataactccc cagagggtgcc 360  
 atcctctaga ttocagggga taaagacgag cacaagaagt actgctgagc actttgtgtg 420  
 ggatgtgtgt ctaaacacga caatctgaag acagaggtgt agaaattggc aagtttccta 480  
 aagcatgaca acacacaccc aaaactcttc cataatgatt ccctttttcc ctgtattttt 540  
 cctggatgca ccatcactat gggaaccagg atggttactc ccaattccct gtcacccacc 600  
 gcttatttaa taaacgattt ctactttact gaaattgatg cttcgttttc ttctaattcc 660

attctataact ttacctctgc tctgagttac actgaattta taacccttct tttaaacaga 720  
agtcttgcaa gaacaaacta cagcagtatc agcaaccaac aatgccacca atacagatta 780  
aaaaaacatt cttatctgag gccaggtaac caaatttatg caaaataact caacagatgc 840  
tggtcagtac tagctgaccc atgaatttaa gctcttactt ggaagaaata caacccaaag 900  
aggagagaaa ggaaaaaaat gagtctcata ttaacatata ataaaacctt attaatgat 960  
aactccataa attatgagtg gcaatcagat agataattca 1000

<210> 89  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 89  
tattatgtta ttgtgtaatt atttgaatta ttgtcccctt tccatcaatc ccccaaacac 60  
acacatatta ggtggaaatc cttaggggct gagatgatgt tttatttaca tctgcatgcc 120  
tgatgttaag cccagtgtcg ggcacgaatg cgatttagtg agtgtttctt gaacatgaat 180  
aatgaattca ccagtgaag catgagtgga tctgggtggg gcacaaaagg ctgactccag 240  
gttccaggaa tctgggtgga gaaacttctg ggtctggagg agcagaggac cactgtgtta 300  
ggtctacgtg gttctggctg gcagggttag caaggatgca gaggagtctt tgggtcttgc 360  
tcaaagtata atttaaaaca acaataataa ttaacattca tttagttctt actatgtgtc 420  
agtcccttat tgccctctat gtattcagcc actaatctc aaaattctag gggtagata 480  
tttttccggt ctatactata catatgagaa aaagggtaga acagggagggt gcagaaactt 540  
gccccaggat acacagcaag taaaatggga actgggattg gtcacctagg gattcttggt 600  
tttttagatt tgttttttta atctctctat agccccttag gttatttatt gatattttta 660  
ctttttattt tgaaataatt gtagattcac aggaagtac aagagagagg tcctgtgtac 720  
tcttcaccca gatttctcca atgcttagat tttatataac tgtaatacaa tatgaaaacc 780  
aggaaactga tattggttca atatatgtgt atacttctat gccatttcat catgtgtaga 840  
tgtaaccacc atcatgacca agctgcagaa ctgttccatc accacgaaga tctgccacct 900  
gttgctcctt taaagtcata ccagccctct tcctgtccc caccactgt cactatgctt 960  
aacccttggg aaccactaat ctgttttccc atctctatag 1000

<210> 90  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 90  
atgcatacac cagagccgac ccgcagactc tgcaaccag gccagctgc acggtcagtt 60  
tggaagtcta cacaagcatc tagaggacct ggacacaaac agggctaatt cagggtgcca 120  
attcatgtcc caactctgtc ctgtcaggcg actaaggcag ggctctggga atccagggac 180

```

agggtggagta actcgtacac agtcagtgtg ggagtccttag caggtgactg ggtcctgccc 240
ggactcgtgt gggatggagg gctgggtaaa ctcatgtctg caataaaagg gacagaatct 300
cagtgcaaaa gagactagaa aaaatgttag gttccagag agaggctgga attcagaggg 360
gaagatggaa gcccatTTga tatagtagtg gtgaagatgg aaggtggccc ctgccgtgag 420
gaagacacct gagctatgaa gagtggagta taagcttgga accagatgtg cacataccca 480
gagttcatgt ccaacatatt tcaaaatctt tgcaaagtct gtgtggatcc ttaaaaactg 540
gggagggcag agccagcagt gggcaggtgg cccccacctg gaggaatggg attatagagt 600
ccaggagtga ggcagcgccc tacagtttgt cctcatcctt ccattttcca cacttccagt 660
ttcctttcaa ccacttcaga aaaaaaaaaa agtccagaaa gtctaattgtt gccaagtTTa 720
gaaaccaggt cgtcattagt gtgagtggaa tcaacgttga ttacagtctg gtccttttca 780
agtttctttg atatcttcaa aagcccaatc atcctgttcc atctaggaca ttaagaaaaa 840
tacacccaaa gaatagtctt tcaagtacat tgccaccgta gctagatgat tattatcctg 900
actattaatt actattatga ttactgttgc catgggtttt atgtttttct gtgtgcccac 960
ccaatcccac atccagccac cacagccact gctgggtttt 1000

```

```

<210> 91
<211> 1000
<212> DNA
<213> Homo sapiens

```

```

<400> 91
tattatgtta ttgtgtaatt atttgaatta ttgtcccctt tccatcaatc ccccaaacac 60
acacatatta ggtggaaaac cttaggggct gagatgatgt tttatttaca tctgcatgcc 120
tgatgttaag ccagtgctg ggcacgaatg cgatttagtg agtgtttctt gaacatgaat 180
aatgaattca ccagtgaag catgagtgga tctgggtggg gcacaaaagg ctgactccag 240
gttccaggaa tctgggtgga gaaacttctg ggctggaggg agcagaggac cactgtgtta 300
ggtctacgtg gttctggctg gcagggtag caaggatgca gaggagtTtc tgggtcttgc 360
tcaaatagata atttaaaaca acaataataa ttaacattca tttagttctt actatgtgtc 420
agtcccttat tgcccttctat gtattcagcc actaatcctc aaaattctag gggtagata 480
tttttccggt ctatactata catatgagaa aaagggtaga acagggaggt gcagaaactt 540
gccccaggat acacagcaag taaaatggga actgggattg gtcacctagg gattcttggt 600
tttttagattt tgttttttta atctctctat agccccttag gttatttatt gatattttta 660
ctttttattt tgaaataatt gtagattcac aggaagttac aagagagagg tcctgtgtac 720
tcttcaccca gatttctcca atgcttagat tttatataac tgtaatacaa tatgaaaacc 780
aggaaactga tattggttca atatatgtgt atacttctat gccatttcat catgtgtaga 840
tgtaaccacc atcatgacca agctgcagaa ctgttccatc accacgaaga tctgccacct 900

```



gttgctcctt taaagtcata ccagccctct tccctgtccc caccactgt cactatgctt 960  
aacccttggt aaccactaat ctgttttccc atctctatag 1000

<210> 92  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 92  
tagtttctct ggtctgcctt ggggaagaaa ggagagcagg agaaagaaag gtgggagaag 60  
gccagaaaga ctttgtttct gaagctcttt cagtttcctt cagttcaaag cactcatcac 120  
accaagacac catactgtgg ggtatcacat tctgagccct aacacttcca atattatgct 180  
atgaatttac atcatgattt caggtaatta ttccaacaat gccacaaggt gagcatttgt 240  
gttatccagt ttcacagatg cagaaactga agtggaaaaa attgactagc attatatggc 300  
tggcaagtga tcaaacagga ttttctcatt atttcattca ctcaatagtt attgagctca 360  
taatatatgc caggcattat gtcagacttc atggatacag acagggtacac agtaaacaag 420  
gtggccactg cccaaatgga gcttgcattc tgggtgggaa gacagataat aaacaacaag 480  
aaagaagcaa tataacagat tgggacagt ctattaatat aagtaaatga aggagggata 540  
tcatcaggag aatctgggaa ggagtggatg ctacctgaga caggatggtc aaggatctgc 600  
ctagttgcaa agcactagac tttccacaac cccttctacc ctccagtggg cctctgcagt 660  
atatatggca accaattctg gtttcatgta ttctaccact tactccaact ctagtaaata 720  
tctgcaaagc ttaccattgc ctacgactct cagattattt cccaagatg ctgcagaatc 780  
cttataatgt ttctcagcct caatagaatg aaaagcaggt ctgtgcttat atcacttaat 840  
gaccaaagag gaaggaaatt tacaattaaa gtgtactttg ccaactgtgg atgaattagt 900  
taggtcactg tgatctacag gttagatgtc tgttcagcag tgcctctac ttgagattcc 960  
aaggaggttg aagctcacta ctgcgccccc ctgcgacccc 1000

<210> 93  
<211> 1000  
<212> DNA  
<213> Homo sapiens

<400> 93  
atcagcgaca ccatacctggt tgtcatccct gagctgtgtt aagtaggcac ttcccctaag 60  
agagttaaag gggcactcgt gagatactaa gaagactcct tccccagcc ccaggcctcc 120  
ttgtaccttt tgccctctca ttctgtctgc tgccttctgg gaaatgatgg gactggcagg 180  
ctgtactatg cagcagggat agcagggtg tttgtctgc ctcaggaag gcagataacc 240  
cctagaaaca ggaagagcca aatgaggttg tgtaagtctg aggcagaaac attagtcgtg 300  
agagcaagac ttgcatttgc aagagccagg ctgtgtgtgt gtttgtgtgt gtgcgtgtgt 360

```

gtgtgtgcat gtgtgtgcac gtgtgtgcat gtgcgtgtgt gtgtccgtgt gtgtgtgtgt 420
aaaactggat ggccaagagc caaccctgg agggcacgga gacagggag aaaacagagt 480
gaaacaaaaa tttttgtgta gaaggcataa aagttatcat cacagactcc actgtgtaaa 540
ggcataactt gctttattta tctctagtgt atatgaactt agcctccctt tccattcagc 600
ctgtgaaagg agatagtgtc tggggcattt ggtagaagaa ggggatggga gatgatcaaa 660
acccaagta aggttcatat ccaatatagt gtctaagcag caaatgacta atggccgaag 720
aaggagacta gacagaggat tagaggcagc catggggctg gtgcagctgt ggagagctct 780
gagcaaagaa acaagggttg caggtgagga ggcctaggat agaggccaga aggccaaacc 840
tggggctgtg cagccagtgg tcatgggtgg acagcaggca ctggctgggc attggctggg 900
catgcagatg cccaaggcca gctgtgccac atagaagccc tgaggaagtg agggtaatta 960
acccctgaac aaccagatc atcttcaggg gaacagccag 1000

```

```

<210> 94
<211> 388
<212> DNA
<213> Homo sapiens

```

```

<400> 94
ctgtgttggg ttcctccagt ggtttgtgga tggagatcat ggggtgtgtt agtccatttg 60
cattgctata aaggactatc tgagcctagg tagtttgtaa tgaaaagagg tttatttggc 120
tcagggttca gcaggctgta caggaagcat ggcctggca tctgcttggc ttccgtgag 180
gccccaggaa gcttccaatc atggcagaag gtaaacggga accagcatgt tacatggcaa 240
gagggaaagc aagagatggg ggaaggtagc aggcctttt aaacaatcac atctcacatg 300
aactcttttc tttctttctt tttttttttt tttttttgaa atggagtctt gctctgtcac 360
ccaggctaga gtgcagcggc acagtctt 388

```

```

<210> 95
<211> 662
<212> DNA
<213> Homo sapiens

```

```

<400> 95
atgttaaaat aaatattcat atatgcaacg gcaggacttt aaaaaatca cacaatagg 60
aaaacaaaag agaccattcc agaagtcaac aagaaaaata agtttagttt tacaagaagt 120
tcacgatctc gtccttattt taccacgtgc tagaatttgg tgaccaaagt accagaacat 180
tagttttag aatagtaatt tttaactaa atttttagca cagacatta aaaaaaatt 240
atctggcagc tgaatacaaa acgcaacaac aaaaaccaa acacaaatgg agctactcta 300
gttagagtca gagaggcaga tctctgaacc atgcctgcct gcacacaact caaaaaacta 360
gtaatgtaga gtgatttctc aagcctcttc tggtagtcta aacattacag attcttctga 420
ctaaaaagag aggcaatccc tgagactctc catagaaacc ccaggctctg tagaagccat 480

```

gaacatttgg tattgagggg ggaggcaaca gagtctccag ctgtagtttt gttttgaacg 540  
aatctggaaa ataaactgaa aaacaattta aaacaaaaag actttttaat agtaaagtta 600  
aagttgatgt gagatgttgg aataaaaatg aaggccattt caaaaccac cacaggcaga 660  
tg 662

<210> 96  
<211> 644  
<212> DNA  
<213> Homo sapiens

<400> 96  
cctgcaaagt ctcttcctgc tgcaccttc ttctgaaacc attaatacacc acgacccact 60  
gaatgaagcc caatctcaaa tcacagtga aaatcctgca acgtgcaggg tgatgagtgt 120  
ttacattagc tgaaatgaaa tgatgtaata ccagaaatcg agggagggtc gcgatccaga 180  
gtcagggcac tgcaaaaacc tctgtgaaac ataacttttc tacattacaa aaaaatgtcc 240  
ttgcgtttta gtaatctggc ttctgtaaat ttaggattac ttggattttt ctgatctcat 300  
caatttgttt tccaaataga aattcagaac ttcccaatta ctactgttt tagtcaagtt 360  
taaaaaaag ggtagcaaat agaaccctaa gtgtatacat gtgcaaaga cccagtatca 420  
aggaataat aatagaagcc agccatccag gtatgtgggc acctgccatg ctgcagaata 480  
gcagagcctc ccaagggtct aagtgccttc aaagtaaaga caactcctaa gaaagacagt 540  
atttgtttaa gccagtggcc aatttttctt cctataactg atgatgaaca agaaaacca 600  
ggagttccta gccctattat tgatgggcaa ctgctattga ttac 644

<210> 97  
<211> 582  
<212> DNA  
<213> Homo sapiens

<400> 97  
acaaggtcgg tgtacacccc ctgtgattct gggagtaata tcttctcctc ccttggatat 60  
taggaacaat atcacggcgg ggggtggggg tgtgtacagc ctctgcaata ttgggagtaa 120  
tatcatcctt tctccccact ggatattagg aacaatatca caggagggtc ggacaccccc 180  
tgcgatattg ggagtaacat cattttcttt tcccagtga tattaggaac aatattgcat 240  
tggggtgtac accccttccg acattaggag taatatcatc ctctcccaca gtggatatta 300  
ggaacaatat ctgagaagga gtgtagaacc cctgcggtat taggagtaat atcatcctct 360  
ccctccctgg atattaggaa caataacaca gggagagtat acagcccctg tgatattgag 420  
agtaataata tctctcccc atctgaatat taggaacaat atcagggggg tggggtacac 480  
catttgcgat agtgggagga atatcatcct ctccccacct ggatattagg aacaatatca 540  
caagtggagt atacaccccc tgcgatattg ggagtaatat ct 582

<210> 98  
<211> 502  
<212> DNA  
<213> Homo sapiens

<400> 98  
tatttaataca tataataacta aatataactgt attcagaagt tttttgtgtt ttagtcaggt 60  
aagatgcagg gtgtagaggt gttaaccttt ccttaaaatt ttaatggcta gatattcttga 120  
gatctgtctg atgtaggagt ggaaagtggg tggttctttt cttcccatc ataaaggctc 180  
acagctgata cccctataaa gaaagactgg ttaacaagag aaaagcacia caaatattatg 240  
aatgtgaata agtatgagag ccatacaaaa atatgaaaat tcaaagaaat ggtagacga 300  
ttgatgctta actaccttct tcattaggga gaggaaagt gggcgaggag tgggggagtg 360  
gggaatgggg cccctccat ctccaggagt ggataatgt ttgtaaataa ttctgttttg 420  
acactgaatg gagcggaatg gaaaggacaa acaataggaa tgtgaggggt ggaactgcat 480  
ggtgaacaaa ggtgtctta tt 502

<210> 99  
<211> 541  
<212> DNA  
<213> Homo sapiens

<400> 99  
ccgaagccga aaagtctgaa actggcccaa agtgggaatt tatatccctg ttctcctgct 60  
ggaatgttgc cttttcctaa accacccatg gtcccgccct acaccatcct gtacctatac 120  
aaaccccata ctgagccagt agacaggact atggttgagc attggagaga agcagcttga 180  
tggtttaaca ccgaagaaaa atccagccag agacggccag aacttccggg gaggggttacg 240  
ctaccgaccc tgtctccttc tcagctcccc ttctgcccga gagccacgtt tcattcacia 300  
taaaatcccc cacatccacc acccttcaat ttattcgtgc aacctatctt ttctggtgctg 360  
gtggacaaga gcgcgggagc cacagggtga gatacaaaaa gctgtcacat tggccctttg 420  
cccttgctgg cggagggcag ccgcctcaca cagaggcaga gggccactg aactgttaac 480  
acttaagcca tctgcagatg gcagagcaaa aacagcactg gaacatgcc tctggggctt 540  
c 541

<210> 100  
<211> 610  
<212> DNA  
<213> Homo sapiens

<400> 100  
atagagacga agttaaacac ttaatttgca aaactactga gaagtaaatt tcttggtcca 60  
aggtaactgg agtaattgcc aaatgcagat aaatcctccc cctgagtagg aagccccaca 120  
ctgttttgaa aacaattcct agactttgcc cctgttgaag ctgattgaat gctcaaccac 180

aagactccac tgttggttagc tctcgcttac tgcttttagg ggcggagtta acacttttca 240  
aaaatccgag cttccctaataa aaatacaggg atttagtgaa gatttttgatt gtctgggggtt 300  
ggcattcctg aggacagaat aatttatttt gctctaagca ggtgtgttat gagaacagag 360  
gctatgttga taagagatcc ctgggagctg gtaatatatt atcttctgta atttcttcca 420  
aaaatagact taatggaaag aggatgcata atataccccc tctcaaagga agcgttcccc 480  
aatacaacag aagcagtcata tctaaaaaca gctttatggc tctgcagtca ataactctat 540  
tttctcccct ttcacaactt ccttccttct gctatgtaag aacttatgtg agggcacaca 600  
cacattcacg 610

<210> 101  
<211> 524  
<212> DNA  
<213> Homo sapiens

<400> 101  
aaaaaaaa acccccatga tatggatatt gttatcattc cttttctcac aaatggtaat 60  
attgaaatta atagagggtg tatatcgtgt ccacagtcac acagttagaa agcgtcagag 120  
ccagggtttg aactcaagta gcctaactat agaaccata tttttaatca ctatacagta 180  
ttttactatc tgttccatca aaagaaatca tttttcagag tggagatgat agaacataca 240  
tgagaacaag agtattttaa tccaagatac ctgcaaagca tctagacact ctagatttag 300  
acttttagct ccttgcccaa gattaattac ctttcaggaa aataaaacta cataccaatg 360  
agatcactag acctctcgca atgatctatg aagaataatg ggaacagcta tctgggtatc 420  
taatgggcta gagtcagata aatgggttct caatagattt ccagaataat ggggaaattt 480  
ggttttgcata taacaatagg ctacgtatgt tatattcatt ctag 524

<210> 102  
<211> 677  
<212> DNA  
<213> Homo sapiens

<400> 102  
tcctttctct ctttcaatcg tgtggagaaa ataattatca gttgggaacc atcatttttc 60  
tactaccatg aatgcaaag tacttccatg acccatcttc ctttacgaat aaagttacaa 120  
tataagaaat accactacac atatctgagt ttatctttta actgtctttt agagccatt 180  
ctcttctgcc ttcctagaac ctctactatg gattatccct ttaccatagc attgtcattc 240  
tcttcctttt aatgcatttg tttcccactg atttttaaac atgattgagt cattttcatt 300  
agagactaaa taaacatcct cattacatgg ttcactagga ccaactccctc ttcagttgtg 360  
tgagaaacta agctttttaga aagagacgtc caaactcagt atctctattt ctgcatgcca 420  
cacaaatcca gtttgatttt catcctcatc agtctactaa aagatgtcac taaggacacc 480  
aatgaattcc aaaaaagccc ctgaaatcca atggaaattt gacatttttg accactttct 540

ctttcttcaa acattcttcc cttagttttc caagatagtt ttcttctttc cttcttactc 600  
actctatttt gatcttcttt gaaaattcat ccacctctac ccagtcataa aatgttaaga 660  
gttgaggggg gcagtcc 677

<210> 103  
<211> 428  
<212> DNA  
<213> Homo sapiens

<400> 103  
caggctaaat atcataaata aaacatctct catttctgtg aataggaaaag cacacttgag 60  
tgaagcacag acatgacagt tgagcatgta agagatccat tgggtgctac ttgagaaaagc 120  
agttggactg cattctgggt ctctctgaag tttgctttta ggcaagtacc agatggattg 180  
tatttttagaa aagatttgtc tggaacattt cctgatgtca ttatccagag acaatgagac 240  
aactcatttg cttatgaggt ttttactaca gcaatctaga gatggaattt ccaatggaaa 300  
taaaaaaggg tttttataat ttctatatgt acactggcag ctccgccttt taaaaaatta 360  
gttcctttta atgaatgtat tttgggagta gattatagtg tatttagtaa attggcactg 420  
tgtttaga 428

<210> 104  
<211> 657  
<212> DNA  
<213> Homo sapiens

<400> 104  
tctcattttg aaaatgtaag tggatatcac tactgcattg cctggaaatc ccacgaggaa 60  
gataatgccaa taaataacag ggaggtagtg catcttgagt gggatgtttt catcagtgca 120  
atttcacaaa gcagctgcat aatcggggaa atcagaagca tttgctaaat agtctagtgg 180  
ctcattcatg gttgtctcct ttcatcttgc aagaaaacaa gagagttcag tttggcaata 240  
tgaatcaaat gagcagtaac tcgctgataa aggaaaacag aaaacattaa tgatagggta 300  
ataaaaacaa ggatctactt ttaaatgaaa attattctaa catcctaaat ttgccacttc 360  
tctctcttta atctcaaaag agaccctgtg gagaagaaat tgaatttcca agaaaatgac 420  
tatgaggcaa gttactaaat gcatctaata aaaatataaa agttaaatta ccatgagagt 480  
taaaatgagg gattgggaga aaaaagccac atgtcgcttt ggaaaacaat ttggcaaggt 540  
caccatttgg agaagccata gggatatgcc attagagact taacaacagg acctactatt 600  
aaccaagtgt gatgcatgcc accatcactt acttctacat gtcacaaaat actgaaa 657

<210> 105  
<211> 533  
<212> DNA  
<213> Homo sapiens

&lt;400&gt; 105

ccacctgctg tcctgctaga cgtggaaaga ttcgcagcaa cagagcaggg aaaatgagtc	60
aaatggaggc caaaaatgag aactaagaga tttgtgagaa tattcaagca aggcaaggag	120
aaaaataagag aaggaaagta aaatatagcc acaagcaaaa gtggtaacaa aatgcttgat	180
atgaagtcct atttaccagt gataagccac atggatagtt agttatgagc ttttttgtaa	240
tcaacaggaa aaggaaaatc acaattttca agattcccag tgtctctaag gtataaagcc	300
caagtaattg gagagaagca caactatttg tggaactaag ataaaaatga attgcctcta	360
gtcagttttt gaagagccac ttgtccaggg tctcacagct gctcggccag aatttgaacc	420
ccaaccacat agttccagag cccacattct cagacatagc cccaatact gcctctgggc	480
tggaagctggt attctcaata actgtttgtt gagtgatag gtgaatcacc att	533

&lt;210&gt; 106

&lt;211&gt; 595

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 106

tatccacata aatgtgcatt ttcttttggg ccaaaatgag gcagaggtgt catgtgaatt	60
tttcattcct tcacacaacg atagtctctc acaaaacaaa gaacaaaagg aaacatatgt	120
tcacagtggg aaggattatt actcgatcat ctgtataagc atggcccaag gagcctttgc	180
caacctactg gggatgtcac atgtaaaaag gtttctccaa aaggttggca atatgattta	240
ttaaaggagt cagatgacat gggagttaag ggcagcaaac ttcatttga tggaaggat	300
ctaagctgct ccagcaaaat gaaaggatta tggttcacct gccaaactg tgcaatttat	360
ggatgaaacc tcaaccacga aaagtgaac ttctttgtgt gtgtgtatgg ggttgcgagg	420
ggagacatag gaaaggaaag gcagacagac cgtggaaaac agatatttcc cctggataag	480
agtggaatgg ccagtctcat aacactcatg tattatagaa ttaaataaa acctgtttca	540
gaaagtacaa tattaagacc ctttttaaat cttagatattc tttgatgata tctct	595

&lt;210&gt; 107

&lt;211&gt; 596

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 107

tttcctactg atcagagtta ctgtagaatt tgatttaggt gtgtaaatta gtctgaggca	60
cacattcagt cttaggcaac cctctctgtg atggcatgcc tcaaagcagt ggtttgaatt	120
aggggcaacc ttcaacctg agggacactt ggcaacatct tgaaatattt caatggtctt	180
aagtgagaaa gtgctatttg catctggtag attcaagcca gggatgatgc caaagatttg	240
aaaaaacaca gaacaggcca tacaacagag aattatctgg tccaaaatgt caatggtgcc	300
atggttgaca aaacctgaga taagcttagg gaaggatcca gcacagagca gaatgtattc	360

tctctgtaaa gaagccaatc ccaaagagaa agaagttgag taatgctgcg tatatttact	420
cactttctct ttccaaatct cttagtttga taattcactc gacttgccct ggtaaggaat	480
gagggaggaa gcaaaaaaga ccaagcttgt gttacactaa ttactgtccc tcaacagaaa	540
aacgtgaggt gaggggtaag aaagtccccc cattctcaca tctatatcca atacat	596

<210> 108  
 <211> 603  
 <212> DNA  
 <213> Homo sapiens

<400> 108	
tttgcctctt ttctatggt catcatctca ttgaatggca ccccatctg catggtagcc	60
tgggaaatat attaaggtat tatccttgaa ccttctttct ttatcatccc tatgtccagg	120
taatctgaaa ttctgtcaga atatgcatct ttaatctatc ttaaactggc ccatttttaa	180
aaatttctat ctatcttgac cttactttac ctaaatgatt atcactctcc taattgtttc	240
ctaattgggc tcataggcaa gacaaatctg ttctttatc tgcctctaga attatctttt	300
caaacacgga tgtggcctc cttctttctt acaaatgacc tcatagtccc aaagacaaag	360
tctatactct ccctaaataa cattcaaggc cctcactcac gcagctccct gattcccacg	420
tcagtatttt tgctctctc cccttcccaa agcacactct cacatacgcg ttattctacc	480
tggagtcata ttaagctact ttcaattctg ggctttctct tagccttcaa ccctctctta	540
ggctggtgca ttcttgggga gtgggtccaat ccatgcacgt gctaccatgc acccaccttt	600
ctt	603

<210> 109  
 <211> 575  
 <212> DNA  
 <213> Homo sapiens

<400> 109	
ctgcatgttg tctattgggc tgatccatgg gttgcttttg ctccaaggtc caggctaaag	60
gagatgccct ctcttgggga atgtcatgcc cctgctagag gtagtctctg cttggactgg	120
gcacactgct acttcgctgc tcatttcac aaccccagcc agccactgtg gggcaagcca	180
gtgttccttg cttgtcagag atgtgtact ttgcatacaa tggatgaag agtgaacagc	240
aggggtgaat taaacagtca accacaacct gaagccactt tccctgctaa gtggacctca	300
actcaatggt ctcatctga aagatgtggc ctaaatctt gcttggatg gtaattctc	360
tctaatagac tctgtgttc tcttgccagt caagaggact gaaggggatt gaaggctga	420
acctaggctc agtggctact gccctctctc cacagccgct ggcttcagc agacattcct	480
gatgtgatg tgctccttg agtgtgagc tttgggggaa atcctgttgc atgggtgccag	540
accctccttc cccatctcat aactccatca cagag	575



<210> 110  
<211> 402  
<212> DNA  
<213> Homo sapiens

<400> 110  
ttgtggagca gtttagagaca catggcagtg tccttgagtg gctctgagtg tgggaccatt 60  
ttctaggtga tcactcagca tagcttaccg atcagactca agtgaatgga acctgccctc 120  
ttccctttcc tcttggtttt ggaacagttg ctaccaggtg agtggttttt ccctccagac 180  
agttactgag agtaatccct gagcactcac tgggtgcctg ttctgtgctg acagtcacat 240  
cattcatcct aacagcaatt ccattctgca tcttctctgg acacccccag gaccatccag 300  
gacaaccctg cctgacacca ggcctagtgt ggctccatga taacaaagac gcaggtccag 360  
agacaatccc cctacatggt gcctgcatct gattccccct gg 402

<210> 111  
<211> 564  
<212> DNA  
<213> Homo sapiens

<400> 111  
tcttgcactc tgggccccca aacaagaggc cactcagaaa tcacagtttg agaacaaggc 60  
accattgccc cctgagcctg ggctttcctg aggcttgggt aagagaaaga gagatgagaa 120  
ggctccctgg gctacagagg tctggagaga agctggcacc tgggaagaac aatttccccca 180  
gcagctagcc aagctgggggt cttccaagtg gatgcagaga cctgcctcgc tgccctcccc 240  
atcctctgag agtgcccttct ctgggctttt gcttcaaaga gccatctttt tcccatggtc 300  
actcatcttc cttgtccttt gcttcatgac acctgagcg tgttagaagc taatcctgaa 360  
caagcataga aggggcactt ggggtaggag ctgcagtggc accacccgag aggccagctt 420  
tacctcccc aaagatccac tgcccagaag ggaagaccag gggcctccct ggtgccaagg 480  
gcttgagagt atgcatccaa tgcagctagg tctccacac actgtggtgg ggccccctcac 540  
cctcagatca gcatcttact ctca 564

<210> 112  
<211> 433  
<212> DNA  
<213> Homo sapiens

<400> 112  
taacaaaaca ctttttatca tatatgaaac tctgtacaa tgatttggct agaagaaaaa 60  
aatagttgga aggtcaaatt tgttttaaaa catctgttca aaagcctgca ttaaactttt 120  
atctgtcctg acaaaacatg tctcaatttc tttctaaagc agctctattg tcctagcata 180  
tgcctcacca agttctttta agggcatttc caaccttagt tctgacaatg aagacacaaa 240  
gtagggttagg ttccaaaacc acccttecta gccctccctg tagaaaatac catgttgac 300  
agttacatgt gtcccctgac acaaacgaca ctcatcttac gtaggtcact ggacctcaaa 360

ctgttggtgc ttgctgtccc agccaattca agagtgaagg aagatgtaac cagacataca 420  
tatctccctt tct 433

<210> 113  
<211> 461  
<212> DNA  
<213> Homo sapiens

<400> 113  
cagtccaatg ctccagtttt atagattggg aaaactgaga gcctaagggg tcacttgta 60  
tagctcctat ccccaaactt aaaaaacaaa gagttttaca gaatgagtc aatataattt 120  
gtttgggcta ctatttcatt ttaccatttt atccctatta gtatttatca ccatacattc 180  
aaaggaattc atacatgtag acacatctga ggtgttcctg atttctcctg ttcgacctgt 240  
ggtaaaactc ctgtggcact atagcacctt tagcttatca gtcttctttc cctcacctca 300  
tagatcagaa cttatcagcc cccatcctgg tccttctgaa tcttttgta agtcattgct 360  
ttccaatctc tgataaagtg ttgaaagggg accattatgc ctctcagaga tacacacagt 420  
catgtgccac ctaactatgt ttcagtcagt gagggaccat a 461

<210> 114  
<211> 444  
<212> DNA  
<213> Homo sapiens

<400> 114  
ccaataccac catctgaggg tctagagaag gcttgattta ctttcatgag tcccggaata 60  
agatctcctc aaacaaggaa ttttttttta atcatggaag tatggcaatg ggcaactaaa 120  
ccaaaagtct cagtgtcctc ctcatagata gcttcgctca gaaacaggca gcctgggtag 180  
agagatggaa tgtaaaagtct tattaatgc tcagctgaag tgtcaagtag ggggctttgg 240  
tgctgtcctt caggatgtaa tatatgtact aaaccagtga ccgaatacta tacagaatca 300  
gtagtaccta aaatacatgg atttttatca caaggcttag acatagaatc agcacttgta 360  
actatcaaat ggttgaggaa tttctacttc atttgtccac aattacgctg gattagaagt 420  
gtttgcatcc ttgcatctgt gtgt 444

<210> 115  
<211> 473  
<212> DNA  
<213> Homo sapiens

<400> 115  
tttgttacaa tattaagggt gtgtccaagg tccagagata gcatgtaaca ctaacaaatt 60  
ctgtgggatg gtggtgatgt caataccaag aaaagctttg cagagagctt ggggtttcag 120  
ccaagactcc acaaaggcat aggggctttg tgggagaatg gcagtcctcc tggagaagtg 180  
gcagataaaa aggtaaagat ctgtgagcaa cgtcatcttg agttcaggaa ttgacaatag 240

tttgggtatta gaagaagagt aagagtgtca aaaggagcat ttgtgtaatc ttctactcca 300  
gagatttttaa tctccttaat agaaagttgt ttgtattgat tgaatgatta acctttatta 360  
agaattttgt tgtctcaggc actggattag tagctttaca catttcattt aaatctcaca 420  
ttttgatagc ttctactatg gttattatct tacagaagaa actgaagtta aga 473

<210> 116  
<211> 261  
<212> DNA  
<213> Homo sapiens

<400> 116  
cctgaaacca tgggctcttc gtacctccag tgccgctcac atcttatgac acatagtagg 60  
ggcgtaata aatgcttatt aagttgacga ctatgccaga aaaaggggtga gggattacac 120  
aaagttttta caaaatctca cggtaactct tcagaagcaa aaataaaata ataacattta 180  
ataaaagtgc ctgctcaagg cctgcagccc aattccaggt ttgctccaaa tgttgatggc 240  
cttgagcttt ctgtgtgaa a 261

<210> 117  
<211> 193  
<212> DNA  
<213> Homo sapiens

<400> 117  
ctgctccatg gggatgggcc tcagtgagtg tatgtgccag gcttgaaatg gcttcacggt 60  
atggggttgca ggagcaccat gaggttcac tcattcttgc ctctctctgc cagcatgtgt 120  
gccatctgca atgtctcact gagcactgag tggggcctgc tatgtgggca gtatccctgc 180  
catcttcata tca 193

<210> 118  
<211> 364  
<212> DNA  
<213> Homo sapiens

<400> 118  
atctcattgg tatgtagttt tattttcctg aaaggtaatt aatcttggcc aaggagctaa 60  
aagtctaaat ctagagtgtc tagatgcttt gcaggatatc tggatttaaa tactcttggt 120  
ctcatgtatg ttctatcatc tccactctga aaaatgattt cttttgatgg aacagatagg 180  
aaaatactgt atagtgatta aaaatatggg ttctatagtt aggctacttg agttcaaacc 240  
ctggctctga cgctttctaa ctgtgtgact gtggacacga tatacaacct ctattaattt 300  
caatattacc atttgtgaga aaaggaatga taacaatatc catatcatgg tgggttcttt 360  
tttt 364

<210> 119  
<211> 425

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 119

```

agagatcttt aaaatactca aagaaaattg tcacctagaa tttgataact cttgaaaata    60
tcttgcaaaa atgaaggcta aataaatgat tttttgacaa agaaaagctg aaaaaattta    120
ttgtgagcag acctgtacta caagaaaggt taaaagaagt tatttaggta gaaagaaaat    180
gatatcaaat aagcagatct acacaaagga atgaagatct tcagaaatcg taaaattgtg    240
ggtaaatcta aaagccattt taaaaatttt gagtcattct aagattattg tctatagcaa    300
agaaaaatgc tagcaatttg ttatgagggt taaaatatgc agaagcagaa gtaaatcata    360
taatgatagc aacatgacaa ctgggggaaa atgaaagtcc actgaagaaa tgcttaataa    420
atgtt                                         425

```

&lt;210&gt; 120

&lt;211&gt; 438

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 120

```

actttccttt ccaggcattt cttgatgtgg aagagattta ctgagtctga tacctttaaa    60
ggtctgacaa gagacatttg ctgcctatgc cttctgttct cttggaggag tgctaccaat    120
aaggcttcgt caacataaca aggccacctt agctagacag gcctcttcct ttcttcctct    180
cataacctgt cttgccacta aacctgaatt accagcacia cctctttggg gccatgctct    240
gagcccatat tctttctata acctcaagta ggtatataag cttctgcgcc ttattgtctt    300
cattctgaag gctcttatgt acatgcatta aacaaatttg tatctcctat taatgtgcct    360
tttgcgagtt gatttttcag tgaaacttca gaggtccaac ggcagtagcc cctaccaagt    420
tcaagatgct ccacttac                                         438

```

&lt;210&gt; 121

&lt;211&gt; 482

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 121

```

gtgatgtaag actggtggac ttaaattaat tttttaaaagg catcatggga ttttgtatcg    60
gctatctctg tatctagaag atgtcagact catggaagtt ttgtccattt tattcccttt    120
gcttatccat tctttcttgt ttacagaaag acttaatttt ctgtctcata tctctgtcct    180
tcttgcccca ctatttttcc ccttctcca aaaatcccag ccccaaaaac agtctacata    240
ttgtgaaaaa gatttctcaa accacaaggg tgatgtaact ttaggcctgt gttttctctc    300
tcacacacac aaaatattgg atatgagtga gattttaaaa aattggtttt taaatgtgat    360
gaaaagagtg tccttttcac cagaacaaaa caacccttaa tgctgaagcc tccttcccga    420
tatgggtggc ttccaaatat gaagaaatct gtgcattggg ccacaggctc cagacaaagt    480

```

ct 482

<210> 122  
 <211> 568  
 <212> DNA  
 <213> Homo sapiens

<400> 122  
 ccttggcagc tccaacttga acatgtaaag ggtgtattca acagacaagt gagagaagga 60  
 acctcacaca gcctgagtgg gcctgagata ggctgagggg cctaagcttc aattgcataa 120  
 gcagggctag gtcactccag ttaccaaaga cagaaacaga tagtccagag cgtccaggg 180  
 gatgctagcc actgccagg agatgatcag agaacacaca acagaaatca gaaaatgtag 240  
 tacaagaaga atttgctgat aggtgcaatc gcctcagcaa ggcacaggaa actcaactca 300  
 gaaggcagtc tgtctgtcat ccaccaattc tctgggtcaa gtctgatgtg cactcataaa 360  
 gtaaaaatgc actgttattg tgactgagaa aaaaaataaa gctaaaaggt aagtgcctat 420  
 aaaataagat ttactaatg caaacaaaag ccctaaagaa gtgtgggttg agccagtggt 480  
 cctcctctat tagcaccaac aatggatagg tggttgagtc tgtcaaaatg cctctgggtt 540  
 tacagaaatg aaagcttggt ctgtgcc 568

<210> 123  
 <211> 413  
 <212> DNA  
 <213> Homo sapiens

<400> 123  
 cattttttac cacatatact ataagaatta gtattatttt tgattaaaaat aaatgttatt 60  
 ttcagaggtg caattttttg ctttcagtaa gattttcta ttaaggaagt catttttaag 120  
 gctaaattta aatgagaaaa agagcttggt gcaacttggt atccagttgg atccagtttt 180  
 ctctgctggt ccattttttg tatccctttt gagtttgc atctttttta catttttttg 240  
 tatagcagat tttttatttt tgggtacatt gtgcacataa acttcttggt gtggaggaga 300  
 ggttaaattt taatagctaa tgggacaaag gtatataggg atatataggt acaaccctag 360  
 ctcttattct ttcttttcct ccatagtatt ctggtgatgt agggataaaa ttt 413

<210> 124  
 <211> 525  
 <212> DNA  
 <213> Homo sapiens

<400> 124  
 ccaagcaaag ttatatattgt attttatttt acatttattt tgttatattc cttttatcta 60  
 cttaggtttc ttctctactt ccctttttta ttgaagagtt taatgcatgt atctgtgtgt 120  
 ttgcttgaaa aaaaacacca agtataacat gttctatcta tgaatacttc tggccattaa 180  
 ctcaaaaagg actatattac agacagaaaa gcaccagaaa gcaatcaggg acttcatcta 240

agaggtagga cagcatagtt ggtaaaaata cagaccctgg aggcaaactg cctgggcttg 300  
aatcccagct ttattacttt gggaaaacta cttatcttct ttacttgttt tggatccat 360  
gtctgtgaaa tggaagtaat aataatcttc tcatagcatt gttgtgaggt ttcaatagat 420  
gaagtgaaga ctttagaagg gcacatgata agaattatat aagggttacc tattattgct 480  
atccaatttg tcatagcaag ctaagggacc ttgggcaagt tactc 525

<210> 125  
<211> 575  
<212> DNA  
<213> Homo sapiens

<400> 125  
actggtagaa tgggctcatt caagcatgta acgcccttaa atttttcatt taaattttct 60  
gtgccttaga aatgaacttt acagtaatct ttgctttcta aaaataaatg tgtttcttgt 120  
taagcattta gtctcatcac aaattctggt ttagaaaaaa acaacagaaa atagtgaatg 180  
agaagggtag gagacttagg actcagcgaa ttctatctca gtgcccaagac tttaaaactg 240  
ggaataaatg ctacttctcc atgacctggg tctgataatt tgtctgcagg aacactgttt 300  
ctagaggggtg gtgtggtaca gtgggaggaa tggactttgg agtgagatcc atgttcaaatt 360  
cccaagtcac ttaccttctc tgatcctcag tttcctcatc tgtaaaatga ccataatcaa 420  
caccatctcg aagatttgtg gtgacaacac agcatttact tcctgctgta tacttcccat 480  
ttcctcttgt agagacagaa ttttccactt tattttaatc tataattatg taatcccat 540  
taaaaatcac ccttcgactt tcagttccac aaggc 575

<210> 126  
<211> 638  
<212> DNA  
<213> Homo sapiens

<400> 126  
attgctctct tctagatttt ctaatgttgg tcggtgccct tcgtaagttg tgtacaaagc 60  
tggatccagt actccaaggg tgatctgacc tcacagagca cagtgcctgg ggagtgccct 120  
taatctggac ttggaattcc atcatacaga ggccaagtct ctgaccatga tgttctctct 180  
gtgtaactgg ggctgctgaa acccaagtat tgtcagccag tgccggtctc cagccatgct 240  
tgtgtctttt aagaagtgac agtaactgct atttgtggag atggctattc atagggactc 300  
cttttctttg cctgacagag gccagtggt ctaagctcta agaggggctc tgatgccagc 360  
atgtgagtca cactcacttg ctactgttct tttccagagt tttgggccac ttgtgctgc 420  
acatcactac ctctctctcc cctgccagc ttgcattgtc gcccttcccc atctaccatg 480  
ctgtccttga acataaggcg cttctctgca ttccatgtgt ctactttgta gttatgtgct 540  
gcattttgaa agagctgaat ctatgtccag gttcaagaaa gaatgctgat caactgttgg 600

caatagatgg gtttaataata tcttatgatt ggttcttg 638

<210> 127  
<211> 573  
<212> DNA  
<213> Homo sapiens

<400> 127  
tagtctagac tctttttccc cttttaaggt cagctgatta accttaattc catctaatac 60  
cttgatttcc ctttgccatg tatgtcctgg ggaatgaggat gtggatggat ctaggggggc 120  
cgggtattctg gctaccatag ctatcttgct ctttttgttt ataattatga tatgttccaa 180  
aaaggagtaa aacgtaatac aagaagataa aaatacattt accattaagt aagaaaaaag 240  
acaagggaga agagaataag aaaatgagtc aggagtggga ttatacaaa aaattagtga 300  
gtccacttta ctctctggaa gtggatggtg agcttttctt gccagccttc ttgaagaggg 360  
aagcactgtc agttatgttg tagtgtgtcg atctagtaaa atccaactgg ttgttcagat 420  
acctagatga atattcttga taggaagatg aaaaaaaaat ttcttccaaa gtcttcatgg 480  
atacataaag tgtataatga gcaaacctt tgacatgttt acagtaaacc caatggtgtg 540  
tttcacctgg cctttctctt ctttcgttta ctg 573

<210> 128  
<211> 461  
<212> DNA  
<213> Homo sapiens

<400> 128  
catctattcg acgaccttga gttaccgctg agacatttct gaggcacaac actaagaaaa 60  
cgcatgtaat tgtcaagcgt ggcagggcag tattgtcttc aaagtcccgct ctgactgaca 120  
gggcagaggt tcttctcac tgcccgaatc tgcttcccgag cagctccagg gttccctcag 180  
gaagccgccc tccaccttca cctcaggcat gtctctgcaga gccctctgga gaaccagctt 240  
caggttctgc ctattttgac gctgcctaaa ggagcccacg aagaagtaaa tgacgggggtt 300  
ggcactaccg ttagaggag acaggaaaaat ggaaactaga tggacatgac agaaaatgac 360  
ttccaaatcc aggtgtatcc cagtagacag agcccaccga atgccgaagg gcaggctgcg 420  
gagtaggaag actagcactg tgagcaggat cgtcacgtac a 461

<210> 129  
<211> 655  
<212> DNA  
<213> Homo sapiens

<400> 129  
tcactggaga agcctagtca cctgggcaga atatcttgaa ctaggataa gttcatccat 60  
ggtagaccaa ctctgtgatg gaggttatgag atggggaagg agggctctggc accatgcaac 120  
aggatttccc ccaaagctca gactccaag gagcacatca gcatcaggaa tgtctgctgg 180

aagccagcgg ctgtggagga ggggcagtag ccactgagcc taggttcaga gcttcaatcc	240
ccttcagtcc tcttgactgg caagagaaca gcagagtcta ttagagagga attaccattc	300
caagcaagaa ttagggccac atctttcaga atgagaccat tgagttgagg tccacttagc	360
agggaaagtg gcttcaggtt gtggttgact gtttaattac accctgctgt tcaactctctt	420
caccattgta tgcaaagtac agcatctctg acaagcaagg aacactggct tgccccacag	480
tggttggtg gggttgatga aatgagcaac gaagtagcag tgtgcccagt ccaagcagag	540
actacctcta gcaggggcat gacattcccc aagagagggc atctccttta gcctggacct	600
tggagcaaaa gcaacccatg gatcagacca atagacaaca tgcagccctc atcta	655

<210> 130  
 <211> 657  
 <212> DNA  
 <213> Homo sapiens

<400> 130	
aagagttaga gcaggatttt accttgtttt acaaaaaaga aaagtttatt ttgaaaaaaa	60
ttccaacctt gcctcctccg aactatagtg aaaagataat ttccacatc cctttgttca	120
ggaaatgagg acacagtggg gtcatgggtt tttgattgtc cacttgaaa aggttaaaac	180
ctgtcctaca gtcgatga cttcagttcc atttaagtgg ggtcctgtct ctctcactct	240
ccaccgactg tacctttact ataacatggc cttatataga tagctttgag taagtgtgtg	300
ttaaagtact gcccaagtga atggaaaatt gagaagggcc tccagcactg gagtatggaa	360
aggagcactg ggttcattga ctctttggat ttctcccttg ctacgtaagt ccgttcccta	420
aaggacatgg atcttgacag tgttggaatc ttcagaaata attgcaatac cagaagttat	480
ttaagatttt accattttca aagtatttgt acgtaacact ttcatatgtt tttgtttcct	540
agctacctca gtttccctgt tggcttgagc agattagtgt aaagaggtag tgacatcagg	600
ggaaacaggt ttactcagcc atcttcatta ccatattatc actgacttga ggctcct	657

<210> 131  
 <211> 566  
 <212> DNA  
 <213> Homo sapiens

<400> 131	
tagtcgctgc tttctgttcc cgcttaaaga tggagatatt ttttccttcc atgcttgagg	60
agtctcgaaa gttttgcaca ctcttccacc tcctggaact tcaactgtgc attcaggggtg	120
actactgctg tctggctcca ctcgaggga gccaggtaac ctgtgttagg ccgcgctttt	180
cctggcggcc ttgtaaatct gttagtacat gaaaagcatg acgcacatgg ggattaggat	240
gccaatgcgg tggagtaaat cgtgtagcca aagtcttgac tgaccaagca caccttatca	300
tcgtttacat tctgagcccg accaaaaatg gtaggtaag tgacaaaggc ggaaagaagg	360
cagacagaaa gaatcatctt cgtcatgcat ttccccttct gcctcatagg gtacgtgaga	420



ggcttcatga tcccaaggta cctgtcgatg ctgatcacgt acaagggtcaa gatccaggcc 480  
 gtgcagcaca tgacattcac ggagaagacg ttacagaaaa agtgtccaaa gatccacttg 540  
 cccccgatga ggtcgggtgac actgat 566

<210> 132  
 <211> 575  
 <212> DNA  
 <213> Homo sapiens

<400> 132  
 agtgttacag ctgggcagcc agagagacag catgtagtcc tcattgaagc agaaagacag 60  
 agggttctga gacagaggtc tccaggaaaa aaaaaagaa cctgacttac tggataaaca 120  
 agtctttagt ttaaaaaaca acaaaaaact gtatacacat atatataaa aatcaggtag 180  
 tataaagaaa aacagaactc cagagattcc tgggtcacag aaggggaaag ggctgttcaa 240  
 gaaagtgaag ttgaactaac tgaaaataca gctatcttta tattggaag acagtcagga 300  
 agtcaacaga taaggcctaa actgcataaa gcaggaaaca gcagactaaa gacattatta 360  
 agaaatatgg aacacaacca aaagaaatag caaaaacaat gaaaagtgc tgtttttcat 420  
 aagtgaggca ggggaagaga aggggttatt tttttcccca ttatatgtct ttaagaacta 480  
 cttgctaaaa atattgggca catatgaatt tgataaaagc gaaaaacttt ttacttcaca 540  
 agtgcagctt taacatacgt tgattacagt gaagt 575

<210> 133  
 <211> 651  
 <212> DNA  
 <213> Homo sapiens

<400> 133  
 aaagggtgaca gagaagtagg tgaggaattc agttttaaat ttattcattt ttaagttgtg 60  
 tcaggctctcc ccaagattat ccctcgggtc tgtgattcat aggacttagc atatagttgt 120  
 attcacagct atgacttatt aacagaggga taccgaagca taatcagcaa aaggaaaaga 180  
 tgcatgagga aaagtctgaa gaaaccaggg acagcttcca agattctttt cccagtgaag 240  
 ttacacagga tatgcttaat tctttcagca aggaattgtg acaagacatg tgaaacacta 300  
 cctgccaggg aagttcctta gtgactcagt gcccatggtt attattggg actggtcacg 360  
 tatgccctct ttgcctcata cttagagaat tccagttcca gaaggaaagc aggtattcag 420  
 tataagccat attatttgca tagaccagtt taggatcaag gaattgtagg aagcttttca 480  
 aaatctaaga ccccaaatac cagccaagag ccagccttgc aagcaggaca ttttaagagt 540  
 agcagtcttg ggtctgctgt attaactctt tctgcacag aaatgatagt atgacatcta 600  
 agttattatt atcaaggagc cgagaaatgc atgtttttta ggctaggga g 651

<210> 134

<211> 966  
 <212> DNA  
 <213> Homo sapiens

<400> 134  
 atgaaccaga ctttgaatag cagtgggacc gtggagtcag ccctaaacta ttccagaggg 60  
 agcacagtgc acacggccta cctgggtgctg agctccctgg ccatgttcac ctgcctgtgc 120  
 gggatggcag gcaacagcat ggtgatctgg ctgctgggct ttcgaatgca caggaacccc 180  
 ttctgcatct atatcctcaa cctggcggca gccgacctcc tcttcctctt cagcatggct 240  
 tccacgctca gcctggaaac ccagcccctg gtcaatacca ctgacaaggt ccacgagctg 300  
 atgaagagac tgatgtactt tgcctacaca gtgggcctga gcctgctgac ggccatcagc 360  
 acccagcgct gtctctctgt cctcttcctt atctggttca agtgtcaccg gcccaggcac 420  
 ctgtcagcct ggggtgtgtg cctgctgtgg acactctgtc tcctgatgaa cgggttgacc 480  
 tcttcctctt gcagcaagtt cttgaaattc aatgaagatc ggtgcttcag ggtggacatg 540  
 gtccaggccg cctcatcat ggggggtctta accccagtga tgactctgtc cagcctgacc 600  
 ctctttgtct ggggtcggag gagctcccag cagtggcggc gccagccac acggctgttc 660  
 gtggtgttcc tggcctctgt cctgggtgtc ctcatctgtt ccctgcctct gagcatctac 720  
 tggtttgtgc tctactggtt gagcctgccg cccgagatgc aggtcctgtg cttcagcttg 780  
 tcacgcctct cctcgctccg aagcagcagc gccaaacccg tcactactt cctgggtggc 840  
 agccggagga gccacaggct gccacaccagg tcctgggga ctgtgctcca acaggcgctt 900  
 cgcgaggagc ccgagctgga aggtggggag acgccaccg tgggcaccaa tgagatggg 960  
 gcttga 966

<210> 135  
 <211> 198  
 <212> PRT  
 <213> Homo sapiens

<400> 135

Lys Lys Gln Val Ser Leu Thr Glu Gln Glu Thr Ile Leu His Phe Phe  
 1 5 10 15  
 Lys Trp Gly Lys Thr Glu Gln Leu His Glu Lys Tyr Asn Ser Leu Tyr  
 20 25 30  
 Ile Lys Leu Ile Gly His Glu Leu Ala Leu Gln Val Glu His Asn Asn  
 35 40 45  
 Ser Arg Ser Lys Ser Arg Leu Pro Ser Lys Ser Cys Ser Ile Arg Arg  
 50 55 60  
 Phe Phe Ile Gln Asp Ala Lys Ile Ile Lys His Asn Asn Cys Ile Glu  
 65 70 75 80  
 Leu Asn Glu Asn Arg Gln Cys Phe Ile Ile Glu Lys Phe Ser Asp His  
 85 90 95

His Ala Lys Ile Phe Leu Ile Phe Asn Phe Leu Cys Arg Ile Ile Phe  
 100 105 110  
 Met Ser Met Gly Tyr Phe Glu Tyr Arg Arg Ala Met Cys Asn Asn Tyr  
 115 120 125  
 Ile Arg Val Asn Ile Val Ser Ile Thr Ser Ser Val Tyr His Leu Cys  
 130 135 140  
 Tyr Lys Gln Ser Ser Tyr Ile Leu Leu Val Ile Leu Asn Cys Thr Thr  
 145 150 155 160  
 Lys Leu Tyr Leu Gln Ser Pro Cys Cys Ala Ile Tyr Ile Leu Phe Ile  
 165 170 175  
 Phe Phe Leu Thr Ile Phe Cys Thr His Pro Ser Ser Leu Tyr Ser Pro  
 180 185 190  
 Ser Ala Gln Leu Asn Ser  
 195

<210> 136  
 <211> 214  
 <212> PRT  
 <213> Homo sapiens

<400> 136

Arg Cys Ser Ile Val Ser Ser Val Ser Cys Pro Leu Leu Pro Pro Gly  
 1 5 10 15  
 Val Asp Ser Cys Thr Val His Pro Thr Pro Ala Phe Pro Ser Phe Leu  
 20 25 30  
 Ile Ser Pro Val Ile Phe Pro Val Ala Leu Leu Cys Trp Cys Pro Val  
 35 40 45  
 Arg Ser Cys Gly His Lys Arg Leu His Gly Pro His Pro Gln Leu Gly  
 50 55 60  
 Glu Ser Ser Pro Ser Trp Val Leu Trp Thr Val Lys Lys Asp Gly His  
 65 70 75 80  
 Val Gly Ser Val Glu His Glu Val Val Gln Asp Leu Gly Gly His Arg  
 85 90 95  
 Ser Cys Leu Pro Ala Ser Arg Ala Leu Pro Pro Phe Gly Ser Leu Leu  
 100 105 110  
 His Leu Gly Lys Arg Phe Val Pro Thr Pro Arg Arg Val Asn Arg Ala  
 115 120 125  
 Pro Trp Trp Ser Thr His Cys Pro Ser Glu Gly Pro Ser Ser Leu Met  
 130 135 140  
 Ser Trp Cys Pro Gly Leu Pro Gly Arg Ile Leu Ala Ala Leu Pro Gly  
 145 150 155 160  
 Pro Glu Met Asn His Trp Glu Glu Ile Gly Asn Glu His Thr Ala Ala  
 165 170 175  
 Thr Leu His Pro Asn Pro Val Pro Tyr His Arg Arg Leu Leu Trp Gln  
 180 185 190  
 Asp Asp Ser Ile Ser Val Cys Leu Arg Ser Leu Phe Leu Pro Arg Leu

195                      200                      205  
 Leu Pro Pro Gly Arg His  
 210  
 <210> 137  
 <211> 141  
 <212> PRT  
 <213> Homo sapiens  
 <400> 137  
 Ile Ile Ser His Thr Ala Phe Phe Arg Phe Ser Leu Ser Ile Cys Phe  
 1                      5                      10                      15  
 Cys Asn Ser Tyr Trp Thr Phe Thr Ser Leu Ser His Cys Leu Leu Tyr  
 20                      25                      30  
 Leu Leu Thr Phe Val Phe Ser Val Ser His Cys Cys Ile Val Ser Tyr  
 35                      40                      45  
 Tyr Leu Ala Leu Pro Val Asn Ser Leu Ser Phe Phe Cys Asn Leu Phe  
 50                      55                      60  
 Ile Ser Ser Leu Cys Leu Leu Phe Gln Leu Asn Leu Ile Ala Gln Ser  
 65                      70                      75                      80  
 Phe Ile Trp Ser Phe Lys Ile Cys Phe Cys Leu His Ser Tyr Phe Val  
 85                      90                      95  
 Leu Phe Ser Leu Ser Leu Tyr Leu Phe Leu Met Leu Ser Ser Ala Tyr  
 100                      105                      110  
 Tyr Phe Asp Ile Tyr Phe Leu Ala Ser Leu Arg Tyr Ser Ile Ile Ser  
 115                      120                      125  
 Gly Pro Arg Ile Ile Lys Ser Pro Thr Thr Ser Val Asp  
 130                      135                      140  
 <210> 138  
 <211> 223  
 <212> PRT  
 <213> Homo sapiens  
 <400> 138  
 His Glu Trp Leu Thr Phe Phe Ile Glu Asp Glu Ile Leu Ser Trp Cys  
 1                      5                      10                      15  
 Ile Tyr Val Pro Cys Tyr Phe Pro Ala Asn His Phe Ser Asn Thr Ala  
 20                      25                      30  
 Gln Leu Tyr Ser Asp Thr Val Asp Thr Val Phe Gln Ala Leu Tyr Phe  
 35                      40                      45  
 Gln Phe Ile Cys Gly Ile Leu Asp Ser Phe Gly Ser Ser Thr Glu Val  
 50                      55                      60  
 Thr Phe Ile Tyr Arg His Phe Arg Gly Ile His Thr Thr Ser Tyr Asn  
 65                      70                      75                      80  
 Cys Thr Ala Ile Ala Cys His Cys His Val Phe Ile Asn Phe Gln Phe  
 85                      90                      95  
 Leu Glu Asp Phe Ser Ile Ile Ile Tyr Lys Leu Val Lys Phe Thr Val

100                      105                      110  
 Ile Cys Gln His Leu Glu Gln Glu Lys Met Ser Ala Lys Asp Gly Arg  
     115                      120                      125  
 Thr Leu Tyr Phe Ile Leu Ile Ala Gly Phe Leu Pro Asp Asp Asn Phe  
     130                      135                      140  
 Gln Lys Ile Asn Pro Asn Phe Asn Thr Ser Cys His His Phe Thr His  
     145                      150                      155                      160  
 Ser Asn Ile Lys Ile Ser Asn Phe Thr Tyr Ile Ser Ser Glu Ser Thr  
                     165                      170                      175  
 Asp Lys Leu Phe Tyr Ile Glu Gly Asn Ile Ser Trp Glu Val His Asn  
                     180                      185                      190  
 Cys Thr Cys Arg Ile Ile His Arg Ser Phe Gln Val Leu Leu Leu Gln  
     195                      200                      205  
 Ile Gly Leu Lys Ser Ile Thr Val Gly Leu Ser Val Ala Gln Lys  
     210                      215                      220

<210> 139  
 <211> 173  
 <212> PRT  
 <213> Homo sapiens

<400> 139

Asn Ile Ile Thr Phe Phe Tyr Glu Tyr Ser Trp Ser Phe Gln Asn Lys  
 1                      5                      10                      15  
 Thr Ser Tyr Trp Phe Asn Lys Leu Trp Tyr Asn Gln Ile Met Lys Leu  
                     20                      25                      30  
 Tyr Ala Phe Val Lys Val Thr Phe Gln Lys Asn Ile Leu His Arg Ile  
                     35                      40                      45  
 Thr Asp Pro Ser Ala Leu Pro Thr Leu Trp Ala Leu Ser Leu Phe His  
     50                      55                      60  
 His His Tyr Leu His His Cys Leu Gln Val Phe Tyr Thr Ala Arg Val  
     65                      70                      75                      80  
 Gly Leu Cys Leu Leu Asn Ser Gln Val Lys Arg Gly Arg Lys Leu Thr  
                     85                      90                      95  
 Pro Ser Gly Gly Ser Leu Gly Met Ile His Gly Arg Trp Ser Ile Asn  
                     100                      105                      110  
 Thr Ser Ala Leu Phe Pro Leu Glu Ile Leu Arg Asn Gly Phe Tyr Ile  
     115                      120                      125  
 Val Ser Gln Ser Phe Leu Lys Val Leu Asn Phe Asn His Pro Gln Gly  
     130                      135                      140  
 Val Val Gly Phe Ile Ile Val Tyr Ile Pro Leu Trp Leu Pro Phe Leu  
     145                      150                      155                      160  
 Leu Val Ser Leu Leu His Ser Lys Leu Gly Phe Ile Ser  
                     165                      170

<210> 140  
 <211> 223

<212> PRT  
<213> Homo sapiens

<400> 140

```

Val Phe Leu Ser Arg Lys Glu Glu Lys Gly Trp Val Val Thr Gly Gly
1      5      10      15
Gln Gln Cys Gln Asn Trp Gly Val Trp Thr Gly Ile Gln Glu Asn Glu
20      25      30
Gly Ala Gln Asp Glu Gln Lys Gly Gly Glu Ala Ile Phe Ile Lys His
35      40      45
Leu Leu Cys Ala Ser Gln Ala Arg Leu Gln Ile Ile Thr Leu Leu Lys
50      55      60
Ser Ser Gln Gln Pro Ser Asn Arg Tyr Leu Ser Leu Ile Pro Tyr Pro
65      70      75      80
Cys Ser Ala Ser Pro Pro Ile Thr Met Ala Glu Glu Phe Lys Pro Leu
85      90      95
Ser Lys Ala Ser Thr Val Ile Cys Pro Leu Asp Pro Ile Pro Ser Ile
100     105     110
Phe Leu Phe Ile Glu Thr Phe Ser Met Val Phe Lys His Thr Leu Leu
115     120     125
Ser Leu Leu Leu Asn Arg Gln Met Gln Leu Ile Lys Leu Phe Phe Ser
130     135     140
Leu Gly Tyr Cys Pro Ile Ser Leu Leu Pro Phe Met Ala Glu Leu Leu
145     150     155     160
Glu Arg Val Phe His Asn His Phe Ile Ser Thr Pro Leu Thr Asp Phe
165     170     175
Thr Gln Leu Glu Glu Glu Gly Thr Leu Ile Pro Lys Cys Pro Ile
180     185     190
Lys Pro Asn Pro Leu Lys Val Leu Cys Cys His Asp Gly Cys Glu His
195     200     205
Gly Glu Lys Ile Leu Glu Asp Val Gly Asn His Asp Arg Glu Thr
210     215     220

```

<210> 141  
<211> 176  
<212> PRT  
<213> Homo sapiens

<400> 141

```

Ser Cys Glu Thr Ser Ile Leu Val Ser Trp Gly Gln Gly Asn Gln Gly
1      5      10      15
Pro Ser Met Leu Ile Leu Pro Cys Val Arg Leu Ile Leu Ser Ile Ser
20      25      30
Gly Gly Gln Val Ala Thr Trp Pro Pro Gly His Thr His Gln Glu Phe
35      40      45
Ile Leu Cys Asn Leu Glu Glu Gly Leu Arg Asn Ala Gly Gly Tyr Leu
50      55      60

```

Pro Gly Asp Ile Leu Tyr Pro Leu Ile Gly Asn Trp Gly Arg Ser Gln  
 65 70 75 80  
 Phe Gly His Thr Phe Pro Glu Leu Asn Phe Tyr Glu Gly Asp Leu Gly  
 85 90 95  
 Gly Arg Gly Ser Glu Ala Asn Ile Ala His Val Pro Gln Thr Leu Val  
 100 105 110  
 Cys Leu Thr Glu Ile Tyr Ile Phe Ser Asp Lys Phe Phe Lys Ser Leu  
 115 120 125  
 Leu Tyr Val Phe Arg Thr Ile Ser Gly Asp Phe Leu Lys Asn Asn Phe  
 130 135 140  
 Cys Leu Leu Tyr Leu Phe Ser Ala Val Thr Gly Pro Gln Ser Pro Tyr  
 145 150 155 160  
 Asn Val Asn Pro Glu Val Glu Leu Leu His Tyr Ser Phe Phe Phe Phe  
 165 170 175

<210> 142  
 <211> 209  
 <212> PRT  
 <213> Homo sapiens

<400> 142

Ser Gln Lys Asn Thr Thr Pro Leu Leu Glu His Asn Val Ile His Phe  
 1 5 10 15  
 His Leu Leu Ala Ser Leu Ala Glu Phe Gln Lys Cys Asn His Tyr Glu  
 20 25 30  
 Ala Gly Thr Lys Asp Phe Pro Asn His Phe Val Ile Leu Ile Asn Ile  
 35 40 45  
 Ser Ser Ile Leu Leu Asp Pro Phe Thr His Phe Leu Tyr Cys Phe Pro  
 50 55 60  
 Phe Pro Glu Val Leu Asn Lys Ile Ser Leu Leu Phe Val Leu Glu Lys  
 65 70 75 80  
 Ser Ser Cys Leu Pro His Arg Met Val Val Gly Glu Thr Gln Trp Glu  
 85 90 95  
 Thr Ser Val Lys Gly Gln Lys Thr Leu Thr Phe Val Ile Val Ser Ser  
 100 105 110  
 Phe Phe Gln Asn Thr Ser Ile Ala Trp Leu Leu Tyr Thr Arg Leu Leu  
 115 120 125  
 Lys Ile Tyr Leu Cys Pro Thr Thr Leu Phe Val Val Asn Ile Phe Leu  
 130 135 140  
 Ile Leu Ile Gln Tyr Ile Ser Glu Ile Phe Asp Leu Gln Ser Asn Leu  
 145 150 155 160  
 Ser Ile Thr Met Ile Pro Tyr Leu Asn Thr Gly Met Val Lys Met Arg  
 165 170 175  
 Thr Asn Leu Pro Phe Leu Cys Ser Tyr Arg Gln Ala Ile Leu Ile Thr  
 180 185 190

Asn Val Gln Ser Lys Pro Met His Glu Cys Arg Met Gln Leu Lys Ser  
195 200 205

Arg

<210> 143  
<211> 200  
<212> PRT  
<213> Homo sapiens

<400> 143

Ser Phe Pro Val Ser Glu Lys Ile Lys Pro Cys His Ser Lys His Val  
1 5 10 15  
Leu Pro Lys Phe Lys Lys His Val Asn Leu Leu Val Lys Leu Tyr Val  
20 25 30  
Leu Val Asp Phe Glu Ile Leu Cys Asn His Leu Lys Leu Ala Ser Gly  
35 40 45  
Pro Gln Leu Asp Gln Ile Pro Val Ser Leu Phe Leu Thr Ser Leu Cys  
50 55 60  
Trp Thr Thr Tyr Leu Gln Arg Gln Lys Lys Asp Lys Ser Asn Asn Pro  
65 70 75 80  
Thr Val Ile Leu His Lys Ser Met Thr Lys Leu Pro Leu Gln Lys Leu  
85 90 95  
Asn Ser Ser Ser Leu Asn Phe Leu Thr Ile Thr Trp Lys Ser Ala Thr  
100 105 110  
Met Val Asn Cys Gln Thr Cys Thr Ala Ser Gln Pro Thr Leu Tyr Thr  
115 120 125  
Asn Lys Gly Gly Leu Tyr Ser Asp His Tyr Trp Asn Lys Leu Ser Leu  
130 135 140  
Pro Asn Val Ser Ser His Pro Leu Asn Tyr Leu Leu Leu Tyr Phe  
145 150 155 160  
Tyr Thr Ala Ile Lys Leu Lys Leu Leu Lys His Asn Phe Ala His Val  
165 170 175  
Gln Asn Phe Tyr Ser Val Pro Gln Gln Ser Leu Thr Asn Pro Gln Asn  
180 185 190  
Leu Pro Thr Asn Leu Phe Leu Thr  
195 200

<210> 144  
<211> 170  
<212> PRT  
<213> Homo sapiens

<400> 144

Val Ile Pro Ser Ser Val Cys Pro Thr Val Gly Leu Pro Asp Thr Asp  
1 5 10 15  
Ser Thr Thr Leu Val Ile Cys Asp Phe Leu Phe Thr Gly His Glu Lys  
20 25 30



Pro Phe Thr Asp Trp Leu Gln Cys Ala Ser Leu Pro Tyr Gln Leu Leu  
 35 40 45  
 Phe His Thr Asn Ser His Leu Val Asn Trp Val Pro Cys Ser Ala Lys  
 50 55 60  
 Met Cys Phe Ser Ala Gln Val Ile Leu Tyr Thr Pro Ile Leu Asn Leu  
 65 70 75 80  
 Leu Cys Ala Ser Gln Ser Thr Ile Phe Gln Ser Gln Leu Lys Pro Phe  
 85 90 95  
 Ile Ile Gln Tyr Gly Phe Ser Pro Gln Ser His Val Lys Val Ser Pro  
 100 105 110  
 Cys Phe Phe Gln Thr Val Val Ala Leu Thr Gly Leu Leu Leu Gly Tyr  
 115 120 125  
 Lys Leu Thr Leu Tyr Phe Ser Ile Phe Ser Leu Pro Trp Ser Lys Arg  
 130 135 140  
 Lys Ile Arg Ser Met Asn Leu Arg Thr Tyr Lys Leu Leu Val Glu Gln  
 145 150 155 160  
 Gly Leu Asp Ile Val Cys Ile Asp Ser Arg  
 165 170

<210> 145  
 <211> 214  
 <212> PRT  
 <213> Homo sapiens

<400> 145

Met Gly Thr Ala Leu Phe Lys Val His Phe Pro Asp Ser Ala Val Leu  
 1 5 10 15  
 Phe Ser Ser Ser Ile Pro Thr Asn Ser Gly Leu Gln Ala Phe Pro Leu  
 20 25 30  
 Leu Ser His Ser Ile Leu Pro Glu Pro Ser Ile Lys Ala Pro Thr Ile  
 35 40 45  
 Leu Pro Ser Gly Gly Ala Ile Phe Leu Ser Phe Pro Glu Arg Trp Asp  
 50 55 60  
 Pro Leu His Phe Thr His Leu Ser Pro Arg Pro Ser Thr Cys Leu Ala  
 65 70 75 80  
 Gln His Ser Asn Ile Asn Pro Val Glu Ile Asn Cys Gly Ile Ala Trp  
 85 90 95  
 Phe Pro Trp Met Val Ile Gln Val Val His Cys Thr Thr Met Cys Asn  
 100 105 110  
 Ile Pro Gly Lys Arg Gln Lys Phe Ile Asp Trp Leu Gly Val Leu Asn  
 115 120 125  
 Ser Gln Gly Lys Leu Phe Asp His Cys Met Pro Ser Thr Trp Glu Asn  
 130 135 140  
 His Ile Pro Gln Leu Leu Arg Pro Tyr Cys Met Val Thr Trp Gly Asn  
 145 150 155 160  
 Ile His Thr Val Ser Pro Ala Leu Ser Ala His Lys Gly Asp Ile Val

165                      170                      175  
 Gln Arg Gly Asn Leu Ser Leu Pro Ser Thr Ser Leu Phe Leu Thr Pro  
                     180                      185                      190  
 Lys Ser Leu Ser Leu Leu Thr Lys Asp Ile Ser Ala Ser Ala Ile Leu  
                     195                      200                      205  
 Phe Ala Glu Trp Arg Ile  
                     210

<210> 146  
 <211> 200  
 <212> PRT  
 <213> Homo sapiens

<400> 146

Arg Ile Ser Gln Lys Cys Cys Val Leu Leu His Pro Leu Trp Gln Leu  
 1                      5                      10                      15  
 Phe Val Tyr Leu Ser His Ala Gly Glu Val Asn Thr Asp Pro Leu Val  
                     20                      25                      30  
 Lys Met Met Ser Asp Ile Phe Phe Ser Ala Ala Asn Leu Ser Ile Phe  
                     35                      40                      45  
 Ser Phe Val Ile Met Gly Ile Leu Trp Lys Val Thr Trp Arg Leu Cys  
                     50                      55                      60  
 Lys Ile Tyr Ser Ser Gln Phe Tyr Leu Pro Val Leu Ala Ser Ile Asp  
 65                      70                      75                      80  
 Val Ser Cys Leu Ser Leu Leu Ala Gln Phe Ala Lys Cys His Tyr Leu  
                     85                      90                      95  
 Pro Phe Ser Ser Met Arg Cys Met Tyr Val Tyr Met Tyr Ile Cys Ile  
                     100                      105                      110  
 Asp Ile Ser Val Tyr Leu Glu Thr Tyr Ile Asp Glu Leu Ser Ile Thr  
                     115                      120                      125  
 Met Ile Ile Tyr Phe Asp Val Gln Val Val Pro Asp Leu Thr Ser Asp  
                     130                      135                      140  
 Ser Phe Leu Asn Leu Met Tyr Gln Asp Val His Lys His Val Phe Phe  
 145                      150                      155                      160  
 Pro Cys Pro Asn His Pro Gly Val Gly His Leu Ser Lys Met Ser Cys  
                     165                      170                      175  
 Phe Cys Leu Leu Arg Trp Arg Ser Gly Ile Gln Lys Ser Arg Ser Val  
                     180                      185                      190  
 Cys Leu Val Cys Phe Ile Ala Ile  
                     195                      200

<210> 147  
 <211> 191  
 <212> PRT  
 <213> Homo sapiens

<400> 147

Tyr Leu Ile Leu Lys Tyr Ile Ile Met Lys Ser Ile Asn Val Ser Arg

1                      5                      10                      15  
 Gln Arg Ser Tyr Ile Pro Lys Ile Gly Asn Asn Cys Val His Met Cys  
                     20                      25                      30  
 Tyr His Thr Ile His Pro Ile Leu Leu Tyr Leu Asn Phe Pro Lys Gln  
                     35                      40                      45  
 Pro Val Val Lys Gln Leu Val Met Arg Thr Asn Glu Lys Leu Pro Glu  
                     50                      55                      60  
 Ile Ser Asp Ser Ser Cys Thr Tyr Phe Thr Pro Glu Val Trp Glu Phe  
                     65                      70                      75                      80  
 Thr Glu His Asn Val Arg Phe Phe Ser Ile Ser Tyr Pro Leu Pro Lys  
                     85                      90                      95  
 Ile Val His Lys Ile Gln Asn Ile Ser Ser Leu Thr Phe Leu Glu Cys  
                     100                      105                      110  
 Asn His Thr Leu Asp Asn Tyr Phe Arg Leu Leu Asn Gly Lys Arg Thr  
                     115                      120                      125  
 Gly Arg Arg Val Lys Val Thr Cys Phe His Leu Ser Tyr Phe Arg Leu  
                     130                      135                      140  
 Thr Ser Lys Ser Phe Phe Thr Leu Phe Leu Ile Leu His Arg Pro Phe  
                     145                      150                      155                      160  
 Leu Val Lys Ser Ala Asp Ser Lys Tyr Lys Ala Asn Ala Tyr Ser Tyr  
                     165                      170                      175  
 Val Ile Phe Met Phe Phe Lys Asn Asn Met Val Leu Thr Ser Ser  
                     180                      185                      190

<210> 148  
 <211> 193  
 <212> PRT  
 <213> Homo sapiens

<400> 148

Gly Leu Ser Glu Gly Glu Ala Ser Leu His Leu Asp Phe Phe Leu Lys  
 1                      5                      10                      15  
 Ile Thr Thr Ile Met Asn Thr Ala Ala Thr Ser Leu Leu Cys Thr Arg  
                     20                      25                      30  
 Gly Ile Ile Leu Gly Val Ser Val Tyr Ala Tyr Pro Glu Ile Ser Ser  
                     35                      40                      45  
 Phe Leu Leu Arg Gly Glu Val Leu His Ile Asp Phe Ile Val Arg Asn  
                     50                      55                      60  
 Gly Lys Ile Phe Asn Lys Cys Ile Arg Ala Thr Thr Phe Ser Ala Leu  
                     65                      70                      75                      80  
 Gln Pro Ala Ser Pro Pro Ser Arg Gln Asp Ile Met Asn Pro Leu Phe  
                     85                      90                      95  
 Gly Lys Ala Ala Glu Lys His Val Leu Gln Thr Tyr Tyr His Leu Val  
                     100                      105                      110  
 Asn Asn Ser Gln Trp Thr Asp Gln Asn Ser Arg Arg Phe Pro Leu Ser  
                     115                      120                      125

Leu His Cys Thr Asp Ala Ala Thr His Ala His Ile Pro Leu Asn Leu  
130 135 140

Pro Val Thr Thr Ala Gln Arg Gln Leu Ser Ser Trp Ala Gln Asn His  
145 150 155 160

Trp Gly Thr Phe Trp Gln Leu Ala Asn His Cys Ala Gln Arg Gln Ser  
165 170 175

Gln Phe Thr Leu Pro Gln Arg Gly Thr Glu Tyr Thr Ala His Pro His  
180 185 190

Leu

<210> 149

<211> 195

<212> PRT

<213> Homo sapiens

<400> 149

Ile Leu Asp Ser Phe Arg Asp Phe Leu Glu Gln Gly Gln Glu Ser Phe  
1 5 10 15

Leu Asp Lys Val Arg Ser Asp Leu Ser Gln Gly Arg Ser Ile Phe Ser  
20 25 30

Tyr Thr Arg Arg Asn Phe His His Lys Gln Cys Pro Lys Asp Ala Cys  
35 40 45

Tyr His Phe Tyr Ser Met Leu Phe Ser Val Phe Trp Pro Ile Leu Leu  
50 55 60

Glu Ile Gln Val Arg Lys Met Thr Lys Gly Ile His Glu Thr Arg Ser  
65 70 75 80

Leu Phe Arg Arg Trp Tyr Asp Cys Leu Ser Arg Lys Lys Glu Met Thr  
85 90 95

Pro Ser Phe Trp Glu Phe Thr Asn Ser Gly Trp Val Leu Asp Lys His  
100 105 110

Leu Lys Asn Gln Ser Phe Pro Cys Val Ala Ala Ile Thr Ile Lys Met  
115 120 125

Glu Met Arg Ser Gly Ala Val Asn Ile Gln Gln Glu Leu Leu Ile Cys  
130 135 140

Arg Pro Asp Lys Ser Pro Pro Glu Trp Thr Pro Ala Arg Glu Gly Arg  
145 150 155 160

Ser Leu Glu Gly Arg Arg Glu Asp Thr Glu Asp Leu Pro Leu Pro Gln  
165 170 175

Glu Ala Pro Arg Glu Arg Ala Thr Thr Val Tyr Ser Ser Arg Leu Trp  
180 185 190

Gly Asp Ser  
195

<210> 150

<211> 168

<212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 150

Leu Lys Ser Ser Gln Gln Pro Ser Asn Arg Tyr Leu Ser Leu Ile Pro  
 1 5 10 15  
 Tyr Pro Cys Ser Ala Ser Pro Pro Ile Thr Met Ala Glu Glu Phe Lys  
 20 25 30  
 Pro Leu Ser Lys Ala Ser Thr Val Ile Cys Pro Leu Asp Pro Ile Pro  
 35 40 45  
 Ser Ile Phe Leu Phe Ile Glu Thr Phe Ser Met Val Phe Lys His Thr  
 50 55 60  
 Leu Leu Ser Leu Leu Leu Asn Arg Gln Met Gln Leu Ile Lys Leu Phe  
 65 70 75 80  
 Phe Ser Leu Gly Tyr Cys Pro Ile Ser Leu Leu Pro Phe Met Ala Glu  
 85 90 95  
 Leu Leu Glu Arg Val Phe His Asn His Phe Ile Ser Thr Pro Leu Thr  
 100 105 110  
 Asp Phe Thr Gln Leu Glu Glu Glu Glu Gly Thr Leu Ile Pro Lys Cys  
 115 120 125  
 Pro Ile Lys Pro Asn Pro Leu Lys Val Leu Cys Cys His Asp Gly Cys  
 130 135 140  
 Glu His Gly Glu Lys Ile Leu Glu Asp Val Gly Asn His Asp Arg Glu  
 145 150 155 160  
 Thr Glu Lys Val Val Lys Gly Phe  
 165

&lt;210&gt; 151

&lt;211&gt; 121

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 151

Thr Gly His Pro Arg Leu Pro Pro Thr Leu Lys Gln Pro Ala Arg Gln  
 1 5 10 15  
 Cys Val Thr Tyr Gly Phe Asn Ser Asp Glu Glu Asp Ser Ser Trp His  
 20 25 30  
 Gly Leu Leu Arg Thr Leu Asn His Lys Val Ser Arg Asp Arg Arg Thr  
 35 40 45  
 Val Pro Thr Ala Ala Thr Pro Arg Trp Val Cys Ser Pro Val Ala Thr  
 50 55 60  
 Leu Lys Phe Leu Lys Thr Phe Tyr Gly Val Leu Leu Cys His Leu Gly  
 65 70 75 80  
 Trp Ser Ala Val Thr Cys Leu Ile Pro His Leu Ala Glu Thr His Arg  
 85 90 95  
 Arg Ser Leu Val Arg Thr Arg Glu Gly Ala Gly His Ser Gly Ser Cys  
 100 105 110

Gln His Phe Gly Arg Leu Arg Gln Glu  
115 120

<210> 152  
<211> 211  
<212> PRT  
<213> Homo sapiens

<400> 152

Leu Val Ala Ile Ser Leu Lys Phe Phe Phe Cys Arg Lys Ile Ser His  
1 5 10 15

Arg Trp Leu Ile Ile Cys His Ile Lys Pro Leu Arg Lys Lys Gly Trp  
20 25 30

Gln Met Leu Leu Leu Val Arg Leu Leu Cys Tyr Glu Ile Trp Val Lys  
35 40 45

Cys Ala Gly Val Thr Glu Glu Gly Glu Phe Leu Ser Pro Ser Arg Ile  
50 55 60

Glu Glu Asn Gly Val Arg Asp Arg Glu Gln Leu Ala Arg Lys Ala Gln  
65 70 75 80

Gly Val Asn Leu Thr Arg Lys Phe Lys Gln Trp Leu Leu Leu Tyr Ser  
85 90 95

Leu Phe Val Gln Ile Leu Lys Met Lys Leu Phe Ile Lys Phe Ile Val  
100 105 110

Val Phe Leu Asn Ser Met Arg Asn Gly Arg Asn Leu Arg Tyr Cys Ser  
115 120 125

Lys Gly Ser Ser Ala Pro Asn Leu Phe Leu Thr Lys Phe Ile Leu Leu  
130 135 140

Pro Lys Val Ser Pro Asn Val Thr Pro Thr Ser Ile Arg Gln Glu Tyr  
145 150 155 160

Cys Asn Glu Ala Met Thr Ile His Asn Leu Leu Ser Ile Lys Gln Val  
165 170 175

His Glu Arg Phe Cys Asn Asn Thr Leu Cys Lys Ser Leu Trp Asn Asn  
180 185 190

Asn Lys Ile Asp Val His Phe Met Tyr Tyr Cys Ile Leu His Ile Leu  
195 200 205

Arg His Glu  
210

<210> 153  
<211> 173  
<212> PRT  
<213> Homo sapiens

<400> 153

Val Asp His Trp Ile His Leu Asp Met Phe Lys Met Phe Thr Tyr Gly  
1 5 10 15

Val Leu Ile Leu Leu Gly Pro Glu Asn Ala Tyr Ser Gly Ile Leu Leu  
20 25 30

Ser Ser Gly Lys Arg Ala Pro Phe Ser Pro Asn Leu Lys Asp His Glu  
 35 40 45  
 Asn His Leu Lys Cys Leu Leu Glu Val Arg Ile Pro Gln Pro Val Trp  
 50 55 60  
 Gly Pro Ala Ile Cys Ile Phe Lys Glu Thr Trp Thr Val Thr Cys Glu  
 65 70 75 80  
 Lys Pro Tyr Ala Gln Tyr Val Leu Ala Ile Arg Ile Thr Met Val Asn  
 85 90 95  
 Ile Asn Tyr Leu Phe Arg Glu His Lys Phe Leu Leu Thr Gln Leu Asn  
 100 105 110  
 Ala Lys Cys Phe Lys Ser Lys Thr Pro Cys Leu Lys Asn Ile Gly Phe  
 115 120 125  
 Phe Phe Lys Gln Tyr Lys Thr Gly Tyr Leu Ser His Glu Phe Gly Ala  
 130 135 140  
 Pro Asn Ser His Cys Phe Gln Thr Ile Ser Gln Glu Arg Ser Leu Gln  
 145 150 155 160  
 Ser Pro Pro Val Ala Ser Ile Ala Leu Cys Val Leu Lys  
 165 170  
 <210> 154  
 <211> 172  
 <212> PRT  
 <213> Homo sapiens  
 <400> 154  
 Gln Ile Leu Gly Ser Lys Arg Arg Lys Met Ser Arg Met Lys Arg Tyr  
 1 5 10 15  
 Leu Ile Ile Ser Ser Ala Asp Phe Leu Gly Asn Val Phe Ile Pro Ile  
 20 25 30  
 Phe Ile Thr Tyr Val Val Lys Asp Ser Phe Ser Gly Leu Tyr Ile Gln  
 35 40 45  
 Leu Phe Glu Tyr Ile Tyr Asn Asn Ile Tyr Ser Cys Leu Ile Gly Asn  
 50 55 60  
 Phe Asn Asn Tyr Gln Asn His Lys Glu Ile Phe Phe Ala Cys Phe His  
 65 70 75 80  
 Tyr Phe His His Phe Gly Ile Cys Tyr Val Val Lys Lys Tyr Ser Glu  
 85 90 95  
 Lys Thr Ile Ile Leu Lys Ser Cys Cys Ile Asn Arg Ile Trp Gly Lys  
 100 105 110  
 Glu Gln Thr Thr Lys Arg Gly Arg Leu Met Ser Leu Val Gly Thr Trp  
 115 120 125  
 Glu Val Thr Leu Ile Ser His Phe Leu Asn Leu Lys Glu Glu Lys Val  
 130 135 140  
 Lys Leu Ile Asn His Ser Thr Gln Lys Asn Thr Phe Trp Thr Ile Lys  
 145 150 155 160  
 Asp Ser Ala Ile Tyr Met Asp Tyr Ile Phe Ile Ser

165

170

<210> 155  
 <211> 231  
 <212> PRT  
 <213> Homo sapiens

<400> 155

Arg Cys Glu Pro Leu Pro Gly Leu Glu Leu Leu Leu Asp Cys Ile Pro  
 1 5 10 15  
 Arg Gly Asn Phe Met Thr Glu Phe Arg Ser Ala His Ile Leu Ala Ala  
 20 25 30  
 Ser Lys Arg Glu Arg Glu Ser Pro Ala Leu Ile Ser Val Ile Phe Leu  
 35 40 45  
 Phe Asp Leu Ile Tyr Ser Ile Asn Thr Pro Gln Glu Gly Thr Phe Pro  
 50 55 60  
 Ser Pro Ala Pro Lys Gln Asn Arg Ser Ile Leu Asp Gly Leu Pro Asn  
 65 70 75 80  
 Trp Cys Leu Gln Thr Ser Ser Leu Ser Pro Ser Pro Thr Leu Lys Ser  
 85 90 95  
 Arg Ser Leu Ile Cys Met Gly Cys Ile Ser Thr Leu Met Leu Pro Gly  
 100 105 110  
 Phe Trp Leu Gly Leu Pro Asn Gly Arg His His Trp Arg Arg Met Glu  
 115 120 125  
 Val Gly Gly Gly Arg Trp Glu Gly Arg Gly Trp Gly Ile Val Pro Leu  
 130 135 140  
 Ala Pro Phe Leu Cys Ser Phe Gly Ser Leu Gln His Pro Val Thr Leu  
 145 150 155 160  
 Ser Leu Ser His Gln Val Phe Ile Phe Cys Trp Phe Pro Phe Val Leu  
 165 170 175  
 Pro Thr Phe Thr Thr Cys Pro Phe Leu Lys Asp Pro Ser Ile Ala Leu  
 180 185 190  
 Phe Gly Asn Ile Leu Phe Ser Ala Gly Thr Pro Glu Leu Tyr Arg Arg  
 195 200 205  
 Val Gln Glu Ala Thr Lys Leu Gln Met Pro Thr Thr Trp Trp Asn Arg  
 210 215 220  
 Cys Pro Leu Glu Ala Ala Ala  
 225 230

<210> 156  
 <211> 160  
 <212> PRT  
 <213> Homo sapiens

<400> 156

Pro Ile Cys Leu Asn Ala Ser Cys Ser Gly Gly Leu Thr Pro Ile Asn  
 1 5 10 15  
 Pro Ser Cys Leu Trp Lys Gly Leu Pro Thr Glu Leu Asp Ser Asn Ile



20 25 30  
 Gln Ser Ser Ser Thr His Pro Phe Ser Trp Thr Leu Trp Gly Pro Arg  
 35 40 45  
 Gln Gln Thr Ser Cys Leu Phe Tyr Arg Ala Ala Leu Gln Met Ala Gly  
 50 55 60  
 Ala Thr Val Phe Ser Ala Leu Glu Asp Leu Ser Met Val Val Ser Phe  
 65 70 75 80  
 His Ile Ser Tyr Asp Phe Tyr Ser Gln Glu Ser Leu Ile Cys Leu Leu  
 85 90 95  
 Met His Phe His Leu Ser Val Thr Leu Leu Gln Asn Gln Arg Glu Ile  
 100 105 110  
 Thr Leu Ile Phe Leu Arg Ala Ser Lys Leu Pro Gly Leu Gln Arg Pro  
 115 120 125  
 Cys Arg Ala His Arg Gln Arg Met Thr Arg Gly His Met Pro Cys Met  
 130 135 140  
 His Phe His Leu Ser Val Thr Leu Leu Gln Ala Asn Leu Lys Gly Met  
 145 150 155 160  
 <210> 157  
 <211> 225  
 <212> PRT  
 <213> Homo sapiens  
 <400> 157  
 Val Pro Leu Val Asn Pro Glu Tyr Asn Ile Phe Tyr Lys Thr Cys Phe  
 1 5 10 15  
 Ile Leu Ser Gly Met Arg Cys Ile Phe Glu Gly Leu Leu Lys Leu Ala  
 20 25 30  
 Ile Thr Ile Arg Leu Leu Leu Asn Leu Gly Ile Ser Leu Pro Ser Cys  
 35 40 45  
 Gln Gly Leu Tyr Leu Met Phe Val Ser Leu Lys Lys Lys Arg Asn Gln  
 50 55 60  
 Thr Asp Tyr Thr Leu Leu Lys Thr Glu Asp Met Tyr Phe Asn Met Ser  
 65 70 75 80  
 Leu Leu Pro Val Ile Gln Ser Leu Lys Phe Gln Asn Pro Ser Gly Thr  
 85 90 95  
 Leu Cys Gly Pro Trp Ile Lys His Thr Trp Ala Tyr Glu Cys Val Asp  
 100 105 110  
 His Trp His Met Arg Gly Asn Cys Leu Leu Gly Tyr Val Ala Leu Pro  
 115 120 125  
 Leu Ser Ile Tyr Asn Ser Asn Val Ser Glu Arg Ser Ser Ser Leu Lys  
 130 135 140  
 Leu Phe Ser Arg Ile Arg Gln Thr Val Pro Ala Asn Gln Gly Asp Glu  
 145 150 155 160  
 Phe Trp Pro Met Phe Gly Arg Ser Leu Leu Gln Trp Gly Val Thr Ser  
 165 170 175

His Glu Arg Ile Ile Arg Asn Leu Ser Thr Thr Leu Gly Asn Leu Ala  
 180 185 190  
 Asn Glu Leu Ala Glu Ala Ile Ala Thr Lys Arg Ser Ser Asp Ser Leu  
 195 200 205  
 Asp Arg Ile Val Met Asp Asp Gly Ile Thr Leu Gly Tyr Ile Val Val  
 210 215 220

Lys  
 225

<210> 158  
 <211> 215  
 <212> PRT  
 <213> Homo sapiens

<400> 158

Leu Pro His Leu Cys Cys Ser Leu Leu Thr Ile Lys Pro Asp Met Cys  
 1 5 10 15  
 Leu Ser Pro Cys Leu Pro Thr His Pro Leu Ile Thr Ser Val Pro Cys  
 20 25 30  
 Ser Gln Val Ala Ser Arg Glu Asp Cys Gly Leu Met Ser Ser Phe Met  
 35 40 45  
 Pro Trp Leu Leu Leu Ile Arg Ala Leu Tyr Thr Phe Ser Lys Ala Leu  
 50 55 60  
 Glu Ser Lys Lys Val Leu Leu Gly Ser Ser Pro Gln Met Gln Phe Met  
 65 70 75 80  
 Lys Ser Val Ser Phe Ser Phe Pro Ser Glu Phe Leu Ser Val Ser Ile  
 85 90 95  
 Lys Ala Leu Asp Thr Pro Trp Phe Thr Arg Gln Lys Leu Ile His Pro  
 100 105 110  
 Thr Gln Pro His Gly Tyr Ser Phe Val Leu Leu Asp Asn Asn His Leu  
 115 120 125  
 Arg Lys Pro Asp Leu Phe Pro His Ser Ser Phe Ser Phe Cys Pro Ala  
 130 135 140  
 Glu Asn Lys Arg Thr Ser Cys His Ile Val Ile Cys Ser Ala Leu Leu  
 145 150 155 160  
 Leu Arg Ser Leu Val Gly Lys Thr Gly Pro Ile Lys Arg Asp Thr Ala  
 165 170 175  
 Met Pro Trp Gly Glu Asp Asn Lys Ser Asp Gly Ser Arg Ala Leu Glu  
 180 185 190  
 Ser Arg Gly Gly Val Thr Asn Cys Pro Asn Gly Thr Val Pro Ser Glu  
 195 200 205  
 Leu Leu His Leu Leu Leu Thr  
 210 215

<210> 159  
 <211> 202  
 <212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 159

Leu Lys Val Lys Lys Glu Tyr Pro Phe Ile Leu Asp Asn Cys Cys Gln  
 1 5 10 15  
 Arg His Tyr Asn Ile Ser Val Val Ile Pro Tyr Phe Ser Lys Ala Lys  
 20 25 30  
 Ile Glu Ile Trp Pro Leu Leu Leu Cys Asn Phe Leu Lys Phe Lys Val  
 35 40 45  
 Ser Val Phe Ser Ile Ile Lys Tyr Ser Ser Leu Lys Leu Met Ala Ile  
 50 55 60  
 Arg Tyr Ser Ile Val Trp Ile Ile Tyr Leu Arg Phe Cys Gly Leu Phe  
 65 70 75 80  
 Cys Phe Gln Asn Asn Thr Lys Ile Asn Ile Phe Val Cys Lys Tyr Phe  
 85 90 95  
 Thr Lys Ile Tyr Ser Glu Lys Phe Leu Lys Val Glu Phe Leu Gly Glu  
 100 105 110  
 Val Thr Phe Lys Cys Leu Ile His Leu Leu Ser Gly Lys Thr Val Arg  
 115 120 125  
 Phe Leu His Ser His His Ser Val Tyr Gly His Gln Leu Thr Val Phe  
 130 135 140  
 Phe Pro Thr Leu Leu Ile Phe Ser Leu Ser Met Trp Ile Lys Phe Gly  
 145 150 155 160  
 Phe Tyr Tyr Phe Asn Leu Tyr Ser Ile Thr Leu Leu Ala Ile Ser Leu  
 165 170 175  
 Gly Val Val Asn Ile Cys Pro Cys Pro Phe Leu Phe Gly Met Leu Ser  
 180 185 190  
 Leu Met Thr Asn Cys His Asn Val Ile Asn  
 195 200

&lt;210&gt; 160

&lt;211&gt; 215

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 160

Asn Ile Ser Phe Leu Ser Leu Lys Met Ala Val Ser Cys Val Leu Ile  
 1 5 10 15  
 Asn Leu Lys Ile Asn Leu Ser Ile Gly Glu Ala Gly Lys Leu Ala Trp  
 20 25 30  
 Lys Val Asn Leu Leu Ser Arg Gly Lys Ile Ser Trp Ala Leu Ile Lys  
 35 40 45  
 Val Asp Ile Phe Arg Gly Gly Lys Ser Lys Phe Tyr His Thr Leu Ala  
 50 55 60  
 Phe Val Gln Phe Ser Pro Leu Phe Ser Leu Tyr Tyr Leu Phe Phe Cys  
 65 70 75 80

Phe Thr Leu Gly Lys Ala Asn Tyr Leu Phe Ser His Ile Phe Trp Gly  
                     85                    90                    95  
 Pro Ile Leu Met Ile Leu Ile Phe Phe Ser Cys Leu Thr Cys Arg Pro  
                     100                    105                    110  
 Ser Thr Glu His Cys Arg Ala Ser Ser Gln Arg Ser Ser Gly Asp Glu  
                     115                    120                    125  
 Leu Ser Phe Leu Gly Trp Asp Cys Cys Ala Gly Leu Asp Arg Thr Glu  
                     130                    135                    140  
 Asn Cys Arg Asp Lys Tyr Thr Tyr Glu Gln Thr Ser His Leu Phe Ile  
                     145                    150                    155                    160  
 Lys Ala Leu His Trp Leu Trp Lys Thr Ala Val Gly Leu Arg Lys Leu  
                     165                    170                    175  
 Asn Phe Leu Gly Ile Phe Val Leu Asn Ile Glu Arg Glu Arg Arg Arg  
                     180                    185                    190  
 Phe Leu Phe Lys Arg Val Tyr Glu Thr Leu Ser Leu Lys Ser Asn Leu  
                     195                    200                    205  
 Met Thr Gly Cys Met Cys Ser  
                     210                    215

<210> 161  
 <211> 199  
 <212> PRT  
 <213> Homo sapiens

<400> 161

Lys Ile Gln Ile Leu Cys His Ser Pro Ala Tyr Leu Leu Thr Leu Pro  
 1                    5                    10                    15  
 Leu Leu Ser Lys Phe Ile Ile Leu Thr Val Val Val Asn Ala Leu Leu  
                     20                    25                    30  
 Ser Val Pro Cys Pro Phe Val Tyr Thr His Leu Val Leu Leu Ser Phe  
                     35                    40                    45  
 Phe Ile Asn Met Leu His His Thr Val Ile Phe Leu Leu Ile Phe Phe  
                     50                    55                    60  
 Lys Lys Val Trp Asn Ile Ser Phe Pro Leu Cys Val Leu Cys Asn Leu  
                     65                    70                    75                    80  
 Ser Asp Lys Thr Thr Cys Tyr Ile Phe Ser Thr His Asn Phe Ile Ser  
                     85                    90                    95  
 Gly Leu Cys Ala Leu Tyr Lys Ser Thr Asn Leu Ser Val Trp Ser Val  
                     100                    105                    110  
 Leu Ser Ser Pro Gly Gln Ile Leu Ile Ile Cys Gln Glu Cys Asn Ser  
                     115                    120                    125  
 Ile Ile Ser Ser Val Thr Gln Phe Ser Lys His Arg Ile Leu Cys Val  
                     130                    135                    140  
 Pro Ile Ala Leu His Trp Ile Gly Pro Gln Phe Cys Gln Cys Ile Ile  
                     145                    150                    155                    160  
 Arg Thr Tyr Leu Gln Val Leu Ser Leu Leu Leu Trp Arg Glu Pro Phe

165 170 175

Ser His Met Asn Cys Asp Phe Val Tyr Leu Ala Pro Thr Met Val Leu  
180 185 190

Asn Ser Trp Val Leu Gly Lys  
195

<210> 162  
<211> 213  
<212> PRT  
<213> Homo sapiens

<400> 162

Tyr Trp Phe Asn Lys Leu Trp Tyr Asn Gln Ile Met Lys Leu Tyr Ala  
1 5 10 15

Phe Val Lys Val Thr Phe Gln Lys Asn Ile Leu His Arg Ile Thr Asp  
20 25 30

Pro Ser Ala Leu Pro Thr Leu Trp Ala Leu Ser Leu Phe His His His  
35 40 45

Tyr Leu His His Cys Leu Gln Val Phe Tyr Thr Ala Arg Val Gly Leu  
50 55 60

Cys Leu Leu Asn Ser Gln Val Lys Arg Gly Arg Lys Leu Thr Pro Ser  
65 70 75 80

Gly Gly Ser Leu Gly Met Ile His Gly Arg Trp Ser Ile Asn Thr Ser  
85 90 95

Ala Leu Phe Pro Leu Glu Ile Leu Arg Asn Gly Phe Tyr Ile Val Ser  
100 105 110

Gln Ser Phe Leu Lys Val Leu Asn Phe Asn His Pro Gln Gly Trp Ala  
115 120 125

Leu Ser Tyr Thr Ser Phe Val Ala Ser Leu Pro Ser Cys Leu Thr Ser  
130 135 140

Pro Phe Gln Thr Arg Ile Tyr Phe Phe Ser Leu Lys Gln Asn Lys Met  
145 150 155 160

Phe Asn Leu Lys Pro Leu Gln Asn Thr Asn Leu Tyr Leu Lys Asn Leu  
165 170 175

Asn Ile Gly Glu Asn Glu Thr Val Tyr Ala Gln Val His Asp Trp Trp  
180 185 190

Arg Leu Lys Ser Ser Lys Ile Phe Leu Lys Gly Tyr Pro Ser Arg Arg  
195 200 205

Leu Asn Cys Leu Ile  
210

<210> 163  
<211> 236  
<212> PRT  
<213> Homo sapiens

<400> 163

Leu Ala Ser Glu Ser Leu Leu Val Arg Lys Glu Val Val Leu Phe Pro

1                    5                    10                    15  
 Leu Gln Ala Lys Ala Phe Gln Val Leu Ser Phe Cys Ser Ile Lys Arg  
                   20                    25                    30  
 Gln Leu Arg Gly Arg Tyr Pro Gln Glu Phe Pro Asp Ser Cys Thr Asp  
                   35                    40                    45  
 Leu Ser Ala Glu Ile Ala Glu Val Ser Trp His Leu His Glu His Leu  
                   50                    55                    60  
 Ser Val Ala Gly Arg Ile Asn Gly Lys Arg Ala Thr Glu Ile Pro Gly  
                   65                    70                    75                    80  
 Ala Lys Ser Ser Ser Glu Ser Pro Ile Phe Asp Gln Glu Leu Val Gly  
                   85                    90                    95  
 Ser Leu Arg Ile Cys Ile Ser Ser Asp Ser Arg Leu Ser Gly Leu Ser  
                   100                    105                    110  
 Asn Trp Asp Gln Ser Asn Ser Tyr His Ala Tyr Leu Val Pro Gly Ser  
                   115                    120                    125  
 Leu Leu Arg Ala Ser Trp Thr Pro Ala Arg Val Ser Pro His Ser Asn  
                   130                    135                    140  
 His Met Arg Tyr Val Leu Leu Leu Ser Pro Cys Ala Asp Glu Asp Thr  
                   145                    150                    155                    160  
 Arg His Arg Glu Asn Trp Pro Gln Val Tyr Ser Trp Gly Gly Gln Ser  
                   165                    170                    175  
 Gln Asn Ser Asp Leu Gly Cys Leu Gly Cys Glu Leu Val Trp Ala Ser  
                   180                    185                    190  
 Met Gly His Arg Gly Arg Ile Ser Trp Arg Ser Arg Thr Glu Gly Lys  
                   195                    200                    205  
 Arg Asp Glu Ile Ser Asp Ser Ala Gly Ser Glu Thr Leu Ser Ala Met  
                   210                    215                    220  
 Ile Lys Pro Asp Tyr Gly Thr Cys Phe Ser Leu Ser  
                   225                    230                    235  
  
 <210> 164  
 <211> 193  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 164  
 Phe Gln Asp Ile His His Arg Cys Gly Arg Gly Lys Lys Thr Met Gly  
                   1                    5                    10                    15  
 Met Gly Ile Leu Pro Phe Ile Asn Thr Gly His Phe Asn Leu Leu Asn  
                   20                    25                    30  
 Leu Ser Thr Phe Cys Asn Leu Arg Ile Phe Ile Leu Asp Ser Trp Thr  
                   35                    40                    45  
 Lys Ala Leu Glu Met Ala Ser Phe Ala Arg Phe Leu Cys Ala Leu Glu  
                   50                    55                    60  
 Lys Ile Pro Gly Phe Asn Ala Lys Asn Arg Gln Gln Arg Ala Gln Glu  
                   65                    70                    75                    80

Met Glu Leu Ser Gly Val Leu Leu Gln Leu Arg Thr Val Cys Tyr Ser  
85 90 95  
Pro Phe Lys Ile Ser Pro Asn Leu Tyr Leu Met Val Lys Asp Val Phe  
100 105 110  
Phe Phe Leu Leu Glu Glu Lys Val Thr Arg Ile His Gly Ser Gly Leu  
115 120 125  
Ile Val Leu Leu Leu Met Glu Ile His Lys Gln Phe Leu Lys Tyr Ser  
130 135 140  
Leu Ala Ser Glu Leu Val Trp Asn Leu Ala Val Tyr Leu Leu Asp Trp  
145 150 155 160  
Val Thr Thr Ala Val Ala Gly Ser Ile His Tyr Thr Arg Leu Cys Ile  
165 170 175  
Ser Met Met Ile Val Lys Phe Cys Glu Lys Val Leu His Leu Cys Ser  
180 185 190

Leu

<210> 165  
<211> 199  
<212> PRT  
<213> Homo sapiens

<400> 165

Leu Phe Ser Ala Phe Ser Leu Ile Leu His Leu Thr Gly Leu Val Val  
1 5 10 15  
Asn Ile Leu Lys Val Tyr Val Leu Ile Lys Thr Ser Ser Phe Pro Lys  
20 25 30  
Glu Lys Lys Ser Gln Phe Gly Leu Val Ser Leu Ser Cys Phe Leu His  
35 40 45  
Leu Thr Asn Val Ser Phe Ile Tyr Ser Phe Cys Ser Val Thr Phe Arg  
50 55 60  
Met Ile Leu Met Gly Lys Asn His Gly Ser Tyr Lys Gln Pro Phe Lys  
65 70 75 80  
Thr Ile Val Ile Leu Cys Ser Val Asp Ser Gly Arg Gly Phe Lys Val  
85 90 95  
Ile Ile Ser Leu Lys His Cys Val Asn Ile Pro Pro Thr Val Val Pro  
100 105 110  
Leu Gly Thr Gly Lys Ile Gln Asn Trp Pro Ala Ser Ser Leu Thr Arg  
115 120 125  
Val Ile Lys Val Arg Leu Leu Tyr Ile Lys Gln His Leu Asn Ala Trp  
130 135 140  
Cys Val Ala Ala Gly Lys Gln Pro Arg Ser Pro Ser Cys Ile Arg Gly  
145 150 155 160  
Leu Met Asn Val Ser Ile Ala Val Phe Ala Val Thr Arg Ser Gly Arg  
165 170 175

Val Phe Pro Ser Ser Leu Asp Cys Leu Pro Met His Thr Gly Val Cys  
 180 185 190

Ile Gly Lys Gln Ser Arg Leu  
 195

<210> 166  
 <211> 150  
 <212> PRT  
 <213> Homo sapiens

<400> 166

Ile Trp Cys Phe His Arg Leu Lys Gly Leu Arg Cys Pro Pro Val Ala  
 1 5 10 15

Val Ala Cys Gly Ser Leu Cys Ser Cys Leu Pro Ser Trp Ala Gln Tyr  
 20 25 30

Leu Val Leu Cys Leu Gly Phe Thr Asn Ala Thr Asn Thr Tyr Ala Pro  
 35 40 45

Thr Leu Cys Gln Val Leu Cys Tyr Met Leu Arg Lys Gln Cys Thr Arg  
 50 55 60

Trp Ile Arg Phe Ser Ser Leu Trp Cys Pro Ser Ser Gly Lys Asp Arg  
 65 70 75 80

Leu Ser Val Phe Tyr Gly Gln Ala Tyr Arg Ala Lys Lys Thr Cys Val  
 85 90 95

Gly Met Gly Gln Gly Arg Tyr Pro Trp Ser Ser Pro Val Thr Gly Ile  
 100 105 110

Arg Leu Arg Val Ile Val Gly Arg Ala Leu Gln Ala Gly Gly Ser Ala  
 115 120 125

Cys Ala Arg Val Leu Arg Lys Glu Gly Glu Gln Cys Val Arg Asn Ile  
 130 135 140

Thr Val Val Ala Thr Gln  
 145 150

<210> 167  
 <211> 218  
 <212> PRT  
 <213> Homo sapiens

<400> 167

Ile Ile Ile Arg Ile Ile Arg Ile Leu Lys Tyr Pro Asn Asn Gln Val  
 1 5 10 15

Asn Lys Ala Thr Phe Tyr Gly Ile Ile His Phe Cys Phe Glu Lys Tyr  
 20 25 30

Thr Leu Phe Lys Tyr Tyr Cys Leu Phe Thr Gln Leu Leu Glu His Ser  
 35 40 45

Ser Ala Lys Ala Phe Met Ile Phe Thr Asn Leu Ala Phe Ile Phe Ala  
 50 55 60

Leu Leu Ser Thr Ile Thr Lys Val Ile Thr Thr Cys Ser Pro Thr Asn  
 65 70 75 80



Tyr Ser Asp Gly Ala Leu Arg Ile Asp Leu Tyr Leu Asn Ile Leu Trp  
                     85                    90                    95  
 Tyr Gln Val Phe Leu His Ser Ser Arg Ile Phe His Phe Ala Tyr Ile  
                     100                    105                    110  
 Leu Met Met Ser Ser Arg Ile Ser Ser Leu Thr Tyr Leu Ala Asn Tyr  
                     115                    120                    125  
 Lys Tyr Val Ile Phe Val Lys Tyr Leu Arg Val Cys Ser Ala Ile Tyr  
                     130                    135                    140  
 Leu Val Ile Leu Asn Gln Ile Leu Asn Val Tyr Thr Phe Leu Met Tyr  
                     145                    150                    155                    160  
 Asn Phe Gln Phe Phe Arg Met Arg Leu Asn Asn Cys Pro Tyr Tyr Ser  
                     165                    170                    175  
 Phe Ile Thr Thr Leu Ile Tyr Leu Leu Tyr Leu Gln Met Ile Tyr Lys  
                     180                    185                    190  
 Asn Ala Phe Leu Tyr Leu Ser Leu Ser Gln Val Leu His Ser Glu Leu  
                     195                    200                    205  
 Phe Phe Leu Phe Val Phe Leu Arg Tyr Ile  
                     210                    215  
 <210> 168  
 <211> 204  
 <212> PRT  
 <213> Homo sapiens  
 <400> 168  
 Tyr Cys Glu Leu Arg Cys Tyr Ile Ser Glu Cys Asn Glu Trp Asp Ile  
   1                    5                    10                    15  
 Ala His Trp Leu Glu Lys Pro Pro Lys Gln Ala Ala Ser Ala Ile Glu  
                     20                    25                    30  
 Leu Leu Ala Trp Ser Arg His Ser Ala Ser Gly His Gly Asp Asn Ser  
                     35                    40                    45  
 Ser Glu Ile Asn Ser Ser Thr Lys Val Ser Asn Asp Val Ile Ser Ser  
                     50                    55                    60  
 Gln Arg Gln Gly Cys Pro Val Lys Gln Thr Asp Gly Gln Ser Pro Pro  
   65                    70                    75                    80  
 Arg Leu Lys Gly Gly Gly Glu Thr Gly Arg Lys Arg Met Arg Trp Val  
                     85                    90                    95  
 Arg Lys Arg Tyr Asn Leu Arg Val Thr Met Ser Ser Cys Ser Pro Arg  
                     100                    105                    110  
 Trp Gln Trp Val Gly Gly Pro Gly Lys Asp Cys Phe Arg Gln Met Glu  
                     115                    120                    125  
 Gln Cys Met Arg Arg Ser Arg Glu Lys Ser Gln Ile Val Cys Ile His  
                     130                    135                    140  
 Val Leu Gln Asn Arg Glu Ser Asn Arg Tyr Leu Gly Lys Lys Lys Glu  
   145                    150                    155                    160  
 Val Ser Leu Phe Leu Ser Leu Lys Val Gln Lys Trp Ala Phe Pro Gln

Phe Ile Cys Gln Pro His Glu Val Phe Thr Asp Leu Asp Leu Leu Ile  
180 185 190

Ser Cys Tyr Phe Ile Thr Leu Leu Glu Leu Leu Pro  
195 200

```
<210> 169
<211> 158
<212> PRT
<213> Homo sapiens
```

<400> 169

Lys Val Leu Ile Phe Val Leu Arg Pro Ile Tyr Thr Tyr Lys Cys His  
1                   5                   10                   15

Pro Ser Ile Phe Leu Cys Asn Phe Leu Ser Ala Gly Leu Pro Ser Leu  
20 25 30

Met Cys Val Leu Tyr Phe Pro Tyr Ile Cys Tyr Pro Ile Thr Cys Phe  
35 40 45

Tyr Asn Cys Leu Phe Tyr Phe Pro Phe Phe Ser His Cys Leu His Ala  
50 55 60

Leu-Phe Leu-Val Leu Asn Ser-Ile Thr-Leu Ile-His Cys Ser Ser Asn  
65 70 75 80

Phe Ile Leu Asn Asn Phe Pro Ile Tyr Leu Asp Ile Tyr Leu Asn Val  
85 90 95

His Ile Ser Pro Leu Ile Glu Val Cys Leu Val Ile Phe Gly Met Met  
100 105 110

Leu Asn Leu Phe Leu Trp Lys Gly Thr Asn Thr Cys Met Phe Met His  
115 120 125

Val Gln Lys Cys Ser His Arg Met Ile Ile Lys Ala Asp Leu Gly Lys  
130 135 140

Lys Thr Ser Leu Ile Phe Ile Phe His Ile Arg Phe Phe Glu  
145 150 155

```
<210> 170
<211> 198
<212> PRT
<213> Homo sapiens
```

<400> 170

His Gln Asn Ser Pro Ile Tyr Leu Arg Ile Asn Val Asn Phe Glu Phe  
1 5 10 15

Asp Ile Thr Met Ile Lys Gly Ala Leu Ile Phe Ser Arg Ser Tyr Lys  
20 25 30

Ile Phe Val Asn Glu Leu Ile Gly Arg Ile Cys Leu Leu Lys Ser Glu  
35 40 45

Val Gly Gly Glu Leu Lys Leu Gly Leu Ile Gly Asn Tyr Ile Trp Val  
50 55 60

Met Asn Ala Trp Gly Phe Ile Ile Pro Leu Pro Leu Pro Leu Ser Val

65                      70                      75                      80  
 Phe Glu Leu Cys His Cys Glu Asn Ile Val Leu Lys Ala Val Leu Phe  
                                  85                                   90                                   95  
 Phe Leu Leu Arg Gly Ser Lys Lys Ser Lys Lys Tyr Thr Gly Leu Ile  
                                  100                                   105                                   110  
 Glu Tyr Val Cys Ser Asn Lys Ile Pro Gly Phe Ser Phe Val Leu Ala  
                                  115                                   120                                   125  
 Ser Arg Asn Gln Val Gln Phe Val Ser Lys Asp Phe Ala Thr Cys Gly  
                                  130                                   135                                   140  
 Gly Lys Leu Leu Gln Asp Leu Ile Val His Ser Gln Arg Leu Ser Ala  
                                  145                                   150                                   155                                   160  
 Ala Arg Gln Ala Ala Phe Tyr Glu Asn Asp Asn Gln Lys Ala Gly Ala  
                                  165                                   170                                   175  
 Leu His Thr Gly His Ser Ser Asn Glu Ser Trp Asp Leu Asp His Gly  
                                  180                                   185                                   190  
 Ser Leu Thr Trp Ala Ala  
                                  195

<210> 171  
 <211> 176  
 <212> PRT  
 <213> Homo sapiens

<400> 171

Leu Lys Val His Val Leu Ile Tyr Ile His Gln Ile Thr Thr Thr Ser  
 1                                   5                                   10                                   15  
 Ser Phe Leu Phe Ile Ser Leu Leu Pro Phe Ile Ser Phe Ile His Met  
                                  20                                   25                                   30  
 Leu Ser Leu Asn Thr Leu Leu Leu Leu Thr Val Ile Phe Gln Ile  
                                  35                                   40                                   45  
 Ser Glu Lys Asn Leu Ile Leu Pro Tyr Ser Thr Phe Leu Met Leu Phe  
                                  50                                   55                                   60  
 Leu Phe Tyr Ala Val Leu Phe Asp Ile Ser His Arg Ala Gly Gln Leu  
 65                                   70                                   75                                   80  
 Ala Met Asn Tyr Ser Ser Phe Val Cys Gln Lys Ile Ser Leu Phe Leu  
                                  85                                   90                                   95  
 Ile Arg Ile Ile Leu Leu Asn Ala Glu Phe Gly Ser Phe Phe Val Ala  
                                  100                                   105                                   110  
 Thr Leu His Val Phe Ser Phe Leu Cys Val Cys Met Val Ser Glu Glu  
                                  115                                   120                                   125  
 Lys Asp Asn Val Ile Leu Ile Leu Phe Pro Leu Trp Ile Arg Cys Trp  
                                  130                                   135                                   140  
 Leu Phe Pro Leu Ser Ser Phe Phe Gln Asp Phe Leu Phe Ser Leu Val  
 145                                   150                                   155                                   160  
 Phe Cys Ser Leu Asn Met Ile Cys Leu Gly Gly Asp Leu Asp Leu Leu  
                                  165                                   170                                   175

<210> 172  
 <211> 195  
 <212> PRT  
 <213> Homo sapiens

<400> 172

Ala Tyr Arg Ile Ser Thr Thr Val Phe Ala Lys Glu Lys Ser Val Val  
 1 5 10 15  
 Ile Lys Phe Ile Leu Trp Leu Asn Tyr Val Leu Gln Phe Val Gly Pro  
 20 25 30  
 Val Thr Cys Gly Arg Gln Arg Ala Val Gly His Ser Val Lys Ala Thr  
 35 40 45  
 Thr Arg Val Leu Ser Ile Glu Ser Leu Cys Ile Met Val Leu Ala Arg  
 50 55 60  
 His Cys Ser Leu Thr Ser Ile Phe Leu Ser Gln Ser Ser Leu Arg Asn  
 65 70 75 80  
 Ala Cys Ser Thr Gly Leu Ile Ile Leu Thr Glu Thr Ser Gly His Phe  
 85 90 95  
 Met Ser Tyr Gly Met Leu Ala Glu Asp Ile Lys His Arg Cys Val Gly  
 100 105 110  
 Ile Gly Gly Glu Ser Thr Ala Ile Phe Gln Leu Gly Ala Pro Trp Phe  
 115 120 125  
 Pro Glu Ile Gln Ser His Gly Val Asn Gln Thr Pro Leu Ser Gly Ala  
 130 135 140  
 Leu Cys Ser Thr Gln Asp Pro Thr Leu Ser Gly Lys Leu Lys Thr Lys  
 145 150 155 160  
 Ser Leu Leu Tyr Ile Arg Phe Ile Lys Asn Ala Thr Ile Thr Lys Ser  
 165 170 175  
 Leu Trp Ala Cys Val Glu Asn Ala Val Ile Lys Leu Asn Ile Lys Ala  
 180 185 190  
 Ser Ser Lys  
 195

<210> 173  
 <211> 225  
 <212> PRT  
 <213> Homo sapiens

<400> 173

Gln Arg Leu Thr Tyr Ser Asn Cys Ile Val Asp Trp Ala His Thr Leu  
 1 5 10 15  
 His Val Thr Asn Val Ser Asn Tyr Trp Ile Cys Thr Ala Leu Pro Ala  
 20 25 30  
 Gly Leu Arg Met Ala Cys Leu Gly Thr Tyr Ile Leu Cys Leu Gln Arg  
 35 40 45  
 Thr Gly His Gly Trp Arg Leu Gly Gly Pro Met Ala Asp Ala Trp Asn  
 50 55 60

Ala Thr Trp Gln Leu Trp Thr Lys Asp Ala Ala Arg His Met Val Cys  
65 70 75 80  
Pro Thr Pro Gly Trp Pro Ile Ala Phe Met Met Gly Leu Ala Ser Gly  
85 90 95  
Glu His Val Val Leu Pro Ala Gln Val Pro Gln Cys Ile Glu Gln His  
100 105 110  
Trp Gly Asn Thr Thr Val Gly Trp Val Pro Val Thr Ala Phe Ala Asn  
115 120 125  
Ile Thr His Val Thr Thr Lys Val Arg Pro Leu Thr Leu Cys Pro Leu  
130 135 140  
Gly Val Tyr Gly Ser Val Gly Thr Gln Ser Arg Phe Thr Tyr Pro Thr  
145 150 155 160  
Ala Leu Asp Ile Val Pro Gly Gly Gly Leu Met Cys Leu Pro Leu Phe  
165 170 175  
Ser Pro Cys Cys Pro Asp Ala Arg Ile Thr Gly Arg Cys Tyr Thr Leu  
180 185 190  
Ser Leu Cys Glu Cys Asn Glu Pro Pro Ala Val Leu Pro Phe Gly Ser  
195 200 205  
Asp Tyr Pro Trp Ser Gly Cys His Asn Cys Arg Ser Thr Gly Tyr Cys  
210 215 220

Ser  
225

<210> 174  
<211> 169  
<212> PRT  
<213> Homo sapiens

<400> 174

Phe Met Ile Gln Gln Ile Lys Cys Gly Asn Tyr Leu Lys Arg Lys Lys  
1 5 10 15  
Lys Asn Ile Trp Glu Ala Ala Glu Met Arg Thr Ile Arg Asn Glu His  
20 25 30  
Phe Tyr Phe Leu Ser Phe Leu Asn Gly Ala Ser Asp Ala Val Phe Ile  
35 40 45  
Ala Leu Phe Phe Pro Asn Trp Asn Ile Phe Phe Leu Ile Leu Leu Val  
50 55 60  
Tyr Ser Leu Val Thr Lys Lys Val Phe Arg Lys Tyr His Asn Phe Pro  
65 70 75 80  
Asn Ser Leu Leu Ser Ala Gly Asp Tyr Glu Tyr Ile Leu Gln Asn Gly  
85 90 95  
Lys Gly Gly Ser Ser Gly Pro Ala Thr Ile Cys Ile Leu Lys Asp Leu  
100 105 110  
Val Glu Leu Lys Ser Gln Arg Lys Trp Glu Glu Leu Ser Lys Tyr Phe  
115 120 125

Ile Ile Phe Phe Leu Glu Tyr Gln Val Leu Ile His His Ile Phe His  
 130 135 140

His Val Ser Lys Ser Phe Phe Leu Lys Lys Val Cys Ile Tyr Ile Ser  
 145 150 155 160

Lys Arg Val Ser Val Val Lys Lys Asn  
 165

<210> 175  
 <211> 199  
 <212> PRT  
 <213> Homo sapiens

<400> 175

Glu Asn Thr Tyr Gly Lys Glu Leu Ser Val Arg Phe Gly Ser Gln Ile  
 1 5 10 15

Leu Ile Phe Asn Lys Ile Tyr Ile Cys Ser Pro Cys Thr Lys Gly Asn  
 20 25 30

Ser Thr Glu Ser Met Pro Asn Ser Lys Gly Met Thr Leu Asn Leu Tyr  
 35 40 45

Ser Lys Tyr Ile Gly Pro Ala Ile Leu Cys Gln Met Leu Tyr Leu Tyr  
 50 55 60

Leu Ile Ala Thr Arg Thr Gly Asn Cys Ala Gln Leu His Leu Arg Thr  
 65 70 75 80

Val Ser Ile Leu Lys His Thr Ser Tyr Ser Ser Ser Asp Pro His Trp  
 85 90 95

Met Lys Leu Asn Gln Thr Lys Gln Lys Ser Tyr Leu Ser Pro Asn Asn  
 100 105 110

Glu Arg Val Cys Arg Met His Ile Val Arg Leu Thr Asp Pro Phe Arg  
 115 120 125

Gln Tyr Val Gly Phe Pro Arg Ile Leu Ser Ala Ser Lys Gln Phe Glu  
 130 135 140

Phe Ser Ser Ala Leu Met Ile Trp Phe Pro His Leu Asp Gly Pro Gly  
 145 150 155 160

Ser Asp Ala Arg Gly Pro His Glu Met Ser Trp Ala Phe Ile Gln Asp  
 165 170 175

Pro Val Ala Pro Ala Gln Glu Asn Arg Pro Leu Arg Val Ser Gly Ser  
 180 185 190

Glu Met Ala Ser Val Thr Arg  
 195

<210> 176  
 <211> 204  
 <212> PRT  
 <213> Homo sapiens

<400> 176

Leu Phe Asn Phe Val Phe Val Ala Val Val Cys Ile His Val Cys Trp  
 1 5 10 15

Cys Pro Tyr Val Leu Phe Gly Val Trp Leu Phe Ser Gln Asn Gln Val  
 20 25 30  
 Thr Val Lys Ser Leu Asn Phe Ser Ile Ser Leu Leu Ser Ser Gly Thr  
 35 40 45  
 Val Thr Val Cys Leu Leu Leu Lys Ser Phe Val Phe Leu Thr Arg Gly  
 50 55 60  
 Glu Val Tyr Ser Thr Leu Thr Gly Leu Tyr Phe Gly Leu Arg Pro Tyr  
 65 70 75 80  
 Lys Thr Phe Leu Lys Ser Leu Ile Ile Cys His Ile Ile Lys Lys Leu  
 85 90 95  
 Tyr Gly Ile Phe Ser His Tyr Ile Leu Ala Thr Met Pro Val Tyr Ile  
 100 105 110  
 Ser Lys Gln Thr Ile Cys Gly Asn Asn Leu Lys Lys Lys Ala Ile Gly  
 115 120 125  
 Ser Lys Tyr Leu Ile Lys Tyr Pro Leu Glu Leu Asn Ile Ser Ser Cys  
 130 135 140  
 Gly Ser Ser His Thr Lys Tyr Pro Thr Leu Leu Ser Phe Arg Val Leu  
 145 150 155 160  
 Ala Gly Thr Gly Ser Ile Lys Asp Asn Glu Leu Lys Lys Gly Thr Ile  
 165 170 175  
 Tyr Lys Tyr Val Ala Arg Leu Gly Glu Thr Ser Lys Val Gly Asn Ala  
 180 185 190  
 Ala Gln Asp Ser Asn Lys Ser Glu Asn Leu Phe Leu  
 195 200  
 <210> 177  
 <211> 201  
 <212> PRT  
 <213> Homo sapiens

<400> 177

His Val Thr Leu Met Ser Thr Val Phe Ser Ser Val Ala Ser Thr Pro  
 1 5 10 15  
 Leu Pro Asn Ser Tyr Asp Asn Ser Ala Ser Gln Thr Tyr Gly Leu Arg  
 20 25 30  
 Asn Pro Leu Lys Ser Gln Leu Val Met Thr Pro Lys Arg Phe Phe Ile  
 35 40 45  
 Ile Ile Leu Tyr Ile Asn Ile Leu Leu Glu Val His Phe Tyr Glu Asn  
 50 55 60  
 Asn Leu Phe Ser Lys Ile Ser Glu Lys Asn Ser Ile Ile Leu His Ile  
 65 70 75 80  
 Gly Ile Phe Leu Met Pro Gly Leu Ile Glu Asp Asn Ile Phe Met Ser  
 85 90 95  
 Thr Ser Gly Phe Asp Leu Phe Gln Tyr Val Ser Leu Val Glu Ile His  
 100 105 110  
 Glu Gly Asn Leu Gly Ser Ser Asp Ile Leu Glu Lys Gly Gly Val Phe

115                      120                      125  
 Gln Pro Phe Trp Thr Thr Val Asp Ile Val Leu Tyr Tyr Asn Lys Thr  
 130                      135                      140  
 Gly Glu Val Val Gly Ser Lys Leu Val Ala Thr Trp Asn Leu Lys Pro  
 145                      150                      155                      160  
 His His Glu Leu Phe Val Ile Trp His Ile Lys Ile Tyr Leu Ser Ile  
 165                      170                      175  
 Leu His Phe Glu Trp Asp Pro Leu Leu Met His Leu Phe Val Thr Ile  
 180                      185                      190  
 Ile Ser Asn Thr Leu Val His Val Met  
 195                      200  
 <210> 178  
 <211> 216  
 <212> PRT  
 <213> Homo sapiens  
 <400> 178  
 Ile Lys Ile Pro Ala Val Lys Leu Asp Ser Ala Cys Leu Gly Ile Phe  
 1                      5                      10                      15  
 Lys Arg Ile Met Tyr Arg Gly Cys His Gly Asn Ser Ser Ser Gly Asn  
 20                      25                      30  
 Ser Val Pro Phe Val Lys Thr Leu Lys Gly Glu Asp Lys Gln Phe Gly  
 35                      40                      45  
 Glu Ile Thr Ala Pro Glu Ile Glu Phe Ile Cys Asn Leu Gly Ser Leu  
 50                      55                      60  
 Val Cys Leu Pro Ala Ile His His Val Asp Glu Lys Gln Lys Asp Lys  
 65                      70                      75                      80  
 Lys Asp Ser His Phe Lys Ala Pro Asn Cys Gln Phe His Ser Ile Ala  
 85                      90                      95  
 Asp Ser Gln His Arg Arg Lys Trp Asp Asn Ala Gly Arg His Tyr His  
 100                      105                      110  
 Arg Thr Val Ser Ser Lys Glu Lys Pro Asn Cys Tyr Phe Ser Met Ala  
 115                      120                      125  
 Glu Gly Gly Cys Phe Pro Arg Gly Arg Ile Leu Phe Asn Pro Val Arg  
 130                      135                      140  
 Ala Gln Leu Gln Pro Ser Val Thr Gly Gln Leu Pro Pro Ser Asn Pro  
 145                      150                      155                      160  
 Glu Gly Arg His Glu Pro Tyr Ser Arg Thr Gly Ala Cys Ser Leu Leu  
 165                      170                      175  
 Ser Thr Ser Cys Thr Phe Arg Ala Pro Ala Trp Asp Ala Glu Asn Ser  
 180                      185                      190  
 His Pro Ser Arg Ala Ala Glu Asp His Met Thr Asp His Gln Leu Phe  
 195                      200                      205  
 Leu Thr His Leu Ser Thr Thr Thr  
 210                      215



<210> 179  
 <211> 189  
 <212> PRT  
 <213> Homo sapiens

<400> 179

Ser Gln Asn Phe Asp Leu Thr Asn Gln Arg Gly Gly Leu Val Phe Phe  
 1 5 10 15  
 Tyr Leu Leu Ser Ala Phe Cys Phe Arg Leu Leu Asn Leu Tyr Ile Lys  
 20 25 30  
 Thr Cys Tyr Thr His Leu Ala Val Phe Phe Phe Ala Ala Val Thr Ser  
 35 40 45  
 Phe Trp Leu Arg Phe Phe Phe Lys Lys Met Tyr Lys Thr Leu Gly Leu  
 50 55 60  
 Ile His Cys Ser Phe Phe Val Leu Ile His Pro Gln Glu Arg Lys Trp  
 65 70 75 80  
 Leu Ser Leu Tyr Val Phe Lys Gly Leu Cys Glu Leu Leu Lys Ala Ser  
 85 90 95  
 Val Thr Ala Arg Thr Ser Val His Lys Gln Val Gln Asp Ala Ala Glu  
 100 105 110  
 Gly Val Ser Ser Leu Thr Glu Arg Gly Ile Glu Leu Phe Arg Met Phe  
 115 120 125  
 Cys Val Gly Thr Asp Arg Leu Lys Ala Thr Asp Leu Met Glu Val Trp  
 130 135 140  
 Ser Phe Gln Gln Met Ser Ser Asn Leu Thr Asn Leu Asp Leu Val Phe  
 145 150 155 160  
 Pro His Gly Pro Arg Ser Ala Ile Leu Phe Phe Cys Leu His Leu Ile  
 165 170 175  
 Ser Tyr Ala His His Cys Ala Asn Ser Arg Leu Phe Ser  
 180 185

<210> 180  
 <211> 157  
 <212> PRT  
 <213> Homo sapiens

<400> 180

Val Ala Ile Cys Gln Val Pro Thr Asp Ile Pro Asn Ile Arg Leu Thr  
 1 5 10 15  
 Pro Ser Asn Gln His Pro Glu Phe Lys Val Cys Ile His Phe Leu Tyr  
 20 25 30  
 Phe Tyr Cys Ile Arg Ile Ser Leu Asn Ser Ser Val Phe Ser Thr Phe  
 35 40 45  
 Ile Tyr Gln Pro Tyr Leu Pro Phe Cys Asn Leu Leu Phe Ser Val Ser  
 50 55 60  
 Ile Ile Phe Met Arg Leu Met His Ile Ala Val Tyr Ser Phe Leu Leu  
 65 70 75 80

Leu Tyr Asn Ser Val Ile Pro Gly Met Gly Arg Gly Asn Trp Phe Gln  
                     85                    90                    95  
 Asp Leu Cys Gly Leu Gln Asn Pro Ser Met Phe Lys Ser Leu Ile Asn  
                     100                    105                    110  
 Glu Ala Val Leu Ala Tyr Asn Leu Cys Thr Phe Leu Arg Thr Leu Ser  
                     115                    120                    125  
 Lys Cys Tyr Val Asn Gly Cys Phe Val Ile Cys Ile Ile Phe Ile Val  
                     130                    135                    140  
 Met Phe Phe Leu Leu Phe Ser Pro Glu Phe Phe Phe Phe  
                     145                    150                    155

<210> 181  
 <211> 219  
 <212> PRT  
 <213> Homo sapiens

<400> 181

Val Thr Leu Val Cys Tyr Ser Leu Met Val Arg Ser Leu Ile Lys Pro  
 1                    5                    10                    15  
 Glu Glu Asn Leu Met Arg Thr Gly Asn Thr Ala Arg Ala Arg Ser Ile  
                     20                    25                    30  
 Arg Thr Ile Leu Leu Val Cys Gly Leu Phe Thr Leu Cys Phe Val Pro  
                     35                    40                    45  
 Phe His Ile Thr Arg Ser Phe Tyr Leu Thr Ile Cys Phe Leu Leu Ser  
                     50                    55                    60  
 Gln Asp Cys Gln Leu Leu Met Ala Ala Ser Val Ala Tyr Lys Ile Trp  
 65                    70                    75                    80  
 Arg Pro Leu Val Ser Val Ser Ser Cys Leu Asn Pro Val Leu Tyr Phe  
                     85                    90                    95  
 Leu Ser Arg Gly Ala Lys Ile Glu Ser Gly Ser Ser Arg Asn Gly Arg  
                     100                    105                    110  
 Thr Ser Trp Val Ser Ile Gln Leu Gly Gly Arg Asp Ala Gln Gly Thr  
                     115                    120                    125  
 Asp Leu Gly Asn Ala Lys Val Lys Leu Gly Lys Asn Glu Leu Gln His  
                     130                    135                    140  
 His Gln Gln Leu Val Cys Thr Gln Met Ser Ala Gly Gly Arg Gly Ala  
 145                    150                    155                    160  
 Gln Asp Leu Leu Lys Val Ser Cys Cys Lys Gly His Phe Tyr Ile Asp  
                     165                    170                    175  
 Val Lys Val Asn Lys Ser Met Glu Arg Ala Thr Lys Thr Lys Glu Asn  
                     180                    185                    190  
 Phe Leu Lys Glu Ser His Trp Ser Leu Val Ile Gln Val Ser Ala Gln  
                     195                    200                    205  
 Met Ser Pro Leu Arg Asp His Ser Cys Pro Pro  
                     210                    215

<210> 182  
 <211> 181  
 <212> PRT  
 <213> Homo sapiens

<400> 182

Gln Gly Glu Gly Gly Thr Gly Tyr Lys Arg Ser Ala Ala Ala Ala Pro  
 1 5 10 15  
 Ala Glu Ser Arg Arg Ala Gln His Ser Cys Pro Leu Asp Pro Ala Asp  
 20 25 30  
 Pro Ser Arg Ala Pro Ser Val Pro Gln Ala Gln Pro Pro Gly Gly Arg  
 35 40 45  
 Ala Glu Gly Ser Pro Gly Arg Cys Gln Gly Ala Ile Leu Glu Gly Gly  
 50 55 60  
 Arg Glu Glu Glu Val Arg Ala Ala Met His Thr Val Ala Thr Ser Gly  
 65 70 75 80  
 Pro Asn Ala Ser Trp Gly Ala Pro Ala Asn Ala Ser Gly Cys Pro Gly  
 85 90 95  
 Cys Gly Ala Asn Ala Ser Asp Gly Pro Val Pro Ser Pro Arg Ala Val  
 100 105 110  
 Asp Ala Trp Leu Val Pro Leu Phe Phe Ala Ala Leu Met Leu Leu Gly  
 115 120 125  
 Leu Val Gly Asn Ser Leu Val Ile Tyr Val Ile Cys Arg His Lys Pro  
 130 135 140  
 Met Arg Thr Val Thr Asn Phe Tyr Ile Gly Glu Cys Gly Pro Leu Arg  
 145 150 155 160  
 Arg Thr Cys Cys Arg Pro Gly Gly Leu Arg Gly Pro Ser Gly Leu Gly  
 165 170 175  
 Arg Pro Leu Ala Thr  
 180

<210> 183  
 <211> 227  
 <212> PRT  
 <213> Homo sapiens

<400> 183

Ile Ile Leu Gln Asp Asn Leu Lys Gln Tyr Leu Val His Ile Asn His  
 1 5 10 15  
 Phe Ile Ser Ala Gly Leu Leu Ser Phe Glu Asn Tyr Phe Tyr His Leu  
 20 25 30  
 Leu Leu Ala Thr Val Asn Leu Ser Asn Leu Val Ser His His Ser Leu  
 35 40 45  
 Ile Pro Cys Ser Ala Leu Val Thr Met Asn Leu Ser Leu Leu Leu Lys  
 50 55 60  
 Tyr Ala Ile Tyr His Val Phe Phe Phe Pro Phe Ser Leu Pro Glu Ala  
 65 70 75 80

His Thr Pro Ser Leu Gly Trp Leu Lys Ser His Asn Leu Thr Phe Gly  
                     85                    90                    95  
 Leu Thr Phe Tyr Asn Ser Leu Tyr Gln Pro Gln Asn Met Ala Trp Val  
                     100                    105                    110  
 Met Leu Ala Leu Thr Val Leu Asp Phe Ser Asp Pro Ser Leu Leu Ile  
                     115                    120                    125  
 Tyr Gln Pro Leu Ser Arg Ser Phe Gly Thr Tyr Ser Asp Phe His Thr  
                     130                    135                    140  
 Pro Glu Leu Phe Ala Ile Leu Phe Ile Trp Lys Ser Tyr Trp Val Ile  
                     145                    150                    155                    160  
 Phe Leu Phe Lys Tyr Asn Leu Ile Ile Thr Pro Leu Val Tyr Leu Ala  
                     165                    170                    175  
 Leu Ser Cys Ser Leu Tyr Phe Pro Cys Pro His Leu Asn Ser Leu Thr  
                     180                    185                    190  
 Gly Glu Ile Asn Tyr Arg Tyr Thr Lys Gly Pro Asp Ser Lys Arg Asn  
                     195                    200                    205  
 Ile Gly Lys Ile Ser Ser Pro Ser Gln Pro Gly Tyr Gln Ile Lys Asp  
                     210                    215                    220  
 Arg Arg Leu  
 225

<210> 184  
 <211> 191  
 <212> PRT  
 <213> Homo sapiens

<400> 184

Pro Pro Thr Asp Ile Ser Val Cys Cys Ser Asp Gln Val Leu Gly His  
 1                    5                    10                    15  
 His Gln Cys Pro Val Val Met Gly His Leu Lys Leu Tyr Leu Tyr Pro  
                     20                    25                    30  
 Ser Ala Leu Leu Leu Asp Leu Leu His His Leu Leu His Met Asp Leu  
                     35                    40                    45  
 Leu His Phe Gly Cys Val Val His His Leu His Thr Leu Pro Asn Lys  
                     50                    55                    60  
 Asn Ile Gln Lys Pro Ser Ser Gln His His Cys Pro Gly His His Ser  
 65                    70                    75                    80  
 Ser Leu Phe Phe Leu Asn Pro Ser Leu His Glu Arg Gln Arg Arg Leu  
                     85                    90                    95  
 Thr Gly Ser Pro Leu Leu Val Asn His Met Lys Ile Lys His Ala Tyr  
                     100                    105                    110  
 Ser Val Leu Val Gln Gln Glu Ile Tyr Phe Gln Thr Arg Lys Ala Thr  
                     115                    120                    125  
 Glu Thr Leu Gly Ile Ile Leu Gly Ala Phe Ile Ile Cys Trp Leu Pro  
                     130                    135                    140  
 Leu Phe Ile Val Ser Leu Pro Ala Lys Ile Pro Pro Tyr Asp Ile Phe

145                      150                      155                      160  
 Ile Leu Leu Ser Phe Phe Phe Phe Phe Leu Ile Pro Ser Leu Thr  
                          165                                      170                                      175  
 Leu Val Ser Gln Ala Arg Met Gln Trp Tyr Asn Leu Ser Ser Leu  
                          180                                      185                                      190

<210> 185  
 <211> 76  
 <212> PRT  
 <213> Homo sapiens

<400> 185

Ile Leu Pro Ala His Leu Ile Pro Leu Gly Lys Leu Trp Cys Cys Leu  
 1                                      5                                      10                                      15  
 Ser Arg Thr Glu Ala Glu Gly Trp Leu Ser Pro Thr Gly Ser Tyr Ser  
                          20                                      25                                      30  
 Leu Asn Ser Ala Ser Ser Pro Arg Leu Gly Glu Thr Thr Trp Gly His  
                          35                                      40                                      45  
 Arg Val Phe Ala Arg Cys His Phe Ala Phe Gln Thr Arg Ser Trp Ser  
                          50                                      55                                      60  
 Ser Gly Phe Arg Leu Gly Leu Trp Asn Ser Gly Ala  
 65                                      70                                      75

<210> 186  
 <211> 99  
 <212> PRT  
 <213> Homo sapiens

<400> 186

Cys Arg Ala His His Ser Leu Thr Ser Phe Val Ser Trp Phe Arg Tyr  
 1                                      5                                      10                                      15  
 Asp Leu Pro Tyr Pro Asp His Ser Ile Asn Cys Lys Leu Pro Val His  
                          20                                      25                                      30  
 Ser Ser Leu Ser Tyr Asn Thr Phe Pro Phe Ser Gln Arg Tyr Cys His  
                          35                                      40                                      45  
 Phe Val Ser Tyr Tyr Ile Thr Tyr Tyr Val Tyr Cys Leu Leu Arg Ile  
                          50                                      55                                      60  
 Leu Cys Ser Leu Met Tyr Leu Lys Tyr Leu Gly Gln Cys Ser Val His  
 65                                      70                                      75                                      80  
 Val Thr Gly Val Gln Gln Arg Leu Leu Asn Glu Ile Phe Asp Asn Cys  
                          85                                      90                                      95

Asp Arg Tyr

<210> 187  
 <211> 194  
 <212> PRT  
 <213> Homo sapiens

<400> 187

Ala Glu Gln Val Leu Val Ile Phe Ala Glu Gln Val Leu Asn Glu Cys  
 1 5 10 15  
 Met Asn Lys Cys Met Asn Val Glu Met Lys Gly Asp Ala Asp Gly Asp  
 20 25 30  
 Asp Ala Asp Gly Asp Asp Asp Ala Asp Gly Asp Asp Ala Asp Gly Asp  
 35 40 45  
 Asp Ala Asp Gly Glu Gln Trp Pro Cys Arg Val Phe Ala Asp Leu Gly  
 50 55 60  
 Leu Ala Ser Gly Cys Gly Gly Ser Ala Ser Gln Gly Phe Glu Phe His  
 65 70 75 80  
 Leu Gln Cys Leu Pro Ala Met Pro Pro Trp Val Thr Phe Ile Leu Leu  
 85 90 95  
 Pro Gly Lys Trp Gly Cys Trp Gln Pro Leu Pro Pro Gly Ile Thr Asp  
 100 105 110  
 Thr Ala Trp Ser Gly Cys Asp Pro Phe Gly Tyr Arg Arg Gly Trp Trp  
 115 120 125  
 Thr Ser Gln Val Gly Arg Ser Ser Leu Asp Glu Arg Pro Arg Thr Ile  
 130 135 140  
 His Arg Arg Ala Gln Glu Ser Leu Leu Ser Pro Ser Asn Ser Thr Glu  
 145 150 155 160  
 Pro Ala Val Asn Cys Trp Leu Leu Pro Val Thr Phe Pro Cys Pro Tyr  
 165 170 175  
 Phe His Ser Leu Glu Ala Ala Arg Thr Thr Ala Gly Trp Pro Trp Pro  
 180 185 190  
 Leu Pro

<210> 188  
 <211> 178  
 <212> PRT  
 <213> Homo sapiens

<400> 188

Ser Phe Ser Leu Gly Asn Phe Val Val Ala Ser Leu Tyr Ser Cys Cys  
 1 5 10 15  
 Phe Asn Asn Phe Val Leu Phe His Ser Phe Thr Val Thr Val Cys Val  
 20 25 30  
 Asp Ser Phe Ser Ser Ser Val Lys Ile Met Ser Pro Glu Ser Ser Phe  
 35 40 45  
 Ile Thr Leu Asp Arg Thr Arg Thr Leu Ser Ile Lys Ser Met Leu Phe  
 50 55 60  
 Val Ile Thr Glu Gln Phe Ser Ala Val Ile Ser Leu Ile Val Thr Phe  
 65 70 75 80  
 Leu Phe Ile Pro Phe Ser Leu Ser Lys Met Pro Leu Phe Val Tyr Trp  
 85 90 95  
 Ser His Arg Ser Glu Ile Cys Glu Phe Ala Ile His Val Ser Tyr Leu

100                      105                      110  
 Phe Ala Asn Gly Phe His Val Ser Lys Ser Leu Phe Ser Ile Val Arg  
       115                      120                      125  
 Tyr Tyr Leu Tyr Cys Phe Val Gln Asn Ile Asn Leu Val Leu Phe Ile  
       130                      135                      140  
 Asp Tyr Ser Leu Val Leu Leu Leu Asn Phe Ile Gln Glu Cys Val Phe  
       145                      150                      155                      160  
 Leu Ser Asp Tyr Phe Phe Leu Pro Asn Cys Ile Phe Leu Arg Gly Leu  
               165                      170                      175

Ile Ile

<210> 189  
 <211> 76  
 <212> PRT  
 <213> Homo sapiens

<400> 189

Pro Arg Glu Ala Lys Arg Leu Asp Ile His Ala Pro Leu Leu Ser Leu  
 1                      5                      10                      15  
 Pro Asp Cys His Leu Leu Met Ala Ala Ser Val Ala Tyr Lys Ile Trp  
               20                      25                      30  
 Arg Pro Leu Gly Ser Val Ser Asn Cys Leu Asn Pro Leu Leu Tyr Phe  
               35                      40                      45  
 Leu Ser Arg Gly Ala Lys Phe Glu Ser Gly Ser Ser Arg Asn Gly Arg  
               50                      55                      60  
 Thr Ser Trp Val Ser Ile Gln Leu Gly Gly Arg Asp  
 65                      70                      75

<210> 190  
 <211> 189  
 <212> PRT  
 <213> Homo sapiens

<400> 190

Ser Leu Val Ile Leu Val Cys Tyr Ser Leu Met Val Arg Ser Leu Ile  
 1                      5                      10                      15  
 Lys Pro Glu Glu Pro His Glu Val Gln Ala Thr Gln Pro Glu Pro Gly  
               20                      25                      30  
 Pro Ser Gly Thr Ile Leu Leu Val Cys Gly Leu Phe Thr Leu Cys Phe  
               35                      40                      45  
 Val Pro Phe His Ile Thr Arg Ser Phe Tyr Leu Thr Ile Cys Phe Leu  
               50                      55                      60  
 Leu Ser Gln Asp Cys Gln Leu Leu Met Ala Ala Ser Val Ala Tyr Lys  
 65                      70                      75                      80  
 Ile Trp Arg Pro Leu Val Ser Val Ser Ser Cys Leu Asn Pro Val Leu  
               85                      90                      95  
 Tyr Phe Leu Ser Arg Gly Ala Lys Ile Glu Ser Gly Ser Ser Arg Asn

100                      105                      110  
 Gly Arg Thr Ser Trp Val Ser Ile Gln Leu Gly Gly Arg Asp Ala Gln  
       115                      120                      125  
 Gly Thr Asp Leu Gly Asn Ala Lys Val Lys Leu Gly Lys Asn Glu Leu  
       130                      135                      140  
 Gln His His Gln Gln Leu Val Cys Thr Gln Met Ser Ala Gly Gly Arg  
       145                      150                      155                      160  
 Gly Ala Gln Asp Leu Leu Lys Val Ser Cys Cys Lys Gly His Phe Tyr  
                               165                      170                      175  
 Ile Asp Val Lys Val Asn Lys Ser Met Glu Arg Ala Thr  
                               180                      185

<210> 191  
 <211> 208  
 <212> PRT  
 <213> Homo sapiens

<400> 191

Ser His Ile Ser Pro Gly Thr Gly Cys Leu Ser Leu Pro Ala Ile Val  
       1                      5                      10                      15  
 Trp Ala Leu Ala Gly Ser Ser Pro Trp Glu Met Trp Ala Arg His Ser  
                               20                      25                      30  
 Asp Arg Ser Gln Ser Ala Gly Ala Gly Ala Phe Gly Leu Ser Ser Pro  
                               35                      40                      45  
 Met Glu Val Ser Glu Pro His Ser His Ser Tyr Arg Arg His Gln Asn  
                               50                      55                      60  
 Ser Leu Tyr Val Glu Pro His Lys Val Glu Thr Val Asn Ser Cys Arg  
       65                      70                      75                      80  
 Asn Leu Leu Trp Asn Thr Thr Val Phe Glu Ser Gly Ser Asp Leu Thr  
                               85                      90                      95  
 Ser Ser Val Thr Leu Gly Lys Leu Leu Leu Pro Trp Thr Pro Thr Thr  
                               100                      105                      110  
 His Leu Asp Val Gly Asn Asn Asp Thr Glu Phe Ile Gly Leu Arg Leu  
                               115                      120                      125  
 His Leu Met Gly Thr Leu Glu Gln Cys Gln Thr Gln Thr Thr Asn Ala  
                               130                      135                      140  
 Gln Lys Leu Val Phe Ile Ile Ala Phe His Phe Asn Cys Gly Leu Leu  
       145                      150                      155                      160  
 Gly Leu Asn Cys Val Pro Ser Lys Arg Tyr Ile Gly Val Leu Thr Leu  
                               165                      170                      175  
 Ser Thr Ser Glu Cys Asp Cys Thr Trp Arg Leu Gly Leu Tyr Arg Asp  
                               180                      185                      190  
 Asn Arg Val Lys Met Glu Leu Gln Gly Trp Ser Leu Ile Gln Cys Asp  
                               195                      200                      205

<210> 192  
 <211> 211

)



<212> PRT  
<213> Homo sapiens

<400> 192

```

Ile Leu Ser Ser Ser Leu Cys Leu Arg Pro Pro Ser Pro Glu Pro Ser
1          5          10          15
Glu Leu Ser Ala Ser Ser Leu Phe Ala Pro Pro Cys Cys Arg His Arg
          20          25          30
Arg Phe Gly Ser Val Pro Ala Glu Val Gly Lys Asp Thr Trp Asn Ser
          35          40          45
Gly Arg Pro Leu Cys Ser Pro Leu Ala Arg Ser Lys Ala Val Lys Asp
          50          55          60
Thr Ala Ser Pro Gly Ser Cys Ser Ser Leu Asn Pro Thr Val Asp Leu
65          70          75          80
Val Gly Arg Leu Arg Ala Gln Ile Cys Arg Cys Ser Ile Val Ser Ser
          85          90          95
Val Ser Cys Pro Leu Leu Pro Pro Gly Val Asp Ser Cys Thr Val His
          100          105          110
Pro Thr Pro Ala Phe Pro Ser Phe Leu Ile Ser Pro Val Ile Phe Pro
          115          120          125
Val Ala Leu Leu Cys Trp Cys Pro Val Arg Ser Cys Gly His Lys Arg
          130          135          140
Leu His Gly Pro His Pro Gln Leu Gly Glu Ser Ser Pro Ser Trp Val
145          150          155          160
Leu Trp Thr Val Lys Lys Asp Gly His Val Gly Ser Val Glu His Glu
          165          170          175
Val Val Gln Asp Leu Gly Gly His Arg Ser Cys Leu Pro Ala Ser Arg
          180          185          190
Ala Leu Pro Pro Phe Gly Ser Leu Leu His Leu Gly Lys Arg Phe Val
          195          200          205
Pro Thr Pro
          210

```

<210> 193  
<211> 208  
<212> PRT  
<213> Homo sapiens

<400> 193

```

Asn Met Ser Tyr Ser Ser Arg Val Asn Ser Leu Leu Leu Phe Ser Phe
1          5          10          15
Asn Phe Ser Tyr Ile Ile Phe His Ile Asn Phe Arg Ile Ser Leu Val
          20          25          30
Trp Gly Val Ile Gln Val Asn Leu Ile Lys Phe Gly Glu Gly Phe Thr
          35          40          45
Ile His Leu Ile Asn Phe Gly Arg Val Val Met Leu Met Phe Ser His
          50          55          60

```

Tyr Ile Leu Lys Cys Asp Ile Ser Phe His Leu Phe Val Leu Asp Gln  
 65 70 75 80  
 Ala Leu Val Ala Ser Ser Glu Asn Leu Leu Asn Ser Arg Asn Asn Phe  
 85 90 95  
 Phe His Leu Leu Thr His Phe Leu Thr Ile Cys Phe Leu Pro Leu Val  
 100 105 110  
 Leu Cys Leu Val Asn Tyr Phe Leu Leu Ile Ser Pro Leu Gln Ile Leu  
 115 120 125  
 Tyr Ala Ile Arg Lys Gly Val Thr Asp Leu Val Ile Glu Thr Gln Tyr  
 130 135 140  
 Thr Phe Val Gly Met Met Lys Ala Leu Gly Ile Phe Ser Tyr Tyr Val  
 145 150 155 160  
 His Leu Ile Ile Leu Lys Leu Ser Ser Tyr Val Glu Pro Ile His Lys  
 165 170 175  
 Ser Arg Ser Phe Asp Phe Lys Ser Cys Ile Phe Pro Tyr Phe Gln Tyr  
 180 185 190  
 Leu Ile Gly Glu Val Thr Cys Asn Ala Ile Val Leu Gln Phe Tyr Ile  
 195 200 205

<210> 194  
 <211> 213  
 <212> PRT  
 <213> Homo sapiens

<400> 194

Met Thr Gly Asn Ala Val Val Leu Trp Leu Leu Gly Phe Arg Met Arg  
 1 5 10 15  
 Arg Asn Ala Phe Ser Ile Tyr Ile Phe Asn Leu Ser Met Ala Asp Phe  
 20 25 30  
 Leu Phe Leu Arg Ser His Ile Ile Arg Phe Pro Leu Ser Leu Ile Asn  
 35 40 45  
 Ile Leu His Pro Ile Phe Lys Ile Leu Ser Pro Val Met Met Phe Ser  
 50 55 60  
 Tyr Leu Ala Ser Leu Ser Phe Leu Ser Ala Met Ser Thr Glu Arg Cys  
 65 70 75 80  
 Leu Tyr Val Leu Trp Pro Ile Trp Arg Cys Arg Pro Arg Pro Tyr Thr  
 85 90 95  
 Cys Gln Arg Ser Cys Val Ser Cys Ser Gly Pro Cys Leu Cys Cys Gly  
 100 105 110  
 Ala Ser Trp Ser Gly Val Ser Val Thr Ser Cys Leu Val Val Leu Ile  
 115 120 125  
 Leu Phe Gly Val Lys His Gln Ile Ser Ser Gly Gly Phe Phe Tyr Val  
 130 135 140  
 Trp Leu Ser Val Val Pro Ala Trp Ser Cys Trp Ser Gly Ser Phe Val  
 145 150 155 160

```
<210> 195
<211> 190
<212> PRT
<213> Homo sapiens
```

<400> 195

His 1	Thr	His	Thr	His 5	Thr	His	Thr	His	Thr	His	Thr	His	Thr	His	Thr	Arg 15	Thr
His	Pro	Ile	Asn 20	Gly	Phe	Pro	Gly	Gly 25	Arg	Ala	Ser	Val	Pro 30	Leu	Thr		
Ala	Gly	Pro 35	Pro	Gly	Pro	Ala	Lys 40	Gly	Ala	Lys	Ser	His 45	Ser	Asp	Ile		
Asn 50	Ser	Trp	Phe	Gln	Ser	Asn 55	Lys	Gln	Ser	Asn	Val 60	Arg	Lys	Val	Ile		
Arg 65	Leu	Lys	Gly	Phe	Glu 70	Gly	Lys	Ser	His	Gln 75	Lys	Val	Lys	Leu	Asp 80		
Pro	Thr	Ser	Thr	Ser 85	Trp	Met	Ser	Tyr	Leu 90	Ile	Ser	Leu	Ala	Ser 95	Val		
Phe	Ser	Pro	Ile 100	Lys	Lys	Pro	Glu	Asp 105	Leu	Pro	His	Gln	Ala 110	Val	Leu		
Lys	Leu	Asn 115	Glu	Leu	Ile	Pro	Val 120	Gln	Ala	Glu	Asn	Ser 125	Ile	Tyr	Ser		
Ile 130	Ser	Gln	Leu	Leu	Leu	Leu 135	Leu	Leu	Leu	Leu	Cys	Thr 140	Trp	Leu	Ser		
Leu 145	Phe	Ser	Phe	Ile	Asn 150	Tyr	Tyr	Ser	Leu	His 155	Leu	Phe	Ala	Ala	Thr 160		
Trp	Ser	Ser	Trp	Asn 165	Pro	Phe	Thr	Ala	Tyr 170	Ser	Arg	Glu	Thr	Gly 175	Glu		
Gly	Arg	Cys	His 180	Leu	His	Ser	His	Trp 185	Asp	Ala	Pro	Ala	Pro 190				

```
<210> 196
<211> 138
<212> PRT
<213> Homo sapiens
```

<400> 196

Glu Asn Leu Phe Phe Lys Gly Lys Phe Val Ser Asn Thr Leu Pro His  
1 5 10 15

Ser Phe Ile Arg Gln Cys Phe Leu Cys His Phe Ser Ala Arg Ile Leu  
 20 25 30  
 Leu Leu Gly Ile Glu Phe Thr Val His Ser Ser Val Leu Ser Val Leu  
 35 40 45  
 Gln Lys Tyr Tyr Leu Phe Pro Ser Asn Leu His Gly Phe Arg Trp Lys  
 50 55 60  
 Ile Cys Cys Gly Leu His Tyr Cys Phe Ser Val Arg Asn Val Pro Phe  
 65 70 75 80  
 Phe Leu Cys Leu Leu Ser Arg Phe Leu Ile Phe Phe Phe His Phe Gln  
 85 90 95  
 Lys Leu Asn Val Phe Gly Cys Ile Leu Phe Arg Val Cys Ser Cys Phe  
 100 105 110  
 Leu Glu Tyr Leu Gly Leu Cys Ser Ser Ile Leu Ile Trp Glu Gly Ser  
 115 120 125  
 His Tyr Phe Leu Ile Val Phe Ser His Ile  
 130 135

<210> 197  
 <211> 175  
 <212> PRT  
 <213> Homo sapiens

<400> 197

Ser Asp Ser Pro Ile Tyr Asn Leu Cys His Thr Asn Arg Leu Asn Pro  
 1 5 10 15  
 His Cys Glu Phe His Thr Cys Val Asp Val Ser Thr Ser Arg Asp Gly  
 20 25 30  
 Cys Ile Phe Phe Ile Phe Leu His Thr Phe Leu Glu Tyr Phe Ile Ser  
 35 40 45  
 Met Val Leu Gln Ile Leu Leu Pro Thr Tyr Cys Gly Phe Lys Ala Met  
 50 55 60  
 Glu Lys Thr Lys Ser His Arg Ser Lys Tyr Cys Arg Lys Gln Asn Ser  
 65 70 75 80  
 Trp Val Asp Leu Ile Phe Leu Tyr Lys Asn Tyr Gly Tyr Gly Tyr Met  
 85 90 95  
 Tyr Leu Cys Met Ser Val Ala Lys Ile Asn Lys Met Asn Thr Phe Asn  
 100 105 110  
 Leu Arg Val Pro Ile Ile Gln Phe Thr Ser Phe Cys Pro Thr Thr Leu  
 115 120 125  
 Glu Ala Lys Thr Leu Val Glu Thr Leu Met Cys Phe Thr Ser Asn Ser  
 130 135 140  
 Ser Leu Ala Leu Asn Ile Pro Leu Phe Val His Pro Leu Ser Asp Ala  
 145 150 155 160  
 Ile Leu Leu Val Lys Gln Gln Thr Ser Thr His Arg Lys Leu Glu  
 165 170 175

<210> 198

<211> 177  
 <212> PRT  
 <213> Homo sapiens

<400> 198

```

Ser Arg Lys Gly Arg His Trp Arg Gly Cys Leu Leu Thr Leu Leu Met
1          5          10          15
Leu Val Ala Val Val Val Cys Phe Ser Pro Tyr His Leu Asn Ile Lys
          20          25          30
Gln Phe Met Ala Arg Gly Met Leu His Leu Pro Ser Cys Ala Glu Arg
          35          40          45
Arg Ala Phe Leu Leu Ser Leu Gln Ala Thr Val Ala Leu Met Asn Met
          50          55          60
Asn Cys Gly Ile Thr Pro Ser Phe Thr Ser Leu His Pro Pro Ile Thr
65          70          75          80
Gly Asn Gly Ser Trp Ala Phe Ser Ser Lys Gly Leu Pro Pro Pro Pro
          85          90          95
Pro Pro Pro Pro Pro Gln Glu Lys Leu Leu Gln Lys His Gln Val Ser
          100          105          110
Pro Arg Pro Glu Val Leu Cys Ser Arg Ser Thr Trp Ser Asn Val Ser
          115          120          125
Phe Ala Leu Leu Tyr Leu Gly Arg Gly Pro Ala Leu Gly Tyr Ser Tyr
          130          135          140
Asn Leu Gly Lys Arg Phe Phe Lys Glu Lys Asn Thr Glu Glu Ile Gln
145          150          155          160
Asn Ala Gly Arg Gly Gly Ser Arg Leu Ser Pro His Phe Gly Arg Pro
          165          170          175

```

Arg

<210> 199  
 <211> 202  
 <212> PRT  
 <213> Homo sapiens

<400> 199

```

Val Tyr Glu Cys Tyr Ile Phe Gly His Cys Trp Asp Val Ala Ser His
1          5          10          15
His Leu Thr Ser Leu Asn Leu Ser Gly Leu Thr Cys Glu Met Gly Ala
          20          25          30
Leu Thr Phe Thr Cys Leu Gln Ala Cys Ser Gln Ile Arg Cys His Leu
          35          40          45
Lys Asp Phe Ser Ser Pro Gly Asp Phe Lys Arg Leu Leu Arg Gly His
          50          55          60
Phe Phe Ser Gly Cys Gly Arg Ser Met Ile Arg Val Ile Arg Met Gly
65          70          75          80
Leu Leu Glu Glu Arg Gly Gly Gln Arg Leu Leu Phe His Phe Met Ala

```

85                      90                      95  
 Pro Ser Gly Gln Arg Thr Asp Ser Ala Thr Ala Ala Thr Arg Ala Leu  
                          100                      105                      110  
 Pro Gly Leu Trp Ser Gln Leu Ser Gln Gln Glu Phe Gln Lys Ala Lys  
                          115                      120                      125  
 Gly Ser Glu Leu His Pro Ser Phe Leu Ala Asp Cys His Pro Ala Ser  
                          130                      135                      140  
 Ser His Ser Pro Gln Gly Tyr Val Met Leu Ala Leu Lys Ala Ser Leu  
                          145                      150                      155                      160  
 Gly Arg Gly Cys Ile Cys His Pro Leu Pro Cys Lys Ile Phe Glu Val  
                          165                      170                      175  
 Gln Arg Ala Leu Gln Ala Glu Pro His Pro Leu Leu His Ser Pro Ser  
                          180                      185                      190  
 Val Gly Met His Ser Pro Ser Val Gly Met  
                          195                      200

<210> 200  
 <211> 175  
 <212> PRT  
 <213> Homo sapiens

<400> 200

Leu Pro Pro Pro Ile Leu Val Pro Thr Val Val Thr Glu Glu Ile  
 1                      5                      10                      15  
 Phe Ser Ser Ser Thr Ala Thr Leu Lys Gly Pro Ser Val Pro Phe Gly  
                          20                      25                      30  
 Gly Leu Gly Ile Asp Leu Pro His Arg Ser Ser Leu Ala Pro Met His  
                          35                      40                      45  
 Thr Phe Arg Asp Leu Arg Thr Gly Pro Leu Cys Leu Pro Leu Ser Leu  
                          50                      55                      60  
 Leu Val Arg Lys Asp Trp Pro Ala Cys Leu His Pro Gln Gln Ser Ile  
                          65                      70                      75                      80  
 Ala Thr Ala Pro Ser Cys Ala Thr Glu Glu Leu Thr Asp Thr Thr His  
                          85                      90                      95  
 Thr Val Tyr Ser Arg Arg Asn Pro Met Gly Pro Ile Ile Leu Cys Pro  
                          100                      105                      110  
 Pro Trp Ile Lys Thr Lys Val Leu Tyr Ala Thr Asn Thr Thr Ala Ile  
                          115                      120                      125  
 Ser Thr Gly Lys Ser Leu Ser Leu Gln Lys Pro Ile Gln Lys Pro Arg  
                          130                      135                      140  
 Arg Ser Asn Cys His Thr Lys Tyr Thr Asp Thr Asn Leu Arg Thr Glu  
                          145                      150                      155                      160  
 Thr Glu Asn Lys Glu Thr Trp His Phe Leu Lys Glu His Asn Asn  
                          165                      170                      175

<210> 201  
 <211> 178

<212> PRT  
<213> Homo sapiens

<400> 201

```

Leu Gly Phe Leu Leu Thr Asp Val Gln Ser Val Phe Gly Tyr Leu Gln
1           5           10           15
His Glu Thr His Tyr Cys Ser Ala Thr Ile Gly Arg His Trp Pro Ala
      20           25           30
His Pro Leu Met Arg Cys Trp Asn Pro Phe Phe Ile Leu Lys Tyr Leu
      35           40           45
Ile Asp Lys Asn Cys Val Cys Ser Arg Cys Asp Val Met Leu Arg Ser
      50           55           60
Arg Tyr Ile Gln Val Tyr Leu Pro Gln Ser Asn Leu Thr Asn Leu Ser
      65           70           75           80
Pro Pro Met Ile Thr Ile Met Leu Arg Gly Gly Ser Glu Asp Thr Lys
      85           90           95
Asp Leu Leu Ser Tyr Gln Ile Ser Ser Gln Gln Tyr Ser Ile Ile Asn
      100          105          110
Thr Val Thr Met Leu Cys Ile Arg Ser Pro Glu His Val Thr Glu Gly
      115          120          125
Leu Tyr Leu Leu Thr Asn Ile Ser Pro Ala Leu His Glu Trp Met Val
      130          135          140
Ser Ile Phe Gln Thr His Ser Glu Asp Phe Ala Trp Leu Ala Thr Ser
      145          150          155          160
Ile Ser Pro Glu Lys Val Gln Lys Ser Arg Pro Ser His Arg Asn Ser
      165          170          175

Asp Ala

```

<210> 202  
<211> 196  
<212> PRT  
<213> Homo sapiens

<400> 202

```

Tyr Gly Ala Leu Tyr Lys Tyr Lys Gln Gln Ser Leu Thr Phe Leu Ser
1           5           10           15
Leu Gln Leu Leu Thr Leu Ala Gly Ser Arg Ile Lys Met Pro Asn Ser
      20           25           30
Thr Gln Lys Pro Trp Pro Val Ser Leu Pro Lys Met Glu Phe Arg Leu
      35           40           45
Thr Ala Gly Asn Arg Asn Cys Ser Phe Lys Ala Ile Ala Trp Ala Met
      50           55           60
Val Pro Ile Phe Val Asn Ile Gly Phe Cys Leu Asn Ser Val Ser Arg
      65           70           75           80
Val Asp Tyr Ile Ile Cys Lys Val Cys Lys Met Lys Val Trp Gly Ser
      85           90           95

```

Ser Ser Lys Tyr Lys Gln Lys Val Leu Leu Ser Val Ser Lys Tyr Lys  
 100 105 110  
 Met Phe Pro Leu Ser Val Ile Tyr Phe Ser Thr Cys Tyr Val Phe Gln  
 115 120 125  
 Phe Val Cys Phe Val Phe Pro Leu Leu Phe Tyr Val Leu Leu Cys Lys  
 130 135 140  
 Lys Ile Lys Asn Leu Asn Tyr His Asn Lys Phe Ser His Ser Phe Leu  
 145 150 155 160  
 Cys Cys Ala Val Ser Ile Asn Ala Asn Ile Lys Ala Phe Asn Leu Tyr  
 165 170 175  
 Ile Glu Ser Gln Lys Leu His Asn Thr Tyr Phe Ile Val Cys Thr Cys  
 180 185 190  
 Met Tyr Ile Leu  
 195

<210> 203  
 <211> 212  
 <212> PRT  
 <213> Homo sapiens

<400> 203

Ser Gly Val Ile Asn Leu Leu Tyr Ile Cys Val Tyr Val Cys Ile Phe  
 1 5 10 15  
 Leu Pro Asn Arg Cys Asn Thr Lys Tyr Ser His Gly Val Ile Thr Phe  
 20 25 30  
 Ser Gln Leu Thr Leu His Pro Tyr Ile Ile Glu Glu Arg Ser Thr Ser  
 35 40 45  
 Ile Leu Phe Leu Leu Val Ile Ala Leu Met Ser Glu Tyr Lys Leu Asp  
 50 55 60  
 Ser Ser Val Ala Asn Asn Thr Arg Gln Ser Lys Asp Phe Ser Cys Cys  
 65 70 75 80  
 Arg His Ile Phe Leu Ile Tyr Trp Lys His Lys Cys Val Pro Pro Asn  
 85 90 95  
 Phe Ile Val Asp Arg Asn Met Lys Asn Phe Ile Lys Leu Lys Thr Gly  
 100 105 110  
 Ser Leu Pro Asp Leu Pro Val Ile Leu Pro Thr Leu Gln Ile His Pro  
 115 120 125  
 Ile Val Pro Ala Ser Phe Thr Met Lys Lys Tyr Glu Thr Cys Leu Thr  
 130 135 140  
 Trp Ser Leu Cys Leu Arg Glu Thr Cys Val Cys Leu Trp Asn Thr Leu  
 145 150 155 160  
 Thr Lys Ile Pro Ala Leu Val Asp Lys Thr Gly Phe Gln Ser Ser Leu  
 165 170 175  
 Asn Ser His Phe Val Leu Asn Lys Val Val Ser Lys Thr Arg Cys Ser  
 180 185 190



Lys Tyr Tyr Cys Ser Asp Ala Ile Ser Lys Thr Val Leu Ile Pro Cys  
 195 200 205

Gly Arg Glu Asn  
 210

<210> 204  
 <211> 172  
 <212> PRT  
 <213> Homo sapiens

<400> 204

Asn Lys Ile Val Phe Ile Phe Ser His Asp Cys Leu Trp Arg Lys Ile  
 1 5 10 15

Ser Lys Asn Leu Pro Lys Thr Asn Ala Ile Leu Ser Arg Val Lys Glu  
 20 25 30

Thr Arg Ser Ser Leu Phe Cys Thr Leu Tyr Phe Cys Ile Ser Val Leu  
 35 40 45

Phe Leu Tyr Gly Ser Asn Asp Gln Leu Glu Ile Lys Ile Leu Lys Gln  
 50 55 60

His Gln Lys His Lys Met Leu Ser Tyr Lys Ser Asn Lys Thr Tyr Thr  
 65 70 75 80

Asp Ser Val Pro Lys Thr Val Asn Val Tyr Leu Lys Asn Gln Arg Arg  
 85 90 95

Ala Glu Gln Arg Ala Thr Ser Cys Leu Leu Leu Glu Asn Ser Ile Glu  
 100 105 110

Leu Arg Tyr Lys Phe Pro Gln Ser Asp Leu Asp Ala Thr Gln Phe His  
 115 120 125

Ser Asn Pro Ser Arg His Phe Leu Leu Lys Ser Thr Ser Cys Phe Ile  
 130 135 140

His Thr Lys Ile His Lys Asn Lys Lys Ala Lys Ile Leu Leu Lys Glu  
 145 150 155 160

Asn Lys Phe Arg Arg Leu Leu Leu Ser Asp Phe Arg  
 165 170

<210> 205  
 <211> 313  
 <212> PRT  
 <213> Homo sapiens

<400> 205

Val Pro Lys Ile Phe Ser Phe Ser Ser Ser Phe Gln Asn Tyr Phe Leu  
 1 5 10 15

Ile Leu Val Lys His Thr Ser Ser Asn Ile Thr Tyr Tyr Leu Val Phe  
 20 25 30

Thr Tyr Ile Thr His Ser Leu Asn Lys Phe Val Glu Met Ile Ile Leu  
 35 40 45

Lys Ile Leu Val Phe Lys Phe Met Ser Ser Gln Lys Leu Leu Pro Arg  
 50 55 60

Ile Ser Ile Leu Asn Ile Trp Ile Asn Ile Leu Phe Tyr Thr Pro Tyr  
 65 70 75 80  
 Asn Ile Leu Leu Ala Ile Ile Ile Phe Phe Arg Ile Cys Ser Thr Ser  
 85 90 95  
 Asn Phe Phe Asp Phe Leu Ile Leu Lys Arg Ile Ile Tyr Ala Asn Gln  
 100 105 110  
 Gln Cys Lys Asp Phe Ser Trp Phe Thr Arg Val Lys Leu Phe Ser Arg  
 115 120 125  
 Met Val Gly Ser Phe Ala Tyr Ile Lys Leu Met Tyr Arg Ser Ala Ser  
 130 135 140  
 Ser His Ile Lys Val Gln Ser Leu Leu Lys Lys His Phe Ile Ser Asn  
 145 150 155 160  
 Gln Phe Val Phe Leu Tyr Thr Leu Lys Pro Phe Asn Cys Phe Tyr Phe  
 165 170 175  
 Ser Ile Leu Thr Ser Ile Ser Cys Tyr Ser Gln Trp Pro Ala Ser Ser  
 180 185 190  
 Leu Ala Ile Arg Gln Leu Phe Val Tyr Leu Ala Lys Tyr Ile His Ala  
 195 200 205  
 Leu Lys Ile Pro Phe Pro Asn Ile Tyr Tyr Asp Phe Phe Lys Gly Phe  
 210 215 220  
 Ser Phe Val Thr Met Thr Leu Lys Ala Lys Val Ser Arg Cys Cys Ile  
 225 230 235 240  
 Thr Val Gly Ser Thr Ile Met Tyr Gln Glu Gly Arg Glu Asn Gln Gly  
 245 250 255  
 Thr Phe Leu Trp Glu Tyr Pro Ile Ile Cys Gln Ile Tyr Ser Asn Ser  
 260 265 270  
 Leu Arg Thr Ile Thr Phe Val Phe Thr Val Phe Pro Met Gln Phe Leu  
 275 280 285  
 Arg Phe Ile Phe Lys Asn Phe Leu Gly Glu Met Asp Tyr Ser Leu Leu  
 290 295 300  
 Ser Ala Val Ile His Asn Phe Tyr Phe  
 305 310

<210> 206  
 <211> 318  
 <212> PRT  
 <213> Homo sapiens

<400> 206

Pro Phe Tyr Tyr Ser Met Leu Val Pro Thr Ser Gly Leu Ser Thr Cys  
 1 5 10 15  
 Cys Ser Phe Cys Leu Glu Ser Ser Ser Pro Asp Leu Leu Arg Phe Pro  
 20 25 30  
 Leu Ser Ile Arg Val Ser Ala Val Ile His Pro Gln Arg Arg Ser Pro  
 35 40 45  
 Asp Pro Val Lys Pro Pro Ile Pro Gln Ser Pro Tyr Val Ser Thr Ser

50                      55                      60  
 Leu Tyr Leu Ile Ser Gln His Leu Leu Ile Ser Leu Thr Leu His Tyr  
 65                      70                      75                      80  
 Met Cys Cys Tyr Met Phe Val Ile Leu Ser Ser Gly Pro Cys Asn Val  
                     85                      90                      95  
 Arg Met Ala Gln Tyr Lys Trp Gln Glu Gly Cys Arg Gly Val Asp Lys  
                     100                      105                      110  
 Ala Glu Ser Gly Trp Gly Ser Trp Arg Asp Gly Gln Gly Pro Glu Leu  
                     115                      120                      125  
 Arg Arg Trp Tyr Leu Gln Cys Ala Leu Asn Cys Pro Gly Met Ile Ile  
                     130                      135                      140  
 Ser Ile Ala Ser Phe His Ser Gln Arg Cys Pro Gly Tyr Tyr Ser Cys  
 145                      150                      155                      160  
 Ser Val Tyr Arg Ala Trp Ala Val Gly Ile Leu Phe Gln Met Gly Cys  
                     165                      170                      175  
 Glu Ala Cys Gly Trp Phe Ala Gly Ser Asp Met Ile Leu Ala Phe Lys  
                     180                      185                      190  
 Asp His Asp Gln Val Leu Glu Thr Leu Phe Trp Leu Leu Pro Thr Pro  
                     195                      200                      205  
 Pro His Thr His Pro Thr Leu Leu His Cys Pro Phe Ser Leu Leu Trp  
                     210                      215                      220  
 Gln Leu Phe Leu Phe Tyr Asn Leu Ile Leu Glu Phe Leu Gln Thr Ser  
 225                      230                      235                      240  
 Gly Ser Gln Leu Gly Ala Ile Ser Pro Pro Arg Asp Ile Trp Tyr Phe  
                     245                      250                      255  
 Ile Trp Arg Tyr Phe Trp Ser Gln Leu Glu Arg Val Leu Ala Ser Ser  
                     260                      265                      270  
 Gly Arg Pro Gly Arg Leu Leu Thr Ile Leu Gln Ser Thr Glu Gln Pro  
                     275                      280                      285  
 Tyr Thr Ile Lys Asn Asp Leu Thr Gln Asn Ala Ser Ser Pro Glu Val  
                     290                      295                      300  
 Lys Lys Pro Cys Thr Arg Leu Ala Pro Ser Asn Arg Asn Ile  
 305                      310                      315  
 <210> 207  
 <211> 318  
 <212> PRT  
 <213> Homo sapiens  
 <400> 207  
 Ile Ser Pro Phe Tyr Tyr Ser Met Leu Val Pro Thr Ser Gly Leu Ser  
 1                      5                      10                      15  
 Thr Cys Cys Ser Phe Cys Leu Glu Ser Ser Ser Pro Asp Leu Leu Arg  
                     20                      25                      30  
 Phe Pro Leu Ser Ile Arg Val Ser Ala Val Ile His Pro Gln Arg Arg  
                     35                      40                      45

Ser Pro Asp Pro Val Lys Pro Pro Ile Pro Gln Ser Pro Tyr Val Ser  
 50 55 60  
 Thr Ser Leu Tyr Leu Ile Ser Gln His Leu Leu Ile Ser Leu Thr Leu  
 65 70 75 80  
 His Tyr Met Cys Cys Tyr Met Phe Val Ile Leu Ser Ser Gly Pro Cys  
 85 90 95  
 Asn Val Arg Met Ala Gln Tyr Lys Trp Gln Glu Gly Cys Arg Gly Val  
 100 105 110  
 Asp Lys Ala Glu Ser Gly Trp Gly Ser Trp Arg Asp Gly Gln Gly Pro  
 115 120 125  
 Glu Leu Arg Arg Trp Tyr Leu Gln Cys Ala Leu Asn Cys Pro Gly Met  
 130 135 140  
 Ile Ile Ser Ile Ala Ser Phe His Ser Gln Arg Cys Pro Gly Tyr Tyr  
 145 150 155 160  
 Ser Cys Ser Val Tyr Arg Ala Trp Ala Val Gly Ile Leu Phe Gln Met  
 165 170 175  
 Gly Cys Glu Ala Cys Gly Trp Phe Ala Gly Ser Asp Met Ile Leu Ala  
 180 185 190  
 Phe Lys Asp His Asp Gln Val Leu Glu Thr Leu Phe Trp Leu Leu Pro  
 195 200 205  
 Thr Pro Pro His Thr His Pro Thr Leu Leu His Cys Pro Phe Ser Leu  
 210 215 220  
 Leu Trp Gln Leu Phe Leu Phe Tyr Asn Leu Ile Leu Glu Phe Leu Gln  
 225 230 235 240  
 Thr Ser Gly Ser Gln Leu Gly Ala Ile Ser Pro Pro Arg Asp Ile Trp  
 245 250 255  
 Tyr Phe Ile Trp Arg Tyr Phe Trp Ser Gln Leu Glu Arg Val Leu Ala  
 260 265 270  
 Ser Ser Gly Arg Pro Gly Arg Leu Leu Thr Ile Leu Gln Ser Thr Glu  
 275 280 285  
 Gln Pro Tyr Thr Ile Lys Asn Asp Leu Thr Gln Asn Ala Ser Ser Pro  
 290 295 300  
 Glu Val Lys Lys Pro Cys Thr Arg Leu Ala Pro Ser Asn Arg  
 305 310 315  
 <210> 208  
 <211> 320  
 <212> PRT  
 <213> Homo sapiens  
 <400> 208  
 Lys Leu Thr Leu Ala Ala Tyr Thr Leu Ile Gln Cys His Leu Pro Cys  
 1 5 10 15  
 Val Ile His Asn Ile Leu Tyr Glu Ser Tyr Phe Leu Cys Val Cys Val  
 20 25 30

Pro Phe Phe Glu Glu Tyr Asp Leu Ser Gln Phe Phe Cys Phe Ser Leu  
 35 40 45  
 Ser Pro Phe Asn Ile Ser Arg Ala Phe Val Val Val Thr Gly Glu Thr  
 50 55 60  
 Thr Tyr Thr Ser Phe Leu Leu Leu Phe Cys Tyr Leu Gln Phe Cys Met  
 65 70 75 80  
 Thr Leu Lys Gln Lys Asn Asn Tyr Leu Thr Ile Ser Phe Val Leu Tyr  
 85 90 95  
 Ser Gly Phe His Ile Gln Ser Pro Phe Ile Met Leu Leu Pro Leu Phe  
 100 105 110  
 Ser Ser Val Phe Glu Asp Gly Lys Ile His Gln His Pro Lys Tyr Gln  
 115 120 125  
 Pro Glu Arg Lys Lys Glu Ser Gly Trp Arg Gln Asp Ser Phe Gln Ser  
 130 135 140  
 Ile Ser Ser Thr Asp His Gly Ala Ala Ala Lys Arg His Ser Lys Arg  
 145 150 155 160  
 Val Glu Arg Gly Lys Thr Ser Ser Leu Arg Cys Leu Pro Phe Lys Phe  
 165 170 175  
 Thr Ile Ile Ile Arg Met Leu Leu Glu Glu Glu Gln Gly Gln Gly His  
 180 185 190  
 Phe Cys Asn Met Thr Gln Lys Asn Ile Asp Leu Lys Phe Asp Thr Tyr  
 195 200 205  
 Glu Leu Ser Lys Cys Arg Glu Lys Leu Pro Pro Cys Cys Thr Cys Met  
 210 215 220  
 Cys Ala Ile His Phe Ile Leu Ile Lys Val Cys Lys His Glu Met Gln  
 225 230 235 240  
 Gly Thr Asp His Leu Phe Met Arg Met Gln His Ser Ser Glu Lys Val  
 245 250 255  
 Tyr Leu Pro Lys Thr Glu Tyr Met Phe Ile Leu Lys Phe Phe Phe Leu  
 260 265 270  
 Phe Leu Phe Leu Ile Val Ile Lys Tyr Lys His Lys Phe Thr Ile Leu  
 275 280 285  
 Ile Ile Phe Lys Tyr Thr Val Gln Tyr Val His Ser His Tyr Cys Ala  
 290 295 300  
 Thr Asn Phe Gln Asn Ser Phe Tyr Leu Ala Lys Met Lys Leu Tyr Thr  
 305 310 315 320

&lt;210&gt; 209

&lt;211&gt; 315

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 209

Gln Pro Phe Ser Met His Ser Leu Glu Glu Lys Phe Phe Phe Phe Leu  
 1 5 10 15

Asn His Tyr Ser Ala Thr Ser Ile Ser Leu Glu Phe Leu Ser Ser Glu

20					25					30									
Thr	Leu	Val	Gln	Val	Ser	Trp	Gly	Ile	Arg	Ile	Val	Cys	Val	Trp	Ile				
35					40					45									
Thr	Lys	Tyr	Tyr	Arg	Leu	Arg	Gly	Glu	Glu	Thr	Leu	Trp	Ser	Phe	Arg				
50					55					60									
Pro	Thr	Leu	Ile	Cys	Leu	Asp	Leu	Phe	Cys	Phe	Lys	Glu	Ser	His	Leu				
65					70					75									
Gln	Arg	Thr	Ala	Ser	Asp	Ser	Pro	Cys	Ser	Val	Phe	Ser	Gln	Glu	Cys				
85					90					95									
Ser	Leu	His	Gln	Pro	Gln	Glu	Val	Leu	Gln	Lys	Glu	Val	Phe	His	Val				
100					105					110									
Gln	Ile	Thr	Leu	Arg	Ser	Asn	Ser	His	His	Ile	Asp	Phe	Glu	Tyr	Ser				
115					120					125									
Cys	Arg	Lys	Thr	Cys	Leu	Tyr	Gln	Leu	Gly	Val	Ser	Pro	Asn	Leu	Phe				
130					135					140									
Gly	His	Gly	Asn	Ser	Phe	Ser	Lys	Lys	Thr	Cys	Phe	Ser	Ile	Ser	Phe				
145					150					155									
His	Arg	Lys	Leu	Thr	Val	Val	Cys	Val	Phe	Phe	Gln	Ile	Ile	His	Ile				
165					170					175									
Tyr	Ser	Lys	Leu	Lys	Leu	His	Trp	Leu	Phe	Gly	Phe	Ile	Asn	Pro	Leu				
180					185					190									
Thr	Ser	Val	Leu	Phe	Phe	Ser	Thr	Thr	Cys	Cys	Leu	Ala	Thr	Ser	Ala				
195					200					205									
Cys	Phe	Val	Trp	Leu	Asp	Phe	Leu	Val	Leu	Ser	Ile	Gly	Leu	Arg	Phe				
210					215					220									
Tyr	Ile	Leu	Ser	Cys	Trp	Asn	His	Pro	Thr	Ser	Pro	Ala	Trp	Leu	Phe				
225					230					235									
Gly	Ser	Arg	Leu	Ser	His	Leu	Val	His	Ser	Ser	Ala	Val	Asp	Leu	Tyr				
245					250					255									
Tyr	Ser	Leu	Met	Ser	Ala	Tyr	Ser	Leu	His	Leu	Tyr	Ser	Phe	Cys	Leu				
260					265					270									
Glu	Met	Met	Ser	Arg	Thr	Gly	Gln	Gly	Trp	Tyr	His	Ser	Ile	Asn	His				
275					280					285									
His	Pro	Leu	Ile	Leu	Thr	Val	Asn	Leu	Pro	Asn	Lys	Ile	Phe	Gln	Lys				
290					295					300									
Arg	Val	Ser	Asn	Asn	Pro	Cys	Leu	Pro	Leu	Trp									
305					310					315									
<210> 210																			
<211> 327																			
<212> PRT																			
<213> Homo sapiens																			
<400> 210																			
Arg	Val	Pro	Ser	Leu	Pro	Gly	Pro	Pro	Ala	Thr	Val	Cys	Pro	Val	Pro				
1					5					10					15				

Ala Ser Glu Phe Ser Gln His Arg Lys Arg Gly Leu Arg Thr Ile Gln  
 20 25 30  
 Pro Val His Ser Arg Glu Ser Leu Ser Val Ser Gln Arg Leu Met Gly  
 35 40 45  
 Cys Leu Trp Cys Arg Val Thr Pro Ala Ser Pro Cys Gly Gly Cys Ala  
 50 55 60  
 Gly Gly Ala Arg Pro Pro Cys Ala Leu Ser Leu Ala Gln Gly Gln  
 65 70 75 80  
 His Thr Ala His Pro Leu Phe Phe Leu Pro Phe Pro Leu Ala Gln Pro  
 85 90 95  
 Leu Val Val Gly Val Thr Arg Gly Ala Glu Arg Ser Trp Arg Ser Arg  
 100 105 110  
 Ala Cys Pro Gly Pro Val Arg Glu Gly Gly Arg Gly Gln Gln His Pro  
 115 120 125  
 Trp Arg Arg Glu Asp Tyr Ile Ile Phe Ile Tyr His Met Pro Lys Ile  
 130 135 140  
 Ala Leu Leu Arg Ala Phe Asp Ile His Pro Lys Ile Phe Lys His Tyr  
 145 150 155 160  
 Gly Ser Met Ser Gly Cys Ile Ser Asn Met Lys Val Glu Ala Ser Cys  
 165 170 175  
 Pro Ala Pro Ser Pro Leu Trp Glu Asn Phe Val His Val Leu Ser Gln  
 180 185 190  
 Leu Phe Gly Lys Gly Gly Pro Ser His Cys Pro Leu Gly Gly Phe Asp  
 195 200 205  
 Val His Cys Val Gly Arg Ser Leu Pro Ser Ile Leu Phe Tyr Phe Cys  
 210 215 220  
 Arg Ile Ser Ala Gln Ser Gly Ser Ala Trp Gln Phe Ser Cys Ser Ala  
 225 230 235 240  
 Arg Glu Val Leu Cys Pro Gly Leu Cys Asp Phe Arg Arg Arg Glu Gly  
 245 250 255  
 Ser Cys Arg Pro Tyr Leu Gln Trp Leu Pro Pro Gly Ile Pro Val Cys  
 260 265 270  
 Ser Leu Cys Thr Val Gln Arg Arg Ser Gly Ser Trp Trp Arg Asp Gly  
 275 280 285  
 Asp Pro Arg Thr Met Ala Ser Thr Lys Ala Gly Gly Ala Cys Asp Arg  
 290 295 300  
 Arg Trp Thr Met Thr Gln Val Pro Ala Arg Tyr Gly Ser Gly Leu Cys  
 305 310 315 320  
 Arg Glu Gly Ala His Pro Gly  
 325

<210> 211  
 <211> 327  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 211

Cys Gln Phe Gly Ala Leu Gly Tyr Ala Gly Pro Val Arg Arg Val Pro  
 1 5 10 15  
 Ser Leu Pro Gly Pro Pro Ala Thr Val Cys Pro Val Pro Ala Ser Glu  
 20 25 30  
 Phe Ser Gln His Arg Lys Arg Gly Leu Arg Thr Ile Gln Pro Val His  
 35 40 45  
 Ser Arg Glu Ser Leu Ser Val Ser Gln Arg Leu Met Gly Cys Leu Trp  
 50 55 60  
 Cys Arg Val Thr Pro Ala Ser Pro Cys Gly Gly Cys Ala Gly Gly Ala  
 65 70 75 80  
 Arg Pro Pro Pro Cys Ala Leu Ser Leu Ala Gln Gly Gln His Thr Ala  
 85 90 95  
 His Pro Leu Phe Phe Leu Pro Phe Pro Leu Ala Gln Pro Leu Val Val  
 100 105 110  
 Gly Val Thr Arg Gly Ala Glu Arg Ser Trp Arg Ser Arg Ala Cys Pro  
 115 120 125  
 Gly Pro Val Arg Glu Gly Gly Arg Gly Gln Gln His Pro Trp Arg Arg  
 130 135 140  
 Glu Asp Tyr Ile Ile Phe Ile Tyr His Met Pro Lys Ile Ala Leu Leu  
 145 150 155 160  
 Arg Ala Phe Asp Ile His Pro Lys Ile Phe Lys His Tyr Gly Ser Met  
 165 170 175  
 Ser Gly Cys Ile Ser Asn Met Lys Val Glu Ala Ser Cys Pro Ala Pro  
 180 185 190  
 Ser Pro Leu Trp Glu Asn Phe Val His Val Leu Ser Gln Leu Phe Gly  
 195 200 205  
 Lys Gly Gly Pro Ser His Cys Pro Leu Gly Gly Phe Asp Val His Cys  
 210 215 220  
 Val Gly Arg Ser Leu Pro Ser Ile Leu Phe Tyr Phe Cys Arg Ile Ser  
 225 230 235 240  
 Ala Gln Ser Gly Ser Ala Trp Gln Phe Ser Cys Ser Ala Arg Glu Val  
 245 250 255  
 Leu Cys Pro Gly Leu Cys Asp Phe Arg Arg Arg Glu Gly Ser Cys Arg  
 260 265 270  
 Pro Tyr Leu Gln Trp Leu Pro Pro Gly Ile Pro Val Cys Ser Leu Cys  
 275 280 285  
 Thr Val Gln Arg Arg Ser Gly Ser Trp Trp Arg Asp Gly Asp Pro Arg  
 290 295 300  
 Thr Met Ala Ser Thr Lys Ala Gly Gly Ala Cys Asp Arg Arg Trp Thr  
 305 310 315 320  
 Met Thr Gln Val Pro Ala Arg  
 325



<210> 212  
 <211> 310  
 <212> PRT  
 <213> Homo sapiens

<400> 212

```

His Glu Leu Ser Leu Pro Cys Gly Gln Ser Pro Val Ile Lys Lys Glu
1      5      10      15
His Thr Pro Ser Leu Thr Glu Thr Ser Leu Asn Lys Lys Asn Ala His
20     25     30
Gln Arg Asn Ile Glu Phe Lys Tyr Leu Glu Gln Met Ser Glu Ile Ser
35     40     45
His Lys Asn Leu Asn Arg Asn Trp Pro Ser Lys Ser Trp Glu Phe Gly
50     55     60
Asp Ala Asn Phe Ile Leu Ser Ile Leu Glu Gln Ser Lys Ile Asn Thr
65     70     75     80
Thr His Phe Ser Leu Arg Lys Ser Ala Tyr Leu Phe Asp Val Pro Ser
85     90     95
Gly Leu Glu Ile Pro Asn Lys Thr Leu Thr Leu Phe Ile Leu His His
100    105    110
Asn Ile Thr Val Asn Lys Asn Asn Leu Asn Leu Cys Ser Asn Phe Pro
115    120    125
Leu Trp Thr Gln Arg Lys Thr Gln Glu Lys Met Val Glu Cys Val Leu
130    135    140
Asn Lys Val His Tyr Leu Tyr Gln Lys Tyr Ala Val Ile Ser Thr Ser
145    150    155    160
Thr Pro Lys Cys Leu Phe Asn Phe Ala Met Met Tyr Lys Ile Leu Val
165    170    175
Thr Cys Gln Ser Ile Asn Phe Ser Gln Leu Ile Leu Lys Ala Glu Asp
180    185    190
Ser His His Phe Val Cys Phe Ser Val Asn Met Ile Val Phe Val Arg
195    200    205
Lys His Ile Tyr Pro Glu Ser Tyr Gly Pro Met Phe Leu Thr Phe Cys
210    215    220
Pro Arg Ser Val Cys Val Ala Ser Cys Val Cys Met Asp Val Asp Asn
225    230    235    240
Lys Leu Asp Ser Tyr Gln Glu Ser Lys Ile Lys Leu Leu Ser Cys Lys
245    250    255
Lys Phe Val Lys Tyr Val Asp Leu Ser Cys Leu Lys Leu Arg His Pro
260    265    270
Gly His Ser Leu Trp Arg Glu Asn Ser Pro Pro Leu His Val Asn Leu
275    280    285
Trp Val Gly Thr Gly Val Gln Gly Phe Arg Val Gly Leu Leu Leu Pro
290    295    300

```

Gly Met Ile Gln Lys Ile  
305 310

<210> 213  
<211> 314  
<212> PRT  
<213> Homo sapiens

<400> 213

Lys Ala Asp Lys Ile Thr Phe Leu Glu Ser Ser Ile Tyr Ser Leu Ile  
1 5 10 15

Val Phe Leu Tyr Ile Thr Leu Ser Gln Leu Trp Ser Lys Glu His Ser  
20 25 30

Thr Glu Glu Gly Gly Ser Leu Ile Phe Pro His Leu Val Thr Pro Met  
35 40 45

Leu Glu Leu His Glu Ile Asp Asn Tyr Tyr Tyr Ile Val Ile Ser Phe  
50 55 60

His Val Leu Ser Phe Ser Ser Ser Leu Leu Leu Phe Phe Lys Ser Arg  
65 70 75 80

Lys Gln Asn Gly His Gln Leu His Glu His Cys Ser Lys Lys Ile Thr  
85 90 95

Val Arg Pro Asn Leu Asn Cys Trp Leu Pro Gly Arg Ala Ile Leu Ile  
100 105 110

Ala Tyr Lys Asp Gln Ile Lys Tyr Gln Ser Gln Val Val Arg Cys Pro  
115 120 125

Cys Thr Glu His Asn Ile Val Tyr Lys Asp Val Glu Leu Leu Leu Leu  
130 135 140

Leu Trp Phe Tyr Thr Val Ala His Asp Lys Glu Leu Ile Phe Tyr Leu  
145 150 155 160

Asn Glu Val Leu Phe Tyr Ile Thr Tyr Phe Met Phe Phe Pro Gln Glu  
165 170 175

Ser Phe Asn Leu Leu Arg Leu Arg Asp Ser Phe Lys Cys Phe Asp Pro  
180 185 190

His Thr Leu Phe Ala Gly Cys Arg Arg Met Cys Met Ile Leu Thr Phe  
195 200 205

Thr Ala Asn Leu Phe Phe Trp Met Gly Tyr Cys Asn Phe Leu Leu Glu  
210 215 220

Asp His Thr Ser Ser Ser Met Phe Arg Arg Gly Leu His Leu Trp Phe  
225 230 235 240

His Gly Trp Thr Leu Asp Pro Leu Trp Leu Ser Lys Ile Leu His Gln  
245 250 255

Cys Asn Ser Phe Val Asn Gly Tyr Met Ile Gln Ala Gly Pro Ile Arg  
260 265 270

Ala Leu Pro Arg Val Leu Leu Glu Leu Leu Gly Arg Glu Ile Leu Ser  
275 280 285

Ser Thr Lys Val Ile Phe Trp Arg Asn His Asp Gln Glu Ser Gln Cys

290                      295                      300  
 Met Glu Asn Lys Ser Arg Glu Lys Lys Lys  
 305                      310  
 <210> 214  
 <211> 320  
 <212> PRT  
 <213> Homo sapiens  
 <400> 214  
 Met His His Val Phe Ile Leu Trp Pro Leu Ile Asp Ser Trp Asp Val  
 1                      5                      10                      15  
 Lys Glu Leu Ile Leu Tyr Thr Tyr Ala Asn Leu Lys Pro Ser Ile Ile  
 20                      25                      30  
 Ser Leu Thr Ser Pro Val Ser Ser Leu Cys Leu Cys Tyr Gln Gln Val  
 35                      40                      45  
 Asn Phe Ser Val Leu Pro His His Lys Pro Gln Leu Pro Leu His Met  
 50                      55                      60  
 Phe Pro Lys Leu Val Ala Asn Ser Val Phe Pro Gly Glu Cys Ile Lys  
 65                      70                      75                      80  
 Tyr Pro Gly Ile His Cys Tyr Thr Val Ser Asn Gly Ser Ser Phe Ser  
 85                      90                      95  
 Leu Leu Trp Arg Arg Thr Pro Glu Glu Ser Thr Ser Pro Gly Pro Ala  
 100                      105                      110  
 Ala Ser Cys Met Gly Asn Leu Leu Leu Leu Leu Gly Phe Thr Leu  
 115                      120                      125  
 His Ile Leu Ser Leu Arg Lys His Thr Lys Ser Phe His Val Phe Val  
 130                      135                      140  
 Pro Val Pro Met Pro Leu Leu Pro Gly Ile Pro Phe Phe Tyr Ser Tyr  
 145                      150                      155                      160  
 Ser Leu Asn Lys Leu Phe Tyr Ser Phe Ser Ser Gly Pro Leu Pro Leu  
 165                      170                      175  
 Ile Gln Leu Arg Asn Asn Tyr Cys Leu Ser Pro Ser Lys Leu Ile Phe  
 180                      185                      190  
 Cys Leu Leu Phe Ser His His Thr Leu Pro Phe Thr Ser Val Ala Tyr  
 195                      200                      205  
 His Phe Phe Cys Tyr Leu Thr Asn Ala Ser Val Phe Ile His Ser Pro  
 210                      215                      220  
 Pro Arg Leu Tyr Ser Ser Trp Val Gln Ser Ile Ser His Ser Phe Leu  
 225                      230                      235                      240  
 Cys Tyr Leu Cys Leu Ser Gln Cys Trp Leu Gln Ser Arg Tyr Phe Arg  
 245                      250                      255  
 Asp Ala Ile Ile Arg Val Arg Val Val Arg Ile Gly Glu Asn Glu Asp  
 260                      265                      270  
 Ser Met Val Leu Arg Cys His Ala Ser Cys Lys Glu Asn Met Lys Gly  
 275                      280                      285

His Phe Phe Phe Leu Gln Leu His Gly Leu Leu Gln Ser Leu Cys Leu  
 290 295 300  
 Leu Gly Leu Glu Leu Pro Ala Ile Ser Val Phe Val Arg Leu Leu Ile  
 305 310 315 320  
 <210> 215  
 <211> 317  
 <212> PRT  
 <213> Homo sapiens  
 <400> 215  
 Pro Val Asn Ala Lys Asp Ile Leu Phe Gly Leu Glu Ile Lys Leu Leu  
 1 5 10 15  
 Met Pro Ile Trp Pro Tyr Ala Leu Arg Thr Leu Leu His Asn Lys Ile  
 20 25 30  
 Ala Val Arg Val Thr Lys Trp Lys Met Asn Asn Met Tyr Arg Glu Arg  
 35 40 45  
 Ile Gln Lys Arg Asn Leu Tyr Phe Ile Phe Ser Lys Leu Pro Gln Ile  
 50 55 60  
 Cys Leu Arg Lys Leu Tyr Asp Leu Val Asn Arg Ile Leu Lys Thr Leu  
 65 70 75 80  
 Ile Tyr Lys Ser Gln Val Trp Ala Leu Val Thr Ser Leu Asn Asp Trp  
 85 90 95  
 Leu Ala Asp Asn Leu Ser Gly Ser Ser Tyr Leu Glu Ile Glu Asn Thr  
 100 105 110  
 Ser Leu Pro Phe Tyr Asn Ser Pro Gln Leu Phe Gln His Thr Gln Cys  
 115 120 125  
 Asp Lys Lys Pro Ser Gln Ala His Phe Ser Asn Asn Glu Phe Val Gly  
 130 135 140  
 Ser Phe Lys Cys Gln Gly Gln Gln Val Arg Ala Gly Ser Glu Ala Asp  
 145 150 155 160  
 Ile Phe Gly Glu His Gly Leu Ala Phe Ser Phe Leu Gly Thr Phe Val  
 165 170 175  
 Leu Trp Met Glu Ser Ile Leu Gly Gln Ala Glu Val Leu Leu Ser Trp  
 180 185 190  
 Trp Gln Asp Gly Tyr Ala Arg Gln Pro Ser Cys Leu Gln Arg Ala Cys  
 195 200 205  
 Leu Val Arg Ser Phe Gly Ile Ser Ser Asp Leu Met Asn Leu Gly Leu  
 210 215 220  
 Met Phe Ile Pro Gly Tyr Ile Ser Phe Ala Gln Val Asn Gly Tyr Val  
 225 230 235 240  
 Asp Cys His Thr Trp Val Ser Val Thr Thr Pro Gly Phe Ser Asp Gly  
 245 250 255  
 Val Ser Pro Lys Gly Pro Thr Arg Val Glu Glu Ser Gly Ser Trp Lys  
 260 265 270

Glu Ser Gln Gly Lys Gly Lys Gly Thr Asn Ala Arg Trp Ala Val Asn  
275 280 285

Gly Ser Cys Pro Asn Phe Met Pro Glu Pro Leu Lys Gly Ile Phe Thr  
290 295 300

Leu Thr Val Gly Ile Asn Ile Gly Arg Gly Asp Ala Trp  
305 310 315

<210> 216  
<211> 319  
<212> PRT  
<213> Homo sapiens

<400> 216

Arg Lys Lys Asp Asp Ser Ile His Val Arg Arg Asn Ser Ala Arg Met  
1 5 10 15

Gln Lys His Lys Tyr Glu Lys Arg Val Tyr Cys Phe His Asn Lys Thr  
20 25 30

Lys Thr Arg Lys Glu Ile Ala Cys Gly Lys Glu Lys Gln Ser Lys Lys  
35 40 45

Arg Lys Thr Asn Leu His Val Ala Asn Leu Phe Val Thr Phe Gln Ile  
50 55 60

His Met Ser Cys Ala Met Ile Thr Arg Gly Phe Pro Asp Lys Phe Cys  
65 70 75 80

Phe Ser Ile Ile Phe Leu Gln Leu Tyr Lys His Gly Phe Tyr Ser Asp  
85 90 95

Asn Leu Ser Phe Asp Ile Phe Phe Ile Asp Tyr Gln Arg Ile Leu Glu  
100 105 110

Thr Asn Gln Ala Gln Tyr Phe Asn Phe Gln Phe Ser Leu Pro Val Ile  
115 120 125

Leu Leu Pro His Thr Ala Ser Thr Pro Ser Trp Tyr Gln Leu Lys Lys  
130 135 140

Tyr Tyr Val Arg Met Thr Ser Val Thr Leu Val Leu Phe Ile Leu Asn  
145 150 155 160

His Ser Glu Pro Tyr His Cys Val Leu Asn Leu His Leu Thr Asp Pro  
165 170 175

Tyr Leu Cys Ser Ser Ser Ala Leu Asp Leu Cys Phe Gln Ala Leu  
180 185 190

Arg Phe Tyr Asn Val Ile Asn Pro Leu Ser Leu Ile Phe Ser Ser Pro  
195 200 205

Leu Thr Cys Met Cys Val Glu Ser Val Tyr Met Leu Glu Asn Tyr Thr  
210 215 220

Thr Phe Thr Arg Phe Ile Leu Leu Val Tyr Leu Thr Leu Thr His Phe  
225 230 235 240

Tyr Ser Leu Gly His Tyr Leu Cys Met Ala Tyr Ala Glu Val Gly Ser  
245 250 255

Gly His Tyr Lys His Gln Glu Thr Ile Ser Ile Thr Pro Cys Ile His

260                      265                      270  
 Val His Val Val Leu Lys Tyr Asn Val Lys Tyr Arg Glu Val Thr Leu  
       275                      280                      285  
 Gly Leu Asn Ser Gly Val Ser Ala Arg Leu Gly Leu Ile Thr Thr Leu  
       290                      295                      300  
 Leu Leu Ala Asn Tyr Ala Ser Leu Asn Pro Cys Ala Ser Lys Leu  
       305                      310                      315  
 <210> 217  
 <211> 313  
 <212> PRT  
 <213> Homo sapiens  
 <400> 217  
 Trp Pro Gln Ile Ser Phe Pro Pro Tyr Val Pro Leu Val Ser Thr Asn  
   1                              5                              10                              15  
 Leu Phe Leu Pro Tyr Trp Ser Gly Gln Cys Pro Pro Asp Thr Ala Val  
                               20                              25                              30  
 Leu Pro Thr Gly Leu Leu Ser Ser Phe Leu Ser Val Ile Ile Leu Ala  
       35                              40                              45  
 Cys Leu Trp Leu Lys Ala His Leu Cys Gly Pro Gln Arg Asn Tyr Leu  
       50                              55                              60  
 Pro Leu His Ser Ser Ser Trp His Leu Ser Leu Met Asp Ser Tyr Tyr  
       65                              70                              75                              80  
 Pro Leu Leu Leu Leu Cys Ala Phe Met His Ile Ile Leu Ala Pro Pro  
                               85                              90                              95  
 Asp Gln Leu Ser Leu Gly Gln Gly Phe Asp Leu Val Pro Ile Tyr Ser  
       100                              105                              110  
 Ser Pro Arg Ala Ser Leu Leu His Thr Val Gly Trp Gly Lys Ile Phe  
       115                              120                              125  
 Ala Tyr Ala Asp Asp Leu Arg Lys Ile Ile Leu Gln Thr Gly Glu Val  
       130                              135                              140  
 Lys Ile Ser Leu Ser Cys Ser Ile Trp Asn Glu Leu Val Ala Gly Asn  
       145                              150                              155                              160  
 Gln Leu Glu Val Ser Ser Glu Gly Asn Thr Trp Thr Tyr Pro Leu Leu  
                               165                              170                              175  
 Gln Val Ser Tyr Leu Tyr Lys Asp Cys Val Pro Val Thr Asn Leu Phe  
       180                              185                              190  
 Leu Asn His Trp Cys Cys Tyr Leu Gln Glu Gly Leu Gly Gln Ile Cys  
       195                              200                              205  
 Glu Glu Thr Ser Met Tyr Thr His Pro Tyr His Leu Lys Asn Lys Phe  
       210                              215                              220  
 Val Cys Val Pro Leu Met Lys Tyr Glu Glu Arg Ser His Ser Phe Gln  
       225                              230                              235                              240  
 Ser Thr Gln Ala Leu Cys Leu Gly Leu Leu Ala Thr His Ala Lys Ile  
                               245                              250                              255

Leu Tyr Gln His Phe Val Lys Pro Thr Ile Leu Thr Val Pro Ala Leu  
260 265 270

Gln Pro Val Ile Asp Ser Asn Phe Asn Ser Pro Leu Val Ala Ile Ser  
275 280 285

Asp Ala Gln Cys Leu Cys Leu Leu Pro Leu Cys Ile Pro Ser Pro Ala  
290 295 300

Leu Asn Ser Ala Gly Cys Ile Gln Glu  
305 310

<210> 218  
<211> 313  
<212> PRT  
<213> Homo sapiens

<400> 218

Thr Cys Ser Ser Thr Asp Ser Lys Val Ile Leu Lys Ser Gln Leu Asn  
1 5 10 15

Val Ile Thr Arg Cys Arg Asp Ser Arg Tyr Val Tyr Ser Glu Arg Asn  
20 25 30

Cys Ser Pro Ser Val Ile Leu Ile Lys Val Lys Ser Phe Gln Asn Ala  
35 40 45

Met Val Gly Gln Thr Asn Arg His Ser His Ser Lys Arg Glu Lys Glu  
50 55 60

Gly Ile Leu Gln Gln Gln Gln Ser Lys Arg Ile Leu Arg Leu Gln Asn  
65 70 75 80

Asn Leu Leu Leu Met Pro His Leu Pro Ile Phe Gln Ala His Leu Gly  
85 90 95

Arg Arg Trp Ala Pro Lys Ala Leu Gly Val Pro Val Pro Ala His Met  
100 105 110

Thr Ala Leu Thr Tyr Ser His Met Pro Gly Trp Lys Cys Pro Leu Val  
115 120 125

Ala Leu Leu Val Tyr Gly Gln Arg Val Gly Leu Leu Leu Leu Cys Gln  
130 135 140

Ala Gln Pro Trp Arg Leu Phe Val Val Ala Pro Pro Leu Cys Gln Phe  
145 150 155 160

Phe Ala Ala Ser Arg Leu Ser Arg Ala Ser Phe Glu Ile Cys Val Glu  
165 170 175

Ser Ala Phe Pro Leu Trp Tyr Cys Thr Val Cys Pro Gly Gly Asp Asp  
180 185 190

Thr Arg Thr Leu Pro Thr Phe Ile Ile Cys Ala Leu Gln Lys Gly Gly  
195 200 205

His Trp Ser Pro His His Thr Trp Thr Leu Trp Ser His Ala Trp Asn  
210 215 220

Asp Ala Val Leu Cys Gln Lys Ala Gly Ser Arg Asp Glu Val Ala Gly  
225 230 235 240

Arg Lys Cys Ala Pro Val Gly Ile Leu Gly Pro Ser Phe Asp Leu Val  
 245 250 255  
 Leu Ser Pro Arg Pro Trp His Ala Gly Pro Val Met Gly Ala Ala Ala  
 260 265 270  
 Val Met Met Ser Glu Met Leu Leu Val Gly Val Ile Pro Pro Leu Pro  
 275 280 285  
 Lys Ala Pro Gly Phe Cys Ser Ser Met Leu Ile Ser Asn Gly Cys Trp  
 290 295 300  
 Ala Thr Ser Leu Val Phe Ser Pro Lys  
 305 310  
 <210> 219  
 <211> 318  
 <212> PRT  
 <213> Homo sapiens  
 <400> 219  
 His Arg Asn Ile Leu Gln Asn Phe Asn Ile Thr Val Leu Asn Ser Val  
 1 5 10 15  
 Lys Thr Lys Asp Asn Pro Leu His Pro Asn Met Thr Ala Phe Asn Ile  
 20 25 30  
 Leu Leu Tyr Phe Ser Leu Phe Ala Met Tyr Ile Ile Leu Gln Ser Cys  
 35 40 45  
 Asn His Thr Gln Tyr Met Ile Leu Ser Cys Phe Pro Thr Tyr His Tyr  
 50 55 60  
 Arg Tyr Phe Tyr Cys Tyr Ile Val Phe Met Val Val Ile Val Asn Ser  
 65 70 75 80  
 Tyr Ala Val Ile Val His Ile Glu Val Leu Tyr Leu Leu Ser Tyr Pro  
 85 90 95  
 Ile Ile Phe Lys Gln Phe Leu Ile Ser Phe Tyr Asn Lys His Gly His  
 100 105 110  
 Ile Ser Asp Arg Gly Val Leu Phe His Ile Leu Thr Tyr Phe Ser His  
 115 120 125  
 Ser Val Thr Ile Thr Pro Lys Asn Thr Asn Phe Leu Ser Leu Asp Val  
 130 135 140  
 Tyr Phe Gln Lys Ile Phe Lys Arg Cys Ile Asn Leu Leu Cys Ser Trp  
 145 150 155 160  
 Cys Lys Arg Pro Phe Cys His Cys Phe Leu Glu Ser Arg Ala Ser Lys  
 165 170 175  
 Ser Arg Asp Met Trp Leu Gly Gly Arg Asn Pro Ala Trp Gly Arg His  
 180 185 190  
 Ser Val Lys Asn Ser Ser Ser His Trp Tyr Thr Gly Phe Ile Phe Leu  
 195 200 205  
 Cys Phe Leu Gln Thr Glu Gln Leu Ile Thr Leu Trp Val Leu Phe Val  
 210 215 220  
 Phe Thr Ile Val Gly Asn Ser Val Val Leu Phe Ser Thr Trp Arg Arg



225                      230                      235                      240  
 Lys Lys Lys Ser Arg Met Thr Phe Phe Val Thr Gln Leu Ala Ile Thr  
                                  245                                   250                                   255  
 Gly Lys Leu Cys Lys Glu Ala Gly Ser Tyr Met Ser Pro Tyr Gly Phe  
                                  260                                   265                                   270  
 Leu Leu Leu Met Asn Phe Ile Lys Lys Lys Lys Met Arg Ile Gly Gln  
                                  275                                   280                                   285  
 Phe Gly Asn Asn Phe Lys Asn Ile Lys Pro Ile Phe Glu Tyr Phe Leu  
                                  290                                   295                                   300  
 Trp His Thr His Ile Met Pro Leu Arg Phe His Tyr Lys Ser  
 305                                   310                                   315  
  
 <210> 220  
 <211> 320  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 220  
  
 Ile Ile Pro Ser Val Ile Phe Phe Tyr Cys Arg His Cys Lys Ser Leu  
 1                                   5                                   10                                   15  
 Asn Leu Asp Lys Ser Tyr Ser Gly Gln Asn Lys Asn Phe Thr Val Ile  
                                  20                                   25                                   30  
 Asn Val Cys Ser Cys Thr Cys Glu Val Lys Ser Phe Ser Leu Leu Ser  
                                  35                                   40                                   45  
 Asn Ser Tyr Val Pro Asn Ile Phe Ser Lys Phe Leu Lys Thr Tyr Asn  
                                  50                                   55                                   60  
 Gly Glu Lys Asn Asn Pro Phe Ser Ser Pro Ala Ser Leu Met Lys Asn  
 65                                   70                                   75                                   80  
 Ser His Phe Ser Leu Phe Leu Leu Phe Leu Leu Val Val Phe His Ile  
                                  85                                   90                                   95  
 Ser Cys Leu Ser Ala Val Ser Cys Phe Met Gln Phe Arg Pro Tyr Leu  
                                  100                                   105                                   110  
 Leu Thr Ser Leu Ser Phe Gln Tyr Lys Asp Ser Cys Ile Phe Ser Phe  
                                  115                                   120                                   125  
 Asn Phe Thr Phe Leu Asn Ser Pro Phe Pro Phe Cys Asp Pro Gly Ile  
                                  130                                   135                                   140  
 Ser Gly Val Leu Phe Phe Phe Ile Leu Pro Asp Phe Ile Tyr Ile Cys  
 145                                   150                                   155                                   160  
 Val Tyr Ser Phe Leu Leu Phe Phe Lys Leu Lys Thr Cys Leu Ser Ser  
                                  165                                   170                                   175  
 Lys Ser Gly Ser Phe Phe Phe Ser Trp Arg Pro Leu Ser Gln Asn Pro  
                                  180                                   185                                   190  
 Leu Ser Phe Cys Phe Asn Glu Asp Tyr Met Leu Ser Leu Trp Leu Pro  
                                  195                                   200                                   205  
 Ser Cys His Trp Ser Ser Ser Leu Cys Cys Tyr Pro Gly Leu Lys Leu  
                                  210                                   215                                   220

Leu Phe Leu Asp Pro Ile Leu Ser Leu Ser Trp Phe Ile Thr Leu Phe  
225 230 235 240

Cys Trp Gly Thr Ser Ser Cys Met Trp Asn Val Met Ser Ala Ser Leu  
245 250 255

Cys Phe Lys Met Tyr Ile Phe Cys Pro Leu Phe Asp Leu Ala Glu Asn  
260 265 270

Arg Ile Leu Asp Cys Lys Ile Gln Lys Leu Leu Gln Arg Leu His His  
275 280 285

Arg Gln Lys Asn Leu Cys Thr His Phe Pro Pro Thr Ser Ser Pro Pro  
290 295 300

Ala Ala Arg Ser Asn His Glu Ser Phe Cys Gln Asn Arg Phe Ala Tyr  
305 310 315 320

<210> 221  
<211> 318  
<212> PRT  
<213> Homo sapiens

<400> 221

Cys Ile Lys Val Phe Ile Leu Lys Gly Lys Ala Thr Met Ile Ala Gln  
1 5 10 15

Leu Trp Tyr Ile Ile Ile Ser His Ile Ile Phe Leu Leu Leu Glu Lys  
20 25 30

Gly Ile Tyr Asp Phe Ser Arg Met His Thr Glu Lys Pro Leu Cys Ile  
35 40 45

Ile Leu Cys Glu Ser Lys Leu Cys Thr Tyr Phe Glu Val Ile Cys Ile  
50 55 60

Leu Cys Arg Arg Lys Glu Asn Asn Leu Leu Tyr Phe Val Cys Gly Ile  
65 70 75 80

Gly Asn Val Phe Leu Thr Lys Pro Lys Asn Ile Ser His Ser Lys Gly  
85 90 95

Lys Met Gly Leu Asn Glu Lys Met Val Asp Leu Lys Tyr Gly Gly Arg  
100 105 110

Phe Phe Trp Gly Thr Leu Asp Leu Ile Met Phe Phe Ser Ile Pro Phe  
115 120 125

Leu Gln Met Phe Ile Ile Leu Leu Leu Phe Ile Tyr Ala Ala Ile Ile  
130 135 140

Tyr Val Cys Ser Cys Phe Ser Cys Ser Gln Thr Leu Tyr Asn Val Ile  
145 150 155 160

Ile Gln His Glu Ser Phe Ser Ile Leu Leu Phe Leu Val Asn Ile Ile  
165 170 175

Ile Trp Gly Tyr Trp Cys Thr His Cys Gln Phe Ile His Phe Asn Tyr  
180 185 190

Ser Thr Gly Phe Trp Ser Met Asn Ile Ser Tyr Phe Ile Tyr Leu Tyr  
195 200 205

Pro Ile Asp Val Tyr Leu Val Pro Ile Phe Ala Val Lys Asn Asn Ala  
 210 215 220  
 Ala Ile Lys Pro Ser Gly Ile Cys Phe Ser Lys Cys Ile Pro Arg Ser  
 225 230 235 240  
 His Arg Phe Ser Gly Cys His Ser Leu Lys Leu Leu Gly Lys Thr Val  
 245 250 255  
 Arg Ile Leu Gly Asn Leu Leu Asn Leu Thr Trp Leu Asn Phe Leu Ala  
 260 265 270  
 Gln Met Arg Val Val Leu Asp Leu Ile Lys Asn Met Val Ile Phe Cys  
 275 280 285  
 Glu Thr Leu Ala Asn Tyr Asp Asn Lys Trp Ser Leu Gly Ile Ser Val  
 290 295 300  
 Ile Thr Ala Ile Lys Arg Gly Leu Lys Tyr Pro Lys Glu Lys  
 305 310 315

<210> 222  
 <211> 317  
 <212> PRT  
 <213> Homo sapiens

<400> 222

Asn Tyr Leu Ser Asp Cys His Ser Phe Met Glu Leu Ser Val Asn Lys  
 1 5 10 15  
 Val Leu Leu Tyr Val Asn Met Arg Leu Ile Phe Phe Leu Ser Leu Leu  
 20 25 30  
 Phe Gly Leu Tyr Phe Phe Gln Val Arg Ala Ile His Gly Ser Ala Ser  
 35 40 45  
 Thr Asp Gln His Leu Leu Ser Tyr Phe Ala Ile Trp Leu Pro Gly Leu  
 50 55 60  
 Arg Glu Cys Phe Phe Asn Leu Tyr Trp Trp His Cys Trp Leu Leu Ile  
 65 70 75 80  
 Leu Leu Phe Val Leu Ala Arg Leu Leu Phe Lys Arg Arg Val Ile Asn  
 85 90 95  
 Ser Val Leu Arg Ala Glu Val Lys Tyr Arg Met Glu Leu Glu Glu Asn  
 100 105 110  
 Glu Ala Ser Ile Ser Val Lys Lys Ser Phe Ile Lys Ala Val Gly Asp  
 115 120 125  
 Arg Glu Leu Gly Val Thr Ile Leu Val Pro Ile Val Met Val His Pro  
 130 135 140  
 Gly Lys Ile Gln Gly Lys Arg Glu Ser Leu Trp Lys Ser Phe Gly Cys  
 145 150 155 160  
 Val Leu Ser Cys Phe Arg Lys Leu Ala Asn Phe Tyr Thr Ser Val Phe  
 165 170 175  
 Arg Leu Ser Cys Leu Asp Thr His Pro Thr Gln Ser Ala Gln Gln Tyr  
 180 185 190  
 Phe Leu Cys Ser Ser Leu Ser Pro Gly Ile Arg Met Ala Pro Leu Gly

195                      200                      205  
 Glu Leu Leu Ser His Met Ile Lys Asp Leu His Tyr Phe Leu Ser Lys  
   210                      215                      220  
 Ser Arg Arg Lys Val Gly Glu Leu Ala Trp His Leu Ala Gly Thr Tyr  
   225                      230                      235                      240  
 Asn Thr Ala Ser Thr Trp His Leu Leu Asp Arg Leu Pro Leu Pro Thr  
                           245                      250                      255  
 Val Val Thr Thr Ser Met Gly Gly Gly Trp Cys Cys Thr Val Pro Met  
                           260                      265                      270  
 Gly Trp Cys Ala Cys Ser Pro Met Pro Pro Ala Leu Pro Gln Cys Cys  
                           275                      280                      285  
 Leu Leu Gln Ser His Leu Phe Arg Trp Ser Ile Leu Ile Glu Lys Val  
   290                      295                      300  
 Leu Gly Thr Ile Cys Leu Lys Cys Ser Pro Ala Asn Val  
   305                      310                      315  
 <210> 223  
 <211> 314  
 <212> PRT  
 <213> Homo sapiens  
 <400> 223  
 Leu Cys Tyr Cys Val Ile Ile Ile Ile Val Pro Phe Pro Ser Ile Pro  
   1                      5                      10                      15  
 Gln Thr His Thr Tyr Val Glu Ile Leu Arg Gly Asp Asp Val Leu Phe  
                           20                      25                      30  
 Thr Ser Ala Cys Leu Met Leu Ser Pro Val Leu Gly Thr Asn Ala Ile  
                           35                      40                      45  
 Val Phe Leu Glu His Glu Ile His Gln Lys His Glu Trp Ile Trp Trp  
   50                      55                      60  
 Gly His Lys Arg Leu Thr Pro Gly Ser Arg Asn Leu Gly Gly Glu Thr  
   65                      70                      75                      80  
 Ser Gly Leu Glu Gly Ala Glu Asp His Cys Val Arg Ser Thr Trp Phe  
                           85                      90                      95  
 Trp Leu Ala Gly Leu Ala Arg Met Gln Arg Ser Phe Trp Val Leu Leu  
                           100                      105                      110  
 Lys Phe Lys Thr Thr Ile Ile Ile Asn Ile His Leu Val Leu Thr Met  
                           115                      120                      125  
 Cys Gln Ser Leu Ile Ala Phe Tyr Val Phe Ser His Ser Ser Lys Phe  
   130                      135                      140  
 Gly Leu Asp Ile Phe Pro Val Tyr Thr Ile His Met Arg Lys Arg Val  
   145                      150                      155                      160  
 Glu Gln Gly Gly Ala Glu Thr Cys Pro Arg Ile His Ser Lys Asn Gly  
                           165                      170                      175  
 Asn Trp Asp Trp Ser Pro Arg Asp Ser Cys Phe Leu Asp Phe Val Phe  
                           180                      185                      190

Leu Ile Ser Leu Pro Leu Arg Leu Phe Ile Asp Ile Phe Thr Phe Tyr  
 195 200 205  
 Phe Glu Ile Ile Val Asp Ser Gln Glu Val Thr Arg Glu Arg Ser Cys  
 210 215 220  
 Val Leu Phe Thr Gln Ile Ser Pro Met Leu Arg Phe Tyr Ile Thr Val  
 225 230 235 240  
 Ile Gln Tyr Glu Asn Gln Glu Thr Asp Ile Gly Ser Ile Tyr Val Tyr  
 245 250 255  
 Thr Ser Met Pro Phe His His Val Met Pro Pro Ser Pro Ser Cys Arg  
 260 265 270  
 Thr Val Pro Ser Pro Arg Arg Ser Ala Thr Cys Cys Ser Phe Lys Val  
 275 280 285  
 Ile Pro Ala Leu Phe Pro Val Pro Thr His Cys His Tyr Ala Pro Leu  
 290 295 300  
 Val Thr Thr Asn Leu Phe Ser His Leu Tyr  
 305 310

<210> 224  
 <211> 321  
 <212> PRT  
 <213> Homo sapiens

<400> 224

Lys Pro Ser Ser Gly Cys Gly Gly Trp Met Trp Asp Trp Met Gly Thr  
 1 5 10 15  
 Gln Lys Asn Ile Lys Thr Met Ala Thr Val Ile Ile Ile Val Ile Asn  
 20 25 30  
 Ser Gln Asp Asn Asn His Leu Ala Thr Val Ala Met Tyr Leu Lys Asp  
 35 40 45  
 Tyr Ser Leu Gly Val Phe Phe Leu Met Ser Met Glu Gln Asp Asp Trp  
 50 55 60  
 Ala Phe Glu Asp Ile Lys Glu Thr Lys Gly Pro Asp Cys Asn Gln Arg  
 65 70 75 80  
 Phe His Ser His Arg Pro Gly Phe Thr Trp Gln His Thr Phe Trp Thr  
 85 90 95  
 Phe Phe Phe Phe Ser Gly Lys Glu Thr Gly Ser Val Glu Asn Gly Arg  
 100 105 110  
 Met Arg Thr Asn Cys Arg Ala Leu Pro His Ser Trp Thr Leu Ser His  
 115 120 125  
 Ser Ser Arg Trp Gly Pro Pro Ala His Cys Trp Leu Cys Pro Pro Gln  
 130 135 140  
 Phe Leu Arg Ile His Thr Asp Phe Ala Lys Ile Leu Arg Tyr Val Gly  
 145 150 155 160  
 His Glu Leu Trp Val Cys Ala His Leu Val Pro Ser Leu Tyr Ser Thr  
 165 170 175

Leu His Ser Ser Gly Val Phe Leu Thr Ala Gly Ala Thr Phe His Leu  
 180 185 190  
 His His Tyr Tyr Ile Lys Trp Ala Ser Ile Phe Pro Ser Glu Phe Gln  
 195 200 205  
 Pro Leu Ser Gly Asn Leu Thr Phe Phe Leu Val Ser Phe Ala Leu Arg  
 210 215 220  
 Phe Cys Pro Phe Tyr Cys Ser Asn Glu Phe Thr Gln Pro Ser Ile Pro  
 225 230 235 240  
 His Glu Ser Gly Gln Asp Pro Val Thr Cys Asp Ser His Thr Asp Cys  
 245 250 255  
 Val Arg Val Thr Pro Pro Val Pro Gly Phe Pro Glu Pro Cys Leu Ser  
 260 265 270  
 Arg Leu Thr Gly Gln Ser Trp Asp Met Asn Trp Ala Pro Glu Leu Ala  
 275 280 285  
 Leu Phe Val Ser Arg Ser Ser Arg Cys Leu Cys Arg Leu Pro Asn Pro  
 290 295 300  
 Cys Ser Trp Ala Trp Val Ala Glu Ser Ala Gly Arg Leu Trp Cys Met  
 305 310 315 320  
 His

<210> 225  
 <211> 314  
 <212> PRT  
 <213> Homo sapiens

<400> 225

Leu Cys Tyr Cys Val Ile Ile Ile Ile Val Pro Phe Pro Ser Ile Pro  
 1 5 10 15  
 Gln Thr His Thr Tyr Val Glu Ile Leu Arg Gly Asp Asp Val Leu Phe  
 20 25 30  
 Thr Ser Ala Cys Leu Met Leu Ser Pro Val Leu Gly Thr Asn Ala Ile  
 35 40 45  
 Val Phe Leu Glu His Glu Ile His Gln Lys His Glu Trp Ile Trp Trp  
 50 55 60  
 Gly His Lys Arg Leu Thr Pro Gly Ser Arg Asn Leu Gly Gly Glu Thr  
 65 70 75 80  
 Ser Gly Leu Glu Gly Ala Glu Asp His Cys Val Arg Ser Thr Trp Phe  
 85 90 95  
 Trp Leu Ala Gly Leu Ala Arg Met Gln Arg Ser Phe Trp Val Leu Leu  
 100 105 110  
 Lys Phe Lys Thr Thr Ile Ile Ile Asn Ile His Leu Val Leu Thr Met  
 115 120 125  
 Cys Gln Ser Leu Ile Ala Phe Tyr Val Phe Ser His Ser Ser Lys Phe  
 130 135 140  
 Gly Leu Asp Ile Phe Pro Val Tyr Thr Ile His Met Arg Lys Arg Val

```
<210> 226
<211> 312
<212> PRT
<213> Homo sapiens
```

<400> 226

Gly 1	Ala	Arg	Gly	Gly 5	Glu	Ala	Ser	Thr	Ser 10	Leu	Glu	Ser	Gln	Val 15	Glu
Asp	Thr	Ala	Glu 20	Gln	Thr	Ser	Asn 25	Leu	Ile	Thr	Val	Thr	Leu 30	Ile	His
Pro	Gln	Leu 35	Ala	Lys	Tyr	Thr	Leu 40	Ile	Val	Asn	Phe	Leu 45	Pro	Leu	Trp
Ser 50	Leu	Ser	Asp	Ile	Ser	Thr 55	Asp	Leu	Leu	Phe	Ile 60	Leu	Leu	Arg	Leu
Arg 65	Asn	Ile	Ile	Arg	Ile 70	Leu	Gln	His	Leu	Gly 75	Glu	Ile	Ile	Glu	Ser 80
Ala	Met	Val	Ser 85	Phe	Ala	Asp	Ile	Tyr	Ser 90	Trp	Ser	Lys	Trp	Asn 95	Thr
Asn	Gln	Asn	Trp 100	Leu	Pro	Tyr	Ile	Leu 105	Gln	Arg	Pro	Thr	Gly 110	Gly	Lys
Gly	Leu	Trp 115	Lys	Val	Cys	Phe	Ala 120	Thr	Arg	Gln	Ile	Leu 125	Asp	His	Pro
Val 130	Ser	Gly	Ser	Ile	His	Ser 135	Phe	Pro	Asp	Ser	Pro 140	Asp	Asp	Ile	Pro

Pro Ser Phe Thr Tyr Ile Asn Ser Thr Val Pro Ile Cys Tyr Ile Ala  
 145 150 155 160  
 Ser Phe Leu Leu Phe Ile Ile Cys Leu Pro His Gln Asn Ala Ser Ser  
 165 170 175  
 Ile Trp Ala Val Ala Thr Leu Phe Thr Val Tyr Leu Ser Val Ser Met  
 180 185 190  
 Lys Ser Asp Ile Met Pro Gly Ile Tyr Tyr Glu Leu Asn Asn Tyr Val  
 195 200 205  
 Asn Glu Ile Met Arg Lys Ser Cys Leu Ile Thr Cys Gln Pro Tyr Asn  
 210 215 220  
 Ala Ser Gln Phe Phe Pro Leu Gln Phe Leu His Leu Asn Trp Ile Thr  
 225 230 235 240  
 Gln Met Leu Thr Leu Trp His Cys Trp Asn Asn Tyr Leu Lys Ser Cys  
 245 250 255  
 Lys Phe Ile Ala Tyr Trp Lys Cys Gly Ser Glu Cys Asp Thr Pro Gln  
 260 265 270  
 Tyr Gly Val Leu Val Val Leu Thr Glu Gly Asn Lys Ser Phe Arg Asn  
 275 280 285  
 Lys Val Phe Leu Ala Phe Ser His Leu Ser Phe Ser Cys Ser Pro Phe  
 290 295 300  
 Phe Pro Lys Ala Asp Gln Arg Asn  
 305 310

<210> 227  
 <211> 321  
 <212> PRT  
 <213> Homo sapiens

<400> 227

Gly Cys Ser Pro Glu Asp Asp Leu Gly Cys Ser Gly Val Asn Tyr Pro  
 1 5 10 15  
 His Phe Leu Arg Ala Ser Met Trp His Ser Trp Pro Trp Ala Ser Ala  
 20 25 30  
 Cys Pro Ala Asn Ala Gln Pro Val Pro Ala Val Pro Pro Pro Leu Ala  
 35 40 45  
 Ala Gln Pro Gln Val Trp Pro Ser Gly Leu Tyr Pro Arg Pro Pro His  
 50 55 60  
 Leu Pro Thr Leu Phe Leu Cys Ser Glu Leu Ser Thr Ala Ala Pro Ala  
 65 70 75 80  
 Pro Trp Leu Pro Leu Ile Leu Cys Leu Val Ser Phe Phe Gly His Ser  
 85 90 95  
 Phe Ala Ala Thr Leu Tyr Trp Ile Thr Leu Leu Gly Val Leu Ile Ile  
 100 105 110  
 Ser His Pro Leu Leu Leu Pro Asn Gly Pro Ser Thr Ile Ser Phe His  
 115 120 125



Arg Leu Asn Gly Lys Gly Gly Val His Ile His Arg Ile Lys Gln Val  
 130 135 140  
 Met Pro Leu His Ser Gly Val Cys Asp Asp Asn Phe Tyr Ala Phe Tyr  
 145 150 155 160  
 Thr Asn Ile Phe Val Ser Leu Cys Phe Leu Pro Cys Leu Arg Ala Leu  
 165 170 175  
 Gln Gly Leu Ala Leu Gly His Pro Val Leu His Thr His Thr Arg Thr  
 180 185 190  
 His Thr Arg Thr Cys Thr His Val His Thr His Ala His Thr His Thr  
 195 200 205  
 His Thr His Lys His Thr His Ser Leu Ala Leu Ala Asn Ala Ser Leu  
 210 215 220  
 Ala Leu Thr Thr Asn Val Ser Ala Ser Asp Leu His Asn Leu Ile Trp  
 225 230 235 240  
 Leu Phe Leu Phe Leu Gly Val Ile Cys Leu Pro Glu Gly Arg Ala Asn  
 245 250 255  
 Ser Pro Ala Ile Pro Ala Ala Tyr Ser Leu Pro Val Pro Ser Phe Pro  
 260 265 270  
 Arg Arg Gln Gln Thr Glu Arg Gly Lys Arg Tyr Lys Glu Ala Trp Gly  
 275 280 285  
 Trp Gly Lys Glu Ser Ser Tyr Leu Thr Ser Ala Pro Leu Thr Leu Leu  
 290 295 300  
 Gly Glu Val Pro Thr His Ser Ser Gly Met Thr Thr Arg Met Val Ser  
 305 310 315 320  
 Leu

<210> 228  
 <211> 123  
 <212> PRT  
 <213> Homo sapiens

<400> 228

Asp Cys Ala Ala Ala Leu Pro Gly Gln Ser Lys Thr Pro Phe Gln Lys  
 1 5 10 15  
 Lys Lys Lys Lys Lys Lys Glu Arg Lys Glu Phe Met Asp Val Ile Val  
 20 25 30  
 Lys Gly Leu Val Pro Ser Pro Ile Ser Cys Phe Pro Ser Cys His Val  
 35 40 45  
 Thr Cys Trp Phe Pro Phe Thr Phe Cys His Asp Trp Lys Leu Pro Gly  
 50 55 60  
 Ala Ser Pro Glu Ala Lys Gln Met Pro Gly Pro Cys Phe Leu Tyr Ser  
 65 70 75 80  
 Leu Leu Asn Pro Glu Pro Asn Lys Pro Leu Phe Ile Thr Asn Tyr Leu  
 85 90 95  
 Gly Ser Asp Ser Pro Leu Gln Cys Lys Trp Thr Asn Thr Pro His Asp

100                      105                      110  
 Leu His Pro Gln Thr Thr Gly Gly Thr Gln His  
     115                      120  
 <210> 229  
 <211> 210  
 <212> PRT  
 <213> Homo sapiens  
 <400> 229  
 Ser Ala Cys Gly Gly Phe Asn Gly Leu His Phe Tyr Ser Asn Ile Ser  
     1                      5                      10                      15  
 His Gln Leu Tyr Ile Tyr Tyr Leu Lys Val Phe Leu Phe Ile Val Phe  
                     20                      25                      30  
 Gln Phe Ile Phe Gln Ile Arg Ser Lys Gln Asn Tyr Ser Trp Arg Leu  
                     35                      40                      45  
 Cys Cys Leu His Pro Gln Tyr Gln Met Phe Met Ala Ser Thr Glu Pro  
     50                      55                      60  
 Gly Val Ser Met Glu Ser Leu Arg Asp Cys Leu Ser Phe Ser Glu Glu  
     65                      70                      75                      80  
 Ser Val Met Phe Ser Ile Pro Glu Glu Ala Glu Ile Thr Leu His Tyr  
                     85                      90                      95  
 Phe Phe Glu Leu Cys Ala Gly Arg His Gly Ser Glu Ile Cys Leu Ser  
                     100                      105                      110  
 Asp Ser Asn Ser Ser Ser Ile Cys Val Leu Val Phe Val Val Ala Phe  
                     115                      120                      125  
 Cys Ile Gln Leu Pro Asp Asn Phe Phe Leu Met Phe Cys Cys Asn Leu  
     130                      135                      140  
 Val Lys Leu Leu Phe Tyr Lys Leu Met Phe Trp Tyr Phe Gly His Gln  
     145                      150                      155                      160  
 Ile Leu Ala Arg Gly Lys Ile Arg Thr Arg Ser Thr Ser Cys Lys Thr  
                     165                      170                      175  
 Lys Leu Ile Phe Leu Val Asp Phe Trp Asn Gly Leu Phe Cys Phe Pro  
                     180                      185                      190  
 Ile Cys Val Tyr Phe Leu Lys Ser Cys Arg Cys Ile Tyr Glu Tyr Leu  
                     195                      200                      205

Phe His  
     210

<210> 230  
 <211> 204  
 <212> PRT  
 <213> Homo sapiens

<400> 230

Val Ile Asn Ser Ser Cys Pro Ser Ile Ile Gly Leu Gly Thr Pro Gly  
     1                      5                      10                      15  
 Phe Ser Cys Ser Ser Ser Val Ile Gly Arg Lys Ile Gly His Trp Leu

20 25 30  
 Lys Gln Ile Leu Ser Phe Leu Gly Val Val Phe Thr Leu Lys Ala Leu  
 35 40 45  
 Arg Pro Leu Gly Gly Ser Ala Ile Leu Gln His Gly Arg Cys Pro His  
 50 55 60  
 Thr Trp Met Ala Ala Phe Tyr Tyr Tyr Ser Leu Asp Thr Gly Phe Phe  
 65 70 75 80  
 Ala His Val Tyr Thr Leu Gly Ser Ile Cys Tyr Pro Phe Phe Thr Leu  
 85 90 95  
 Lys Gln Val Ile Gly Lys Phe Ile Ser Ile Trp Lys Thr Asn Asp Gln  
 100 105 110  
 Lys Asn Pro Ser Asn Pro Lys Phe Thr Glu Ala Arg Leu Leu Lys Arg  
 115 120 125  
 Lys Asp Ile Phe Leu Cys Arg Lys Val Met Phe His Arg Gly Phe Cys  
 130 135 140  
 Asn Ala Leu Thr Leu Asp Arg Ser Pro Pro Ser Ile Leu Gly Ile Thr  
 145 150 155 160  
 Ser Phe His Phe Ser Cys Lys His Ser Ser Pro Cys Thr Leu Gln Asp  
 165 170 175  
 Phe Ser Leu Phe Glu Ile Gly Leu His Ser Val Gly Arg Gly Asp Trp  
 180 185 190  
 Phe Gln Lys Glu Gly Ala Ala Gly Arg Asp Phe Ala  
 195 200

<210> 231  
 <211> 186  
 <212> PRT  
 <213> Homo sapiens

<400> 231

Gln Gly Arg Cys Thr Pro Pro Val Ile Leu Gly Val Ile Ser Ser Pro  
 1 5 10 15  
 Pro Leu Asp Ile Arg Asn Asn Ile Thr Ala Gly Val Gly Val Val Tyr  
 20 25 30  
 Ser Leu Cys Asn Ile Gly Ser Asn Ile Ile Leu Ser Pro His Trp Ile  
 35 40 45  
 Leu Gly Thr Ile Ser Gln Glu Val Trp Thr Pro Pro Ala Ile Leu Gly  
 50 55 60  
 Val Thr Ser Phe Ser Phe Pro Ser Gly Tyr Glu Gln Tyr Cys Ile Gly  
 65 70 75 80  
 Val Tyr Thr Pro Ser Asp Ile Arg Ser Asn Ile Ile Leu Ser His Ser  
 85 90 95  
 Gly Tyr Glu Gln Tyr Leu Arg Arg Ser Val Glu Pro Leu Arg Tyr Glu  
 100 105 110  
 Tyr His Pro Leu Pro Pro Trp Ile Leu Gly Thr Ile Thr Gln Gly Glu  
 115 120 125

Tyr Thr Ala Pro Val Ile Leu Arg Val Ile Ser Ser Pro His Leu Asn  
 130 135 140  
 Ile Arg Asn Asn Ile Arg Gly Val Gly Tyr Thr Ile Cys Asp Ser Gly  
 145 150 155 160  
 Arg Asn Ile Ile Leu Ser Pro Pro Gly Tyr Glu Gln Tyr His Lys Trp  
 165 170 175  
 Ser Ile His Pro Leu Arg Tyr Trp Glu Tyr  
 180 185

<210> 232  
 <211> 157  
 <212> PRT  
 <213> Homo sapiens

<400> 232

Asp Asn Leu Cys Ser Pro Cys Ser Ser Thr Pro His Ile Pro Ile Val  
 1 5 10 15  
 Cys Pro Phe His Ser Ala Pro Phe Ser Val Gln Thr Glu Leu Phe Thr  
 20 25 30  
 Asn His Tyr Pro Leu Leu Glu Met Glu Gly Ala Pro Phe Pro Thr Pro  
 35 40 45  
 Pro Leu Pro Pro Gln Leu Ser Ser Pro Arg Arg Leu Ser Ile Asn Arg  
 50 55 60  
 Leu Thr Ile Ser Leu Asn Phe His Ile Phe Val Trp Leu Ser Tyr Leu  
 65 70 75 80  
 Phe Thr Phe Ile Asn Leu Leu Cys Phe Ser Leu Val Asn Gln Ser Phe  
 85 90 95  
 Phe Ile Gly Val Ser Ala Val Ser Leu Tyr Asp Gly Glu Glu Lys Asn  
 100 105 110  
 His Pro Leu Ser Thr Pro Thr Ser Asp Arg Ser Gln Asp Ile Pro Leu  
 115 120 125  
 Lys Phe Gly Lys Val Asn Thr Ser Thr Pro Cys Ile Leu Pro Asp Asn  
 130 135 140  
 Thr Lys Asn Phe Ile Gln Tyr Ile Tyr Tyr Met Ile Lys  
 145 150 155

<210> 233  
 <211> 178  
 <212> PRT  
 <213> Homo sapiens

<400> 233

Arg Ser Arg Lys Val Asn Trp Pro Lys Val Gly Ile Tyr Ile Pro Val  
 1 5 10 15  
 Leu Leu Leu Glu Cys Cys Leu Phe Leu Asn His Pro Trp Ser Arg Pro  
 20 25 30  
 Thr Pro Ser Cys Thr Tyr Thr Asn Pro Ile Leu Ser Gln Thr Gly Leu  
 35 40 45

Trp Leu Asp Ile Gly Glu Lys Gln Leu Asp Gly Leu Thr Pro Lys Lys  
 50 55 60  
 Asn Pro Ala Arg Asp Gly Gln Asn Phe Arg Gly Gly Leu Arg Tyr Arg  
 65 70 75 80  
 Pro Cys Leu Leu Leu Ser Ser Pro Ser Cys Arg Glu Pro Arg Phe Ile  
 85 90 95  
 His Asn Lys Ile Pro His Ile His His Pro Ser Ile Tyr Ser Cys Asn  
 100 105 110  
 Leu Ile Phe Pro Gly Trp Trp Thr Arg Ala Arg Glu Pro Gln Val Glu  
 115 120 125  
 Ile Gln Lys Ala Val Thr Leu Ala Leu Cys Pro Cys Trp Arg Arg Ala  
 130 135 140  
 Ala Ala Ser His Arg Gly Arg Gly Pro Thr Glu Leu Leu Thr Leu Lys  
 145 150 155 160  
 Pro Ser Ala Asp Gly Arg Ala Lys Thr Ala Leu Glu His Ala Leu Trp  
 165 170 175

Gly Phe

<210> 234  
 <211> 188  
 <212> PRT  
 <213> Homo sapiens

<400> 234

Ile Glu Thr Lys Leu Asn Thr Phe Ala Lys Leu Leu Arg Ser Lys Phe  
 1 5 10 15  
 Leu Val Pro Arg Leu Glu Leu Pro Asn Ala Asp Lys Ser Ser Pro Val  
 20 25 30  
 Gly Ser Pro Thr Leu Phe Lys Gln Phe Leu Asp Phe Ala Pro Val Glu  
 35 40 45  
 Ala Asp Met Leu Asn His Lys Thr Pro Leu Leu Leu Ala Leu Ala Tyr  
 50 55 60  
 Cys Phe Gly Arg Ser His Phe Ser Lys Ile Arg Ala Ser Leu Ile Asn  
 65 70 75 80  
 Thr Gly Ile Arg Phe Leu Ser Gly Val Gly Ile Pro Glu Asp Arg Ile  
 85 90 95  
 Ile Tyr Phe Ala Leu Ser Arg Cys Val Met Arg Thr Glu Ala Met Leu  
 100 105 110  
 Ile Arg Asp Pro Trp Glu Leu Val Ile Tyr Tyr Leu Leu Phe Leu Pro  
 115 120 125  
 Lys Ile Asp Leu Met Glu Arg Gly Cys Ile Ile Tyr Pro Leu Ser Lys  
 130 135 140  
 Glu Ala Phe Pro Asn Thr Thr Glu Ala Val Ile Leu Lys Thr Ala Leu  
 145 150 155 160

Trp Leu Cys Ser Gln Leu Tyr Phe Leu Pro Phe His Asn Phe Leu Pro  
 165 170 175

Ser Ala Met Glu Leu Met Gly His Thr His Ile His  
 180 185

<210> 235  
 <211> 165  
 <212> PRT  
 <213> Homo sapiens

<400> 235

Lys Lys Lys Thr Pro Met Ile Trp Ile Leu Leu Ser Phe Leu Phe Ser  
 1 5 10 15

Gln Met Val Ile Leu Lys Leu Ile Glu Val Val Tyr Arg Val His Ser  
 20 25 30

His Thr Val Arg Lys Arg Gln Ser Gln Gly Leu Asn Ser Ser Ser Leu  
 35 40 45

Thr Ile Glu Pro Ile Phe Leu Ile Thr Ile Gln Tyr Phe Thr Ile Cys  
 50 55 60

Ser Ile Lys Arg Asn His Phe Ser Glu Trp Arg Asn Ile His Glu Asn  
 65 70 75 80

Lys Ser Ile Ile Gln Asp Thr Cys Lys Ala Ser Arg His Ser Arg Phe  
 85 90 95

Arg Leu Leu Ala Pro Trp Pro Arg Leu Ile Thr Phe Gln Glu Asn Lys  
 100 105 110

Thr Thr Tyr Gln Asp His Thr Ser Arg Asn Asp Leu Arg Ile Met Gly  
 115 120 125

Thr Ala Ile Trp Val Ser Asn Gly Leu Glu Ser Asp Lys Trp Phe Leu  
 130 135 140

Asn Arg Phe Pro Glu Trp Gly Asn Leu Val Leu His Gln Ala Thr Tyr  
 145 150 155 160

Val Ile Phe Ile Leu  
 165

<210> 236  
 <211> 218  
 <212> PRT  
 <213> Homo sapiens

<400> 236

Ser Phe Leu Ser Phe Asn Arg Val Glu Lys Ile Ile Ile Ser Trp Glu  
 1 5 10 15

Pro Ser Phe Phe Tyr Tyr His Glu Cys Lys Cys Thr Ser Met Thr His  
 20 25 30

Leu Pro Leu Arg Ile Lys Leu Gln Tyr Lys Lys Tyr His Tyr Thr Tyr  
 35 40 45

Leu Ser Leu Ser Phe Asn Cys Leu Leu Glu Pro Ile Leu Phe Cys Leu  
 50 55 60

Pro Arg Thr Ser Thr Met Asp Tyr Pro Phe Thr Ile Ala Leu Ser Phe  
 65 70 75 80  
 Ser Ser Phe Cys Ile Cys Phe Pro Leu Ile Phe Lys His Asp Val Ile  
 85 90 95  
 Phe Ile Arg Asp Ile Asn Ile Leu Ile Thr Trp Phe Thr Arg Thr Thr  
 100 105 110  
 Pro Ser Ser Val Val Trp Arg Thr Lys Leu Leu Glu Arg Asp Val Gln  
 115 120 125  
 Thr Gln Tyr Leu Tyr Phe Cys Met Pro His Lys Ser Ser Leu Ile Phe  
 130 135 140  
 Ile Leu Ile Ser Leu Leu Lys Asp Val Thr Lys Asp Thr Asn Glu Phe  
 145 150 155 160  
 Gln Lys Ser Pro Asn Pro Met Glu Ile His Phe Pro Leu Ser Leu Ser  
 165 170 175  
 Ser Asn Ile Leu Pro Leu Val Phe Gln Asp Ser Phe Leu Leu Ser Phe  
 180 185 190  
 Leu Leu Thr Leu Phe Ser Ser Leu Lys Ile His Pro Pro Leu Pro Ser  
 195 200 205  
 His Lys Met Leu Arg Val Glu Gly Gly Ser  
 210 215

<210> 237  
 <211> 139  
 <212> PRT  
 <213> Homo sapiens

<400> 237

Thr Gln Cys Gln Phe Thr Lys Tyr Thr Ile Ile Tyr Ser Gln Asn Thr  
 1 5 10 15  
 Phe Ile Lys Arg Asn Phe Phe Lys Arg Arg Ser Cys Gln Cys Gln Tyr  
 20 25 30  
 Arg Asn Tyr Lys Asn Pro Phe Leu Phe Pro Leu Glu Ile Pro Ser Leu  
 35 40 45  
 Asp Cys Cys Ser Lys Asn Leu Ile Ser Lys Val Val Ser Leu Ser Leu  
 50 55 60  
 Asp Asn Asp Ile Arg Lys Cys Ser Arg Gln Ile Phe Ser Lys Ile Gln  
 65 70 75 80  
 Ser Ile Trp Tyr Leu Pro Lys Ser Lys Leu Gln Arg Glu Pro Glu Cys  
 85 90 95  
 Ser Pro Thr Ala Phe Ser Ser Ser Thr Gln Trp Ile Ser Tyr Met Leu  
 100 105 110  
 Asn Cys His Val Cys Ala Ser Leu Lys Cys Ala Phe Leu Phe Thr Glu  
 115 120 125  
 Met Arg Asp Val Leu Phe Met Ile Phe Ser Leu  
 130 135

<210> 238

<211> 213  
 <212> PRT  
 <213> Homo sapiens

<400> 238

Phe Gln Tyr Phe Val Thr Cys Arg Ser Lys Trp Trp His Ala Ser His  
 1 5 10 15  
 Leu Val Asn Ser Arg Ser Cys Cys Val Ser Asn Gly Asp Thr Leu Trp  
 20 25 30  
 Leu Leu Gln Met Val Thr Leu Pro Asn Cys Phe Pro Lys Arg His Val  
 35 40 45  
 Ala Phe Phe Ser Gln Ser Leu Ile Leu Thr Leu Met Val Ile Leu Leu  
 50 55 60  
 Tyr Phe Tyr Met His Leu Val Thr Cys Leu Ile Val Ile Phe Leu Glu  
 65 70 75 80  
 Ile Gln Phe Leu Leu His Arg Val Ser Phe Glu Ile Lys Glu Arg Glu  
 85 90 95  
 Val Ala Asn Leu Gly Cys Asn Asn Phe His Leu Lys Val Asp Pro Cys  
 100 105 110  
 Phe Tyr Tyr Pro Ile Ile Asn Val Phe Cys Phe Pro Leu Ser Ala Ser  
 115 120 125  
 Tyr Cys Ser Phe Asp Ser Tyr Cys Gln Thr Glu Leu Ser Cys Phe Leu  
 130 135 140  
 Ala Arg Lys Glu Thr Thr Met Asn Glu Pro Leu Asp Tyr Leu Ala Asn  
 145 150 155 160  
 Ala Ser Asp Phe Pro Asp Tyr Ala Ala Ala Phe Gly Asn Cys Thr Asp  
 165 170 175  
 Glu Asn Ile Pro Leu Lys Met His Tyr Leu Pro Val Ile Tyr Gly Ile  
 180 185 190  
 Ile Phe Leu Val Gly Phe Pro Gly Asn Ala Val Val Ile Ser Thr Tyr  
 195 200 205  
 Ile Phe Lys Met Arg  
 210

<210> 239  
 <211> 168  
 <212> PRT  
 <213> Homo sapiens

<400> 239

Trp Phe Thr Tyr Pro Leu Asn Lys Gln Leu Leu Arg Ile Pro Ala Pro  
 1 5 10 15  
 Ala Gln Arg Gln Tyr Trp Gly Leu Cys Leu Arg Met Trp Ala Leu Glu  
 20 25 30  
 Leu Cys Gly Trp Gly Ser Asn Ser Gly Arg Ala Ala Val Arg Pro Trp  
 35 40 45  
 Thr Ser Gly Ser Ser Lys Thr Asp Arg Gln Phe Ile Phe Ile Leu Val



50                      55                      60  
 Pro Gln Ile Val Val Leu Leu Ser Asn Tyr Leu Gly Phe Ile Pro Arg  
 65                      70                      75                      80  
 His Trp Glu Ser Lys Leu Phe Ser Phe Ser Cys Leu Gln Lys Ser Ser  
                     85                      90                      95  
 Leu Thr Ile His Val Ala Tyr His Trp Ile Gly Leu His Ile Lys His  
                     100                      105                      110  
 Phe Val Thr Thr Phe Ala Cys Gly Tyr Ile Leu Leu Ser Phe Ser Tyr  
                     115                      120                      125  
 Phe Leu Leu Ala Leu Leu Glu Tyr Ser His Lys Ser Leu Ser Ser His  
                     130                      135                      140  
 Phe Trp Pro Pro Phe Asp Ser Phe Ser Leu Leu Cys Cys Cys Glu Ser  
 145                      150                      155                      160  
 Phe His Val Gln Asp Ser Arg Trp  
                     165  
 <210> 240  
 <211> 185  
 <212> PRT  
 <213> Homo sapiens  
 <400> 240  
 Ser Thr Met Cys Ile Phe Phe Trp Ala Lys Met Arg Gln Arg Cys His  
 1                      5                      10                      15  
 Val Asn Phe Ser Phe Leu His Thr Thr Ile Val Ser His Lys Thr Lys  
                     20                      25                      30  
 Asn Lys Arg Lys His Met Phe Thr Val Gly Arg Ile Ile Thr Arg Ser  
                     35                      40                      45  
 Ser Val Ala Trp Pro Lys Glu Pro Leu Pro Thr Tyr Trp Gly Cys His  
                     50                      55                      60  
 Met Lys Gly Phe Ser Lys Arg Leu Ala Ile Phe Ile Lys Gly Val Arg  
 65                      70                      75                      80  
 His Gly Ser Gly Gln Gln Thr Ser Leu Trp Lys Gly Ser Lys Leu Leu  
                     85                      90                      95  
 Gln Gln Asn Glu Arg Ile Met Val His Leu Pro Thr Leu Cys Asn Leu  
                     100                      105                      110  
 Trp Met Lys Pro Gln Pro Arg Lys Val Lys Leu Leu Cys Val Cys Val  
                     115                      120                      125  
 Trp Gly Cys Glu Gly Arg His Arg Lys Gly Lys Ala Asp Arg Pro Trp  
                     130                      135                      140  
 Lys Thr Asp Ile Ser Pro Gly Glu Trp Asn Gly Gln Ser His Asn Thr  
 145                      150                      155                      160  
 His Val Leu Asn Ile Thr Cys Phe Arg Lys Tyr Asn Ile Lys Thr Leu  
                     165                      170                      175  
 Phe Lys Ser Tyr Ser Leu Met Ile Ser  
                     180                      185

<210> 241  
 <211> 196  
 <212> PRT  
 <213> Homo sapiens

<400> 241

Val Leu Asp Ile Asp Val Arg Met Gly Gly Leu Ser Tyr Pro Ser Pro  
 1 5 10 15  
 His Val Phe Leu Leu Arg Asp Ser Asn Cys Asn Thr Ser Leu Val Phe  
 20 25 30  
 Phe Ala Ser Ser Leu Ile Pro Tyr Gln Gly Lys Ser Ser Glu Leu Ser  
 35 40 45  
 Asn Glu Ile Trp Lys Glu Lys Val Ser Lys Tyr Thr Gln His Tyr Ser  
 50 55 60  
 Thr Ser Phe Ser Leu Gly Leu Ala Ser Leu Gln Arg Glu Tyr Ile Leu  
 65 70 75 80  
 Leu Cys Ala Gly Ser Phe Pro Lys Leu Ile Ser Gly Phe Val Asn His  
 85 90 95  
 Gly Thr Ile Asp Ile Leu Asp Gln Ile Ile Leu Cys Cys Met Ala Cys  
 100 105 110  
 Ser Val Phe Cys Gln Ile Phe Gly Ile Ile Pro Gly Leu Asn Leu Pro  
 115 120 125  
 Asp Ala Asn Ser Thr Phe Ser Leu Lys Thr Ile Glu Ile Phe Gln Asp  
 130 135 140  
 Val Ala Lys Cys Pro Ser Gly Leu Lys Val Ala Pro Asn Ser Asn His  
 145 150 155 160  
 Cys Phe Glu Ala Cys His His Arg Glu Gly Cys Leu Arg Leu Asn Val  
 165 170 175  
 Cys Leu Arg Leu Ile Tyr Thr Pro Lys Ser Asn Ser Thr Val Thr Leu  
 180 185 190  
 Ile Ser Arg Lys  
 195

<210> 242  
 <211> 198  
 <212> PRT  
 <213> Homo sapiens

<400> 242

Phe Ala Leu Phe Pro Met Phe Ile Ile Ser Leu Asn Gly Thr Pro Ile  
 1 5 10 15  
 Cys Met Val Ala Trp Glu Ile Tyr Gly Ile Ile Leu Glu Pro Ser Phe  
 20 25 30  
 Phe Ile Ile Pro Met Ser Arg Ser Glu Ile Leu Ser Glu Tyr Ala Ser  
 35 40 45  
 Leu Ile Tyr Leu Lys Leu Ala His Phe Lys Phe Leu Ser Ile Leu Thr  
 50 55 60

Leu Leu Tyr Leu Asn Asp Tyr His Ser Pro Asn Cys Phe Leu Met Gly  
 65 70 75 80  
 Leu Ile Gly Lys Thr Asn Leu Phe Leu Ile Leu Pro Leu Glu Leu Ser  
 85 90 95  
 Phe Gln Thr Arg Met Trp Pro Ser Phe Phe Leu Thr Asn Asp Leu Ile  
 100 105 110  
 Val Pro Lys Thr Lys Ser Ile Leu Ser Leu Asn Asn Ile Gln Gly Pro  
 115 120 125  
 His Ser Arg Ser Ser Leu Ile Pro Thr Ser Val Phe Leu Ser Ser Ser  
 130 135 140  
 Pro Ser Gln Ser Thr Leu Ser His Thr Arg Tyr Ser Thr Trp Ser His  
 145 150 155 160  
 Ile Lys Leu Leu Ser Ile Leu Gly Phe Leu Leu Ala Phe Asn Pro Leu  
 165 170 175  
 Leu Gly Trp Cys Ile Pro Gly Glu Trp Ser Asn Pro Cys Thr Cys Tyr  
 180 185 190  
 His Ala Pro Thr Phe Leu  
 195

<210> 243  
 <211> 180  
 <212> PRT  
 <213> Homo sapiens

<400> 243

Leu Cys Asp Gly Val Met Arg Trp Gly Arg Arg Val Trp His His Ala  
 1 5 10 15  
 Thr Gly Phe Pro Pro Lys Leu Ser Thr Pro Arg Ser Thr Ser Ala Ser  
 20 25 30  
 Gly Met Ser Ala Gly Ser Gln Arg Leu Trp Arg Arg Gly Ser Ser His  
 35 40 45  
 Ala Val Gln Thr Phe Asn Pro Leu Gln Ser Ser Leu Ala Arg Glu Gln  
 50 55 60  
 Gln Ser Leu Leu Glu Arg Asn Tyr His Ser Lys Gln Glu Phe Arg Pro  
 65 70 75 80  
 His Leu Ser Glu Asp His Val Glu Val His Leu Ala Gly Lys Val Ala  
 85 90 95  
 Ser Gly Cys Gly Leu Phe Asn Tyr Thr Leu Leu Phe Thr Leu Phe Thr  
 100 105 110  
 Ile Val Cys Lys Val Gln His Leu Gln Ala Arg Asn Thr Gly Leu Pro  
 115 120 125  
 His Ser Gly Trp Leu Gly Leu Met Lys Ala Ala Lys Gln Cys Ala Gln  
 130 135 140  
 Ser Lys Gln Arg Leu Pro Leu Ala Gly Ala His Ser Pro Arg Glu Gly  
 145 150 155 160

Ile Ser Phe Ser Leu Asp Leu Gly Ala Lys Ala Thr His Gly Ser Asp  
 165 170 175

Gln Thr Thr Cys  
 180

<210> 244  
 <211> 129  
 <212> PRT  
 <213> Homo sapiens

<400> 244

Val Glu Gln Leu Glu Thr His Gly Ser Val Leu Glu Trp Leu Val Trp  
 1 5 10 15

Asp His Phe Leu Gly Asp His Ser Ala Leu Thr Asp Gln Thr Gln Val  
 20 25 30

Asn Gly Thr Cys Pro Leu Pro Phe Pro Pro Gly Phe Gly Thr Val Ala  
 35 40 45

Thr Arg Val Val Phe Pro Ser Arg Gln Leu Leu Arg Val Ile Pro Glu  
 50 55 60

His Ser Leu Gly Ala Cys Ser Val Leu Thr Val Ile Ser Phe Ile Leu  
 65 70 75 80

Thr Ala Ile Pro Phe Cys Ile Phe Ser Gly His Pro Gln Asp His Pro  
 85 90 95

Gly Gln Pro Cys Leu Thr Pro Gly Leu Val Trp Leu His Asp Asn Lys  
 100 105 110

Asp Ala Gly Pro Glu Thr Ile Pro Leu His Gly Ala Cys Ile Phe Pro  
 115 120 125

Leu

<210> 245  
 <211> 181  
 <212> PRT  
 <213> Homo sapiens

<400> 245

Glu Ser Lys Met Leu Ile Gly Gly Ala Pro Pro Gln Cys Val Glu Asp  
 1 5 10 15

Leu Ala Ala Leu Asp Ala Tyr Ser Gln Ala Leu Gly Thr Arg Glu Ala  
 20 25 30

Pro Gly Leu Pro Phe Trp Ala Val Asp Leu Trp Gly Arg Ser Trp Pro  
 35 40 45

Leu Gly Trp Cys His Cys Ser Ser Tyr Pro Lys Cys Pro Phe Tyr Ala  
 50 55 60

Cys Ser Gly Leu Ala Ser Asn Thr Leu Lys Val Ser Ser Lys Gly Gln  
 65 70 75 80

Gly Arg Val Pro Cys Gly Lys Arg Trp Leu Phe Glu Ala Lys Ala Gln  
 85 90 95

Arg Arg His Ser Gln Arg Met Gly Arg Ala Ala Gly Gln Val Ser Ala  
 100 105 110  
 Ser Thr Trp Lys Thr Pro Ala Trp Leu Ala Ala Gly Glu Ile Val Leu  
 115 120 125  
 Pro Arg Cys Gln Leu Leu Ser Arg Pro Leu Pro Arg Glu Pro Ser His  
 130 135 140  
 Leu Ser Phe Ser Tyr Pro Ser Leu Arg Lys Ala Gln Ala Gln Gly Ala  
 145 150 155 160  
 Met Val Pro Cys Ser Gln Thr Val Ile Ser Glu Trp Pro Leu Val Trp  
 165 170 175  
 Gly Pro Arg Val Gln  
 180

<210> 246  
 <211> 137  
 <212> PRT  
 <213> Homo sapiens

<400> 246

Gln Asn Thr Phe Tyr His Ile Asn Ser Cys Thr Met Ile Trp Leu Glu  
 1 5 10 15  
 Glu Lys Asn Ser Trp Lys Val Lys Phe Val Leu Lys His Leu Phe Lys  
 20 25 30  
 Ser Leu His Thr Phe Ile Cys Pro Asp Lys Thr Cys Leu Asn Phe Phe  
 35 40 45  
 Leu Lys Gln Leu Tyr Cys Pro Ser Ile Cys Leu Thr Lys Phe Phe Lys  
 50 55 60  
 Gly His Phe Gln Pro Phe Gln Arg His Lys Val Gly Val Pro Lys Pro  
 65 70 75 80  
 Pro Phe Leu Ala Leu Pro Val Glu Asn Thr Met Leu His Ser Tyr Met  
 85 90 95  
 Cys Pro Leu Thr Gln Thr Thr Leu Ile Leu Arg Arg Ser Leu Asp Leu  
 100 105 110  
 Lys Leu Leu Leu Leu Ala Val Pro Ala Asn Ser Arg Val Lys Glu Asp  
 115 120 125  
 Val Thr Arg His Thr Tyr Leu Pro Phe  
 130 135

<210> 247  
 <211> 149  
 <212> PRT  
 <213> Homo sapiens

<400> 247

Ser Pro Met Leu Gln Phe Tyr Arg Leu Gly Lys Leu Arg Ala Gly Val  
 1 5 10 15  
 Thr Cys Tyr Ser Ser Tyr Pro Gln Thr Tyr Lys Thr Lys Ser Phe Thr  
 20 25 30

Glu Val Lys Tyr Asn Leu Phe Gly Leu Leu Phe His Phe Thr Ile Leu  
 35 40 45  
 Ser Leu Leu Val Phe Ile Thr Ile His Ser Lys Glu Phe Ile His Val  
 50 55 60  
 Asp Thr Ser Glu Val Phe Leu Ile Ser Pro Val Arg Pro Val Val Lys  
 65 70 75 80  
 Leu Leu Trp His Tyr Ser Thr Phe Ser Leu Ser Val Phe Phe Pro Ser  
 85 90 95  
 Pro His Arg Ser Glu Leu Ile Ser Pro His Pro Gly Pro Ser Glu Ser  
 100 105 110  
 Phe Val Lys Ser Leu Leu Ser Asn Leu Ser Val Glu Arg Val Pro Leu  
 115 120 125  
 Cys Leu Ser Glu Ile His Thr Val Met Cys His Leu Thr Met Phe Gln  
 130 135 140  
 Ser Val Arg Asp His  
 145

<210> 248  
 <211> 145  
 <212> PRT  
 <213> Homo sapiens

<400> 248

Pro Ile Pro Pro Ser Glu Gly Leu Glu Lys Ala Phe Thr Phe Met Ser  
 1 5 10 15  
 Pro Gly Ile Arg Ser Pro Gln Thr Arg Asn Phe Phe Leu Ile Met Glu  
 20 25 30  
 Val Trp Gln Trp Ala Thr Lys Pro Lys Val Ser Val Leu Leu Ser Asp  
 35 40 45  
 Ile Ala Ser Leu Arg Asn Arg Gln Pro Gly Arg Asp Gly Met Ser Leu  
 50 55 60  
 Ile Lys Cys Ser Ala Glu Val Ser Ser Arg Gly Leu Trp Cys Cys Pro  
 65 70 75 80  
 Ser Gly Cys Asn Ile Cys Thr Lys Pro Val Thr Glu Tyr Tyr Thr Glu  
 85 90 95  
 Ser Val Val Pro Lys Ile His Gly Phe Leu Tyr Gln Gly Leu Asp Ile  
 100 105 110  
 Glu Ser Ala Leu Val Thr Ile Lys Trp Leu Arg Asn Phe Tyr Phe Ile  
 115 120 125  
 Cys Pro Gln Leu Arg Trp Ile Arg Ser Val Cys Ile Leu Ala Ser Val  
 130 135 140

Cys  
 145

<210> 249  
 <211> 146  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 249

Leu Thr Ser Val Ser Ser Val Lys Pro Lys Leu Ser Lys Cys Glu Ile  
 1 5 10 15  
 Met Lys Cys Val Lys Leu Leu Ile Gln Cys Leu Arg Gln Gln Asn Ser  
 20 25 30  
 Arg Leu Ile Ile Gln Ser Ile Gln Thr Thr Phe Tyr Gly Asp Asn Leu  
 35 40 45  
 Trp Ser Glu Arg Leu His Lys Cys Ser Phe His Ser Tyr Ser Ser Ser  
 50 55 60  
 Asn Thr Lys Leu Leu Ser Ile Pro Glu Leu Lys Met Thr Leu Leu Thr  
 65 70 75 80  
 Asp Leu Tyr Leu Phe Ile Cys His Phe Ser Arg Arg Thr Ala Ile Leu  
 85 90 95  
 Pro Gln Ser Pro Tyr Ala Phe Val Glu Ser Trp Leu Lys Pro Gln Ala  
 100 105 110  
 Leu Cys Lys Ala Phe Leu Gly Ile Asp Ile Thr Thr Ile Pro Gln Asn  
 115 120 125  
 Leu Leu Val Leu His Ala Ile Ser Gly Pro Trp Thr His Phe Tyr Cys  
 130 135 140  
 Asn Lys  
 145

&lt;210&gt; 250

&lt;211&gt; 84

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 250

Phe Thr Gln Glu Ser Ser Arg Pro Ser Thr Phe Gly Ala Asn Leu Glu  
 1 5 10 15  
 Leu Gly Cys Arg Pro Ala Gly Thr Phe Ile Lys Cys Tyr Tyr Phe Ile  
 20 25 30  
 Phe Ala Ser Glu Glu Leu Pro Asp Phe Val Lys Thr Leu Cys Asn Pro  
 35 40 45  
 Ser Pro Phe Phe Trp His Ser Arg Gln Leu Asn Lys His Leu Leu Thr  
 50 55 60  
 Pro Leu Leu Cys Val Ile Arg Cys Glu Arg His Trp Arg Tyr Glu Glu  
 65 70 75 80  
 Pro Met Val Ser

&lt;210&gt; 251

&lt;211&gt; 62

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 251

Ala Pro Trp Gly Trp Ala Ser Val Ser Val Cys Ala Arg Leu Glu Met  
 1 5 10 15  
 Ala Ser Arg Tyr Gly Leu Gln Glu His His Glu Val His Leu Ile Phe  
 20 25 30  
 Ala Phe Leu Cys Gln His Val Cys His Leu Gln Cys Leu Thr Glu His  
 35 40 45  
 Val Gly Pro Ala Met Trp Ala Val Ser Leu Pro Ser Ser Tyr  
 50 55 60

<210> 252  
 <211> 117  
 <212> PRT  
 <213> Homo sapiens

<400> 252

Lys Lys Glu Pro Thr Met Ile Trp Ile Leu Leu Ser Phe Leu Phe Ser  
 1 5 10 15  
 Gln Met Val Ile Leu Lys Leu Ile Glu Val Val Tyr Arg Val His Ser  
 20 25 30  
 His Thr Val Arg Lys Arg Gln Ser Gln Gly Leu Asn Ser Ser Ser Leu  
 35 40 45  
 Thr Ile Glu Pro Ile Phe Leu Ile Thr Ile Gln Tyr Phe Pro Ile Cys  
 50 55 60  
 Ser Ile Lys Arg Asn His Phe Ser Glu Trp Arg Asn Ile His Glu Asn  
 65 70 75 80  
 Lys Ser Ile Ile Gln Asp Thr Cys Lys Ala Ser Arg His Ser Arg Phe  
 85 90 95  
 Arg Leu Leu Ala Pro Trp Pro Arg Leu Ile Thr Phe Gln Glu Asn Lys  
 100 105 110  
 Thr Thr Tyr Gln Asp  
 115

<210> 253  
 <211> 134  
 <212> PRT  
 <213> Homo sapiens

<400> 253

Thr Phe Ile Lys His Phe Phe Ser Gly Leu Ser Phe Ser Pro Ser Cys  
 1 5 10 15  
 His Val Ala Ile Ile Ile Phe Thr Ser Ala Ser Ala Tyr Phe Lys Pro  
 20 25 30  
 His Asn Lys Leu Leu Ala Phe Phe Phe Ala Ile Asp Asn Asn Leu Lys  
 35 40 45  
 Met Thr Gln Asn Phe Asn Gly Phe Ile Tyr Pro Gln Phe Tyr Asp Phe  
 50 55 60  
 Arg Ser Ser Phe Leu Cys Val Asp Leu Leu Ile Tyr His Phe Leu Ser  
 65 70 75 80



Thr Ile Thr Ser Phe Asn Leu Ser Cys Ser Thr Gly Leu Leu Thr Ile  
                   85                  90                  95  
 Asn Phe Phe Ser Phe Ser Leu Ser Lys Asn His Leu Phe Ser Leu His  
                   100                  105                  110  
 Phe Cys Lys Ile Phe Ser Arg Val Ile Lys Phe Val Thr Ile Phe Phe  
                   115                  120                  125  
 Glu Tyr Phe Lys Asp Leu  
                   130

<210> 254  
 <211> 138  
 <212> PRT  
 <213> Homo sapiens

<400> 254

Thr Phe Leu Ser Arg His Phe Leu Met Trp Lys Arg Phe Thr Glu Ser  
   1                  5                  10                  15  
 Asp Thr Phe Lys Gly Leu Thr Arg Asp Ile Cys Cys Leu Cys Leu Leu  
                   20                  25                  30  
 Phe Ser Trp Arg Ser Ala Thr Asn Lys Ala Ser Ser Thr Gln Gly His  
                   35                  40                  45  
 Leu Ser Thr Gly Leu Phe Leu Ser Ser Ser His Asn Leu Ser Cys His  
                   50                  55                  60  
 Thr Ile Thr Ser Thr Thr Ser Leu Gly Pro Cys Ser Glu Pro Thr Phe  
   65                  70                  75                  80  
 Phe Leu Pro Gln Val Gly Ile Ala Ser Ala Pro Tyr Cys Leu His Ser  
                   85                  90                  95  
 Glu Gly Ser Tyr Val His Ala Leu Asn Lys Phe Val Ser Pro Ile Asn  
                   100                  105                  110  
 Val Pro Phe Ala Ser Phe Phe Ser Glu Thr Ser Glu Val Gln Arg Gln  
                   115                  120                  125  
 Pro Leu Pro Ser Ser Arg Cys Ser Thr Tyr  
                   130                  135

<210> 255  
 <211> 155  
 <212> PRT  
 <213> Homo sapiens

<400> 255

Cys Lys Thr Gly Gly Leu Lys Leu Ile Phe Arg His His Gly Ile Leu  
   1                  5                  10                  15  
 Tyr Arg Leu Ser Leu Tyr Leu Glu Asp Val Arg Leu Met Glu Val Leu  
                   20                  25                  30  
 Ser Ile Leu Phe Pro Leu Leu Ile His Ser Phe Leu Phe Thr Glu Arg  
                   35                  40                  45  
 Leu Asn Phe Leu Ser His Ile Ser Val Leu Leu Ala Pro Leu Phe Phe  
                   50                  55                  60

Pro Leu Leu Gln Lys Ser Gln Pro Gln Lys Gln Ser Thr Tyr Cys Glu  
65 70 75 80

Lys Asp Phe Ser Asn His Lys Gly Asp Val Thr Leu Gly Leu Cys Phe  
85 90 95

Leu Ser His Thr His Lys Ile Leu Asp Met Ser Glu Ile Leu Lys Asn  
100 105 110

Trp Phe Leu Asn Val Met Lys Arg Val Ser Phe Ser Pro Glu Gln Asn  
115 120 125

Asn Pro Cys Ser Leu Leu Pro Asp Met Gly Gly Phe Gln Ile Arg Asn  
130 135 140

Leu Cys Ile Gly Pro Gln Ala Pro Asp Lys Val  
145 150 155

<210> 256

<211> 185

<212> PRT

<213> Homo sapiens

<400> 256

Gly His Arg Pro Ser Phe His Phe Cys Lys Pro Arg Gly Ile Leu Thr  
1 5 10 15

Asp Ser Thr Thr Tyr Pro Leu Leu Val Leu Ile Glu Glu Asp Thr Gly  
20 25 30

Leu Lys Pro His Phe Phe Arg Ala Phe Val Cys Ile Ser Lys Ile Leu  
35 40 45

Phe Tyr Arg His Leu Pro Phe Ser Phe Ile Phe Phe Leu Ser His Asn  
50 55 60

Asn Ser Ala Phe Leu Leu Tyr Glu Cys Thr Ser Asp Leu Thr Gln Arg  
65 70 75 80

Ile Gly Gly Gln Thr Asp Cys Leu Leu Ser Val Ser Cys Ala Leu Leu  
85 90 95

Arg Arg Leu His Leu Ser Ala Asn Ser Ser Cys Thr Thr Phe Ser Asp  
100 105 110

Phe Cys Cys Val Phe Ser Asp His Leu Leu Gly Ser Gly His Pro Leu  
115 120 125

Asp Gly Ser Gly Leu Ser Val Ser Val Phe Gly Asn Trp Ser Asp Leu  
130 135 140

Ala Leu Leu Met Gln Leu Lys Leu Arg Pro Leu Ser Leu Ser Gln Ala  
145 150 155 160

His Ser Gly Cys Val Arg Phe Leu Leu Ser Leu Val Cys Ile His Pro  
165 170 175

Leu His Val Gln Val Gly Ala Ala Lys  
180 185

<210> 257

<211> 128

<212> PRT

<213> Homo sapiens

&lt;400&gt; 257

His Phe Leu Pro His Ile Leu Glu Leu Val Leu Phe Leu Ile Lys Ile  
 1 5 10 15  
 Asn Val Ile Phe Arg Gly Ala Ile Phe Cys Phe Gln Asp Phe Phe Lys  
 20 25 30  
 Glu Val Ile Leu Lys Ala Lys Phe Lys Glu Lys Glu Leu Val Ala Leu  
 35 40 45  
 Val Asp Pro Val Gly Ser Ser Phe Leu Cys Trp Ser Ile Phe Cys Ile  
 50 55 60  
 Pro Phe Glu Phe Ala Phe Leu Phe Asn Ile Phe Trp Tyr Ser Arg Phe  
 65 70 75 80  
 Leu Phe Phe Gly Thr Phe Val His Ile Asn Phe Leu Val Trp Arg Arg  
 85 90 95  
 Gly Ile Leu Ile Ala Asn Gly Thr Lys Val Tyr Arg Asp Ile Val Gln  
 100 105 110  
 Pro Leu Leu Phe Phe Leu Phe Leu His Ser Ile Leu Val Met Gly Asn  
 115 120 125

&lt;210&gt; 258

&lt;211&gt; 168

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 258

Lys Gln Ser Tyr Ile Cys Ile Leu Phe Tyr Ile Tyr Phe Val Ile Phe  
 1 5 10 15  
 Leu Leu Ser Thr Val Ser Ser Leu Leu Pro Phe Leu Ile Glu Glu Phe  
 20 25 30  
 Asn Ala Cys Ile Cys Val Phe Ala Lys Lys Thr Pro Ser Ile Thr Cys  
 35 40 45  
 Ser Ile Tyr Glu Tyr Phe Trp Pro Leu Thr Gln Lys Val Leu Tyr Tyr  
 50 55 60  
 Arg Gln Lys Ser Thr Arg Lys Gln Ser Gly Thr Ser Ser Lys Arg Asp  
 65 70 75 80  
 Ser Ile Val Gly Lys Asn Thr Asp Pro Gly Gly Lys Leu Pro Gly Leu  
 85 90 95  
 Glu Ser Gln Leu Tyr Tyr Phe Gly Lys Thr Thr Tyr Leu Leu Tyr Leu  
 100 105 110  
 Phe Trp Tyr Pro Cys Leu Asn Gly Ser Asn Asn Asn Pro Leu Ile Ala  
 115 120 125  
 Leu Leu Gly Phe Asn Arg Ser Glu Asp Phe Arg Arg Ala His Asp Lys  
 130 135 140  
 Asn Tyr Ile Arg Val Thr Tyr Tyr Cys Tyr Pro Ile Cys His Ser Lys  
 145 150 155 160  
 Leu Arg Asp Leu Gly Gln Val Thr

165

<210> 259  
 <211> 182  
 <212> PRT  
 <213> Homo sapiens

<400> 259

Leu Val Glu Trp Ala His Ser Ser Met Arg Pro Ile Phe His Leu Asn  
 1 5 10 15  
 Phe Leu Cys Leu Arg Asn Glu Leu Tyr Ser Asn Leu Cys Phe Leu Lys  
 20 25 30  
 Ile Asn Val Phe Leu Val Lys His Leu Val Ser Ser Gln Ile Leu Phe  
 35 40 45  
 Lys Lys Thr Thr Glu Asn Ser Glu Glu Gly Glu Thr Asp Ser Ala Asn  
 50 55 60  
 Ser Ile Ser Val Pro Arg Leu Asn Trp Glu Met Leu Leu Leu His Asp  
 65 70 75 80  
 Leu Gly Leu Ile Ile Cys Leu Gln Glu His Cys Phe Arg Val Val Trp  
 85 90 95  
 Tyr Ser Gly Arg Asn Gly Leu Trp Ser Glu Ile His Val Gln Ile Pro  
 100 105 110  
 Ser His Leu Pro Ser Leu Ile Leu Ser Phe Leu Ile Cys Lys Met Thr  
 115 120 125  
 Ile Ile Asn Thr Ile Ser Lys Ile Cys Gly Asp Asn Thr Ala Phe Thr  
 130 135 140  
 Ser Cys Cys Ile Leu Pro Ile Ser Ser Cys Arg Asp Arg Ile Phe His  
 145 150 155 160  
 Phe Ile Leu Ile Tyr Asn Tyr Val Ile Pro Phe Lys Asn His Pro Ser  
 165 170 175  
 Thr Phe Ser Ser Thr Arg  
 180

<210> 260  
 <211> 207  
 <212> PRT  
 <213> Homo sapiens

<400> 260

Cys Ser Leu Leu Asp Phe Leu Met Leu Val Gly Ala Leu Arg Lys Leu  
 1 5 10 15  
 Cys Thr Lys Leu Asp Pro Val Leu Gln Gly Ser Asp Leu Thr Glu His  
 20 25 30  
 Ser Ala Trp Gly Val Pro Leu Ile Trp Thr Trp Asn Ser Ile Ile Gln  
 35 40 45  
 Arg Pro Ser Leu Pro Cys Ser Leu Cys Val Thr Gly Ala Ala Glu Thr  
 50 55 60  
 Gln Val Leu Ser Ala Ser Ala Gly Leu Gln Pro Cys Leu Cys Leu Leu

65                      70                      75                      80  
 Arg Ser Asp Ser Asn Cys Tyr Leu Trp Arg Trp Leu Phe Ile Gly Thr  
                                  85                      90                      95  
 Pro Phe Leu Cys Leu Thr Glu Ala Gln Cys Ser Lys Leu Glu Gly Leu  
                                  100                      105                      110  
 Cys Gln His Val Ser His Thr His Leu Leu Leu Phe Phe Ser Arg Val  
                                  115                      120                      125  
 Leu Gly His Leu Leu Leu His Ile Thr Thr Ser Ser Pro Pro Ala Gln  
                                  130                      135                      140  
 Leu Ala Leu Ser Pro Phe Pro Ile Tyr His Ala Val Leu Glu His Lys  
                                  145                      150                      155                      160  
 Ala Leu Leu Cys Ile Pro Cys Val Tyr Phe Val Val Met Cys Cys Ile  
                                  165                      170                      175  
 Leu Lys Glu Leu Asn Leu Cys Pro Gly Ser Arg Lys Asn Ala Asp Gln  
                                  180                      185                      190  
 Leu Leu Ala Ile Asp Gly Phe Asn Ile Ser Tyr Asp Trp Phe Leu  
                                  195                      200                      205  
  
 <210> 261  
 <211> 187  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 261  
  
 Gln Thr Lys Glu Glu Lys Gly Gln Val Lys His Thr Ile Gly Phe Thr  
 1                      5                      10                      15  
 Val Asn Met Ser Lys Val Leu Leu Ile Ile His Phe Met Tyr Pro Arg  
                                  20                      25                      30  
 Leu Trp Lys Lys Phe Phe Phe His Leu Pro Ile Lys Asn Ile His Leu  
                                  35                      40                      45  
 Gly Ile Thr Thr Ser Trp Ile Leu Leu Asp Arg His Thr Thr Thr Leu  
                                  50                      55                      60  
 Thr Val Leu Pro Ser Ser Arg Arg Leu Ala Arg Lys Ala His His Pro  
 65                      70                      75                      80  
 Leu Pro Gly Ser Lys Val Asp Ser Leu Ile Phe Cys Ile Asn Pro Thr  
                                  85                      90                      95  
 Pro Asp Ser Phe Ser Tyr Ser Leu Leu Pro Cys Leu Phe Ser Tyr Leu  
                                  100                      105                      110  
 Met Val Asn Val Phe Leu Ser Ser Cys Ile Thr Phe Tyr Ser Phe Leu  
                                  115                      120                      125  
 Glu His Ile Ile Ile Ile Asn Lys Lys Ser Lys Ile Ala Met Val Ala  
                                  130                      135                      140  
 Arg Ile Pro Ala Pro Leu Asp Pro Ser Thr Ser Ser Ser Pro Gly His  
 145                      150                      155                      160  
 Thr Trp Gln Arg Glu Ile Lys Val Leu Asp Gly Ile Lys Val Asn Gln  
                                  165                      170                      175

Leu Thr Leu Lys Gly Glu Lys Glu Ser Arg Leu  
180 185

<210> 262  
<211> 149  
<212> PRT  
<213> Homo sapiens

<400> 262

Tyr Val Thr Ile Leu Leu Thr Val Leu Val Phe Leu Leu Arg Ser Leu  
1 5 10 15

Pro Phe Gly Ile Arg Trp Ala Leu Ser Thr Gly Ile His Leu Asp Leu  
20 25 30

Glu Val Ile Phe Cys His Val His Leu Val Ser Ile Phe Leu Ser Pro  
35 40 45

Leu Asn Gly Ser Ala Asn Pro Val Ile Tyr Phe Phe Val Gly Ser Phe  
50 55 60

Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys Leu Val Leu Gln Arg Ala  
65 70 75 80

Leu Gln Asp Met Pro Glu Val Lys Val Glu Gly Gly Phe Leu Arg Glu  
85 90 95

Pro Trp Ser Cys Arg Glu Ala Asp Ser Gly Ser Glu Glu Glu Pro Leu  
100 105 110

Pro Cys Gln Ser Asp Gly Thr Leu Arg Ala Ile Leu Pro Cys His Ala  
115 120 125

Gln Leu His Ala Phe Ser Cys Cys Ala Ser Glu Met Ser Gln Arg Leu  
130 135 140

Lys Val Val Glu Met  
145

<210> 263  
<211> 207  
<212> PRT  
<213> Homo sapiens

<400> 263

His Trp Arg Ser Leu Val Thr Trp Ala Glu Tyr Leu Glu Pro Arg Ile  
1 5 10 15

Ser Ser Ser Met Val Asp Gln Leu Cys Asp Gly Val Met Arg Trp Gly  
20 25 30

Arg Arg Val Trp His His Ala Thr Gly Phe Pro Pro Lys Leu Ser Thr  
35 40 45

Pro Arg Ser Thr Ser Ala Ser Gly Met Ser Ala Gly Ser Gln Arg Leu  
50 55 60

Trp Arg Arg Gly Ser Ser His Ala Val Gln Ser Phe Asn Pro Leu Gln  
65 70 75 80

Ser Ser Leu Ala Arg Glu Gln Gln Ser Leu Leu Glu Arg Asn Tyr His  
85 90 95

Ser Lys Gln Glu Phe Arg Pro His Leu Ser Glu Asp His Val Glu Val  
 100 105 110  
 His Leu Ala Gly Lys Val Ala Ser Gly Cys Gly Leu Phe Asn Tyr Thr  
 115 120 125  
 Leu Leu Phe Thr Leu Phe Thr Ile Val Cys Lys Val Gln His Leu Gln  
 130 135 140  
 Ala Arg Asn Thr Gly Leu Pro His Ser Gly Trp Leu Gly Leu Met Lys  
 145 150 155 160  
 Ala Thr Lys Gln Cys Ala Gln Ser Lys Gln Arg Leu Pro Leu Ala Gly  
 165 170 175  
 Ala His Ser Pro Arg Glu Gly Ile Ser Phe Ser Leu Asp Leu Gly Ala  
 180 185 190  
 Lys Ala Thr His Gly Ser Asp Gln Thr Thr Cys Ser Pro His Leu  
 195 200 205

<210> 264  
 <211> 204  
 <212> PRT  
 <213> Homo sapiens  
 <400> 264

Gly Ala Ser Ser Gln Tyr Gly Asn Glu Asp Gly Val Asn Leu Phe Pro  
 1 5 10 15  
 Leu Met Ser Pro Pro Leu Tyr Thr Asn Leu Leu Lys Pro Thr Gly Lys  
 20 25 30  
 Leu Arg Leu Gly Asn Lys Asn Ile Lys Cys Tyr Val Gln Ile Leu Lys  
 35 40 45  
 Trp Asn Leu Lys Leu Leu Val Leu Gln Leu Phe Leu Lys Ile Pro Thr  
 50 55 60  
 Leu Ser Arg Ser Met Ser Phe Arg Glu Arg Thr Tyr Val Ala Arg Glu  
 65 70 75 80  
 Lys Ser Lys Glu Ser Met Asn Pro Val Leu Leu Ser Ile Leu Gln Cys  
 85 90 95  
 Trp Arg Pro Phe Ser Ile Phe His Ser Leu Gly Gln Ser Phe Asn Thr  
 100 105 110  
 His Leu Leu Lys Ala Ile Tyr Ile Arg Pro Cys Tyr Ser Lys Gly Thr  
 115 120 125  
 Val Gly Gly Glu Glu Arg Gln Asp Pro Thr Met Glu Leu Lys Ser Ser  
 130 135 140  
 Leu Asp Arg Phe Pro Phe Pro Ser Gly Gln Ser Lys Pro Asn Asp Thr  
 145 150 155 160  
 Thr Val Ser Ser Phe Pro Glu Gln Arg Asp Val Glu Asn Tyr Leu Phe  
 165 170 175  
 Thr Ile Val Arg Arg Arg Gln Gly Trp Asn Phe Phe Gln Asn Lys Leu  
 180 185 190

Phe Phe Phe Val Lys Gln Gly Lys Ile Leu Leu Leu  
195 200

<210> 265  
<211> 186  
<212> PRT  
<213> Homo sapiens

<400> 265

Ile Ser Val Thr Asp Leu Ile Gly Gly Lys Trp Ile Phe Gly His Phe  
1 5 10 15  
Phe Cys Asn Val Phe Ser Val Asn Val Met Cys Cys Thr Ala Trp Ile  
20 25 30  
Leu Thr Leu Tyr Val Ile Ser Ile Asp Arg Tyr Leu Gly Ile Met Lys  
35 40 45  
Pro Leu Thr Tyr Pro Met Arg Gln Lys Gly Lys Cys Met Thr Lys Met  
50 55 60  
Ile Leu Ser Val Cys Leu Leu Ser Ala Phe Val Thr Leu Pro Thr Ile  
65 70 75 80  
Phe Gly Arg Ala Gln Asn Val Asn Asp Asp Lys Val Cys Leu Val Ser  
85 90 95  
Gln Asp Phe Gly Tyr Thr Ile Tyr Ser Thr Ala Leu Ala Ser Ser Pro  
100 105 110  
Cys Ala Ser Cys Phe Ser Cys Thr Asn Arg Phe Thr Arg Pro Pro Gly  
115 120 125  
Lys Ala Arg Pro Asn Thr Gly Tyr Leu Ala Ser Leu Glu Trp Ser Gln  
130 135 140  
Thr Ala Val Val Thr Leu Asn Gly Thr Val Lys Phe Gln Glu Val Glu  
145 150 155 160  
Glu Cys Ala Lys Leu Ser Arg Leu Leu Lys His Glu Arg Lys Lys Tyr  
165 170 175  
Leu His Leu Ala Glu Thr Glu Ser Ser Asp  
180 185

<210> 266  
<211> 184  
<212> PRT  
<213> Homo sapiens

<400> 266

Phe Thr Val Ile Asn Val Cys Ser Cys Thr Cys Glu Val Lys Ser Phe  
1 5 10 15  
Ser Leu Leu Ser Asn Ser Tyr Val Pro Asn Ile Phe Ser Lys Phe Leu  
20 25 30  
Lys Thr Tyr Asn Gly Glu Lys Asn Asn Pro Phe Ser Ser Pro Ala Ser  
35 40 45  
Leu Met Lys Asn Ser His Phe Ser Leu Phe Leu Leu Phe Leu Leu Val  
50 55 60



Val Phe His Ile Ser Cys Leu Ser Ala Val Ser Cys Phe Met Gln Phe  
65 70 75 80  
Arg Pro Tyr Leu Leu Thr Ser Leu Ser Phe Gln Tyr Lys Asp Ser Cys  
85 90 95  
Ile Phe Ser Phe Asn Phe Thr Phe Leu Asn Ser Pro Phe Pro Phe Cys  
100 105 110  
Asp Pro Gly Ile Ser Gly Val Leu Phe Phe Phe Ile Leu Pro Asp Phe  
115 120 125  
Ile Tyr Ile Cys Val Tyr Ser Phe Leu Leu Phe Phe Lys Leu Lys Thr  
130 135 140  
Cys Leu Ser Ser Lys Ser Gly Ser Phe Phe Phe Ser Trp Arg Pro Leu  
145 150 155 160  
Ser Gln Asn Pro Leu Ser Phe Cys Phe Asn Glu Asp Tyr Met Leu Ser  
165 170 175  
Leu Trp Leu Pro Ser Cys Asn Thr  
180

<210> 267  
<211> 201  
<212> PRT  
<213> Homo sapiens

<400> 267

Phe Pro Ser Leu Lys Asn Met His Phe Ser Val Pro Leu Arg Cys His  
1 5 10 15  
Thr Ile Ile Ser Val Gln Lys Arg Val Asn Thr Ala Asp Pro Arg Leu  
20 25 30  
Leu Leu Leu Lys Cys Pro Ala Cys Lys Ala Gly Ser Trp Leu Val Phe  
35 40 45  
Gly Val Leu Asp Phe Glu Lys Leu Pro Thr Ile Pro Ser Thr Gly Leu  
50 55 60  
Cys Lys Tyr Gly Leu Tyr Ile Pro Ala Phe Leu Leu Glu Leu Glu Phe  
65 70 75 80  
Ser Lys Tyr Glu Ala Lys Arg Ala Tyr Val Thr Ser Pro Gln Pro Trp  
85 90 95  
Ala Leu Ser His Gly Thr Ser Leu Ala Gly Ser Val Ser His Val Leu  
100 105 110  
Ser Gln Phe Leu Ala Glu Arg Ile Lys His Ile Leu Cys Asn Phe Thr  
115 120 125  
Gly Lys Arg Ile Leu Glu Ala Val Pro Gly Phe Phe Arg Leu Phe Leu  
130 135 140  
Met His Leu Phe Leu Leu Leu Ile Met Leu Arg Tyr Pro Ser Val Asn  
145 150 155 160  
Lys Ser Leu Ile Gln Leu Tyr Ala Lys Ser Tyr Glu Ser Gln Asn Arg  
165 170 175  
Gly Ile Ile Leu Gly Arg Pro Asp Thr Thr Lys Ile Asn Leu Lys Leu

180                      185                      190  
 Asn Ser Ser Pro Thr Ser Leu Ser Pro  
     195                      200  
 <210> 268  
 <211> 321  
 <212> PRT  
 <213> Homo sapiens  
 <400> 268  
 Met Asn Gln Thr Leu Asn Ser Ser Gly Thr Val Glu Ser Ala Leu Asn  
 1                      5                      10                      15  
 Tyr Ser Arg Gly Ser Thr Val His Thr Ala Tyr Leu Val Leu Ser Ser  
                     20                      25                      30  
 Leu Ala Met Phe Thr Cys Leu Cys Gly Met Ala Gly Asn Ser Met Val  
                     35                      40                      45  
 Ile Trp Leu Leu Gly Phe Arg Met His Arg Asn Pro Phe Cys Ile Tyr  
 50                      55                      60  
 Ile Leu Asn Leu Ala Ala Ala Asp Leu Leu Phe Leu Phe Ser Met Ala  
 65                      70                      75                      80  
 Ser Thr Leu Ser Leu Glu Thr Gln Pro Leu Val Asn Thr Thr Asp Lys  
                     85                      90                      95  
 Val His Glu Leu Met Lys Arg Leu Met Tyr Phe Ala Tyr Thr Val Gly  
                     100                      105                      110  
 Leu Ser Leu Leu Thr Ala Ile Ser Thr Gln Arg Cys Leu Ser Val Leu  
                     115                      120                      125  
 Phe Pro Ile Trp Phe Lys Cys His Arg Pro Arg His Leu Ser Ala Trp  
 130                      135                      140  
 Val Cys Gly Leu Leu Trp Thr Leu Cys Leu Leu Met Asn Gly Leu Thr  
 145                      150                      155                      160  
 Ser Ser Phe Cys Ser Lys Phe Leu Lys Phe Asn Glu Asp Arg Cys Phe  
                     165                      170                      175  
 Arg Val Asp Met Val Gln Ala Ala Leu Ile Met Gly Val Leu Thr Pro  
                     180                      185                      190  
 Val Met Thr Leu Ser Ser Leu Thr Leu Phe Val Trp Val Arg Arg Ser  
                     195                      200                      205  
 Ser Gln Gln Trp Arg Arg Gln Pro Thr Arg Leu Phe Val Val Val Leu  
 210                      215                      220  
 Ala Ser Val Leu Val Phe Leu Ile Cys Ser Leu Pro Leu Ser Ile Tyr  
 225                      230                      235                      240  
 Trp Phe Val Leu Tyr Trp Leu Ser Leu Pro Pro Glu Met Gln Val Leu  
                     245                      250                      255  
 Cys Phe Ser Leu Ser Arg Leu Ser Ser Ser Val Ser Ser Ser Ala Asn  
                     260                      265                      270  
 Pro Val Ile Tyr Phe Leu Val Gly Ser Arg Arg Ser His Arg Leu Pro  
                     275                      280                      285

Thr Arg Ser Leu Gly Thr Val Leu Gln Gln Ala Leu Arg Glu Glu Pro  
290 295 300

Glu Leu Glu Gly Gly Glu Thr Pro Thr Val Gly Thr Asn Glu Met Gly  
305 310 315 320

Ala

<210> 269  
<211> 9  
<212> PRT  
<213> Artificial

<220>  
<223> Novel Sequence

<400> 269

Ala Pro Arg Thr Pro Gly Gly Arg Arg  
1 5

<210> 270  
<211> 20  
<212> DNA  
<213> Artificial

<220>  
<223> Novel Sequence

<400> 270  
ctgtctctct gtcctcctcc 20

<210> 271  
<211> 22  
<212> DNA  
<213> Artificial

<220>  
<223> Novel Sequence

<400> 271  
gcaccgatct tcattgaatt tc 22

<210> 272  
<211> 33  
<212> DNA  
<213> Artificial

<220>  
<223> Novel Sequence

<400> 272  
gatcaagctt ggatgaacca gactttgaat agc 33

<210> 273  
<211> 31  
<212> DNA  
<213> Artificial

<220>

<223> Novel Sequence

<400> 273

gacctcgag ctcaagcccc catctcattg g

31